



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C12Q 1/100, G01N 33/53, 33/48, C12P 21/06, C12N 15/00, C07H 21/02, C07K 16/00, 1/00, A61K 39/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/39327</p> <p>(43) International Publication Date: 6 July 2000 (06.07.00)</p>
<p>(21) International Application Number: PCT/US99/30736</p> <p>(22) International Filing Date: 22 December 1999 (22.12.99)</p> <p>(30) Priority Data: 60/113,809 23 December 1998 (23.12.98) US</p> <p>(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Olney, MD 20832 (US). YOUNG, Paul, E. [US/US]; 122 Beckwith Street, Gaithersburg, MD 20878 (US). OLSEN, Henrik, S. [DK/US]; 182 Kendrick Place, #24, Gaithersburg, MD 20878 (US).</p> <p>(74) Agents: HOOVER, Kenley, K. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).</p>		
<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>		
<p>(54) Title: PEPTIDOGLYCAN RECOGNITION PROTEINS</p> <p>(57) Abstract</p> <p>The present invention relates to three novel peptidoglycan recognition binding proteins expressed by keratinocytes, wound-healing tissues and chondrosarcoma tissue. More specifically, isolated nucleic acid molecules are provided encoding human peptidoglycan recognition protein-related proteins, referred to herein as PGRP-K (Keratinocytes), PGRP-W (Wound-healing), and PGRP-C (Chondrosarcoma) of Figures 1A-B, Figures 2A-C, and Figure 3, respectively, each having homology to both human peptidoglycan recognition protein (PGRP) as well as murine Tag-7. PGRP-K, PGRP-W, and PGRP-C polypeptides are also provided. Further provided are vectors, host cells and recombinant methods for producing the same. The invention also relates to both the inhibition and enhancement of activities of PGRP-K, PGRP-W, and PGRP-C polypeptides and diagnostic methods for detecting PGRP-K, PGRP-W, and PGRP-C gene expression.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Peptidoglycan Recognition Proteins

Field of the Invention

The present invention relates to three novel peptidoglycan recognition binding
5 proteins expressed by keratinocytes, wound-healing tissues and chondrosarcoma
tissue. More specifically, isolated nucleic acid molecules are provided encoding
human peptidoglycan recognition protein-related proteins, referred to herein as
PGRP-K (Keratinocytes), PGRP-W (Wound-healing), and PGRP-C
(Chondrosarcoma) of Figures 1A-B, Figures 2A-C, and Figure 3, respectively, each
10 having homology to both human peptidoglycan recognition protein (PGRP) as well as
murine Tag-7. PGRP-K, PGRP-W, and PGRP-C polypeptides are also provided.
Further provided are vectors, host cells and recombinant methods for producing the
same. The invention also relates to both the inhibition and enhancement of activities of
PGRP-K, PGRP-W, and PGRP-C polypeptides and diagnostic methods for detecting
15 PGRP-K, PGRP-W, and PGRP-C gene expression.

Background of the Invention

Peptidoglycan, as well as Lipopolysaccharide (LPS), is a surface component of
20 many bacteria which illicit a wide range of physiological and immune responses in
humans. Specifically, peptidoglycan has been shown to manifest itself clinically by
reproducing most of the symptoms of bacterial infection, including fever, acute-phase
response, inflammation, septic shock, leukocytosis, sleepiness, malaise, abcess
formation, and arthritis (see Dziarski *et al.*, JBC, 273 (15): 8680 (1998)). Furthermore,
25 the type of peptidoglycan (i.e.- the specific stereoisomers or analogs of muramyl
dipeptide, N-acetylglucosaminyl-beta(1-4)-N-acetylmuramyl tetrapeptides, etc.),
were shown to elicit a broad range of activities, including exhibiting greater
pyrogenicity, inducing acute joint inflammation, stimulating macrophages, and causing

hemorrhagic necrosis at a primed site (See Kotani *et al.*, Fed Proc, 45(11): 2534 (1986)).

It has been demonstrated in humans that a lipopolysaccharide binding protein exists that was discovered as a trace plasma protein (See Schumann *et al.*, Science, 5 249(4975):1429 (1990)). It is thought that one of the modes of action by which this lipopolysaccharide binding protein functions is by forming high-affinity complexes with lipopolysaccharide, that then bind to macrophages and monocytes, inducing the secretion of tumor necrosis factor. Dziarski and Gupta (See Dziarski *et al.*, JBC, 269(3): 2100 (1994)) demonstrated that a 70kDa receptor protein present on the 10 surface of mouse lymphocytes served to bind heparin, heparinoids, bacterial lipoteichoic acids, peptidoglycan, and lipopolysaccharides.

Recently, Dziarski *et al.* demonstrated that the CD14, a glycosylphosphatidylinositol-linked protein present on the surface of macrophage and polymorphonuclear leukocytes, bound peptidoglycan and lipopolysaccharide. 15 Furthermore, the binding affinity of CD14 for lipopolysaccharide was significantly increased in the presence of a LPS-binding protein present in plasma. It is thought that the LPS-binding protein functions as a transfer molecule, whereby it binds LPS and presents it to the CD14 receptor (See Dziarski *et al.*, JBC, 273(15): 8680 (1998)).

Yoshida *et al.* isolated a peptidoglycan binding protein from the hemolymph of 20 the Silkworm, *Bombyx mori*, using column chromatography. This protein was found to have a very specific affinity for peptidoglycan (See Yoshida *et al.*, JBC, 271(23): 13854 (1996)). Additionally, Kang *et al.* recently cloned a peptidoglycan binding protein from the moth *Trichoplusia ni*. The peptidoglycan binding protein was shown to bind strongly to insoluble peptidoglycan (See Kang *et al.*, PNAS, 95(17): 10078 25 (1998)). In this study the peptidoglycan binding protein was upregulated by a bacterial infection in *T. ni*. The insect immune system is regarded as a model for innate immunity. Thus, Kang *et al.* were able to clone both mouse and human homologs of the *T. ni* peptidoglycan binding protein. All of these peptidoglycan binding proteins shared regions of homology, as well as four conserved cysteine

residues which may function in the tertiary structure of the protein, possibly in helping to form binding domains. Given that peptidoglycan is an integral component of bacterial cell walls, and that it induces many physiological responses from cytokine secretion to inflammation and macrophage activation, it appears as if this family of proteins may be a ubiquitous group involved in the binding and recognition of peptidoglycan, the presentation of antigens (e.g., cell wall components, etc.), and the activation of the immune system, such as the secretion of cytokines, such as TNF.

TNF is noted for its pro-inflammatory actions which result in tissue injury, such as induction of procoagulant activity on vascular endothelial cells (Pober, J.S. *et al.*, *J. Immunol.* 136:1680 (1986)), increased adherence of neutrophils and lymphocytes (Pober, J.S. *et al.*, *J. Immunol.* 138:3319 (1987)), and stimulation of the release of platelet activating factor from macrophages, neutrophils and vascular endothelial cells (Camussi, G. *et al.*, *J. Exp. Med.* 166:1390 (1987)).

Recent evidence implicates TNF in the pathogenesis of many infections (Cerami, A. *et al.*, *Immunol. Today* 9:28 (1988)), immune disorders, neoplastic pathology, e.g., in cachexia accompanying some malignancies (Oloff, A. *et al.*, *Cell* 50:555 (1987)), and in autoimmune pathologies and graft-versus host pathology (Piguet, P.-F. *et al.*, *J. Exp. Med.* 166:1280 (1987)). The association of TNF with cancer and infectious pathologies is often related to the host's catabolic state. A major problem in cancer patients is weight loss, usually associated with anorexia. The extensive wasting which results is known as "cachexia" (Kern, K. A. *et al.* *J. Parent. Enter. Nutr.* 12:286-298 (1988)). Cachexia includes progressive weight loss, anorexia, and persistent erosion of body mass in response to a malignant growth. The cachectic state is thus associated with significant morbidity and is responsible for the majority of cancer mortality. A number of studies have suggested that TNF is an important mediator of the cachexia in cancer, infectious pathology, and in other catabolic states.

TNF is thought to play a central role in the pathophysiological consequences of Gram-negative sepsis and endotoxic shock (Michie, H.R. *et al.*, *Br. J. Surg.* 76:670-671 (1989); Debets, J. M. H. *et al.*, *Second Vienna Shock Forum*, p.463-466

(1989); Simpson, S. Q. *et al.*, *Crit. Care Clin.* 5:27-47 (1989)), including fever, malaise, anorexia, and cachexia. Endotoxin is a potent monocyte/macrophage activator which stimulates production and secretion of TNF (Kornbluth, S.K. *et al.*, *J. Immunol.* 137:2585-2591 (1986)) and other cytokines. Because TNF could mimic
5 many biological effects of endotoxin, it was concluded to be a central mediator responsible for the clinical manifestations of endotoxin-related illness. TNF and other monocyte-derived cytokines mediate the metabolic and neurohormonal responses to endotoxin (Michie, H.R. *et al.*, *N. Eng. J. Med.* 318:1481-1486 (1988)). Endotoxin administration to human volunteers produces acute illness with flu-like symptoms
10 including fever, tachycardia, increased metabolic rate and stress hormone release (Revhaug, A. *et al.*, *Arch. Surg.* 123:162-170 (1988)). Elevated levels of circulating TNF have also been found in patients suffering from Gram-negative sepsis (Waage, A. *et al.*, *Lancet* 1:355-357 (1987); Hammerle, A.F. *et al.*, *Second Vienna Shock Forum* p. 715-718 (1989); Debets, J. M. H. *et al.*, *Crit. Care Med.* 17:489-497 (1989); Calandra,
15 T. *et al.*, *J. Infec. Dis.* 161:982-987 (1990)).

Passive immunotherapy directed at neutralizing TNF may have a beneficial effect in Gram-negative sepsis and endotoxemia, based on the increased TNF production and elevated TNF levels in these pathology states, as discussed above. Antibodies to a "modulator" material which was characterized as cachectin (later
20 found to be identical to TNF) were disclosed by Cerami *et al.* (EPO Patent Publication 0,212,489, March 4, 1987). Such antibodies were said to be useful in diagnostic immunoassays and in therapy of shock in bacterial infections. Rubin *et al.* (EPO Patent Publication 0,218,868, April 22, 1987) disclosed monoclonal antibodies to human TNF, the hybridomas secreting such antibodies, methods of producing such
25 antibodies, and the use of such antibodies in immunoassay of TNF. Yone *et al.* (EPO Patent Publication 0,288,088, October 26, 1988) disclosed anti-TNF antibodies, including mAbs, and their utility in immunoassay diagnosis of pathologies, in particular Kawasaki's pathology and bacterial infection. The body fluids of patients with Kawasaki's pathology (infantile acute febrile mucocutaneous lymph node

syndrome; Kawasaki, T., *Allergy* 16:178 (1967); Kawasaki, T., *Shonika (Pediatrics)* 26:935 (1985)) were said to contain elevated TNF levels which were related to progress of the pathology (Yone *et al.*, *supra*).

Accordingly, there is a need to provide molecules that are involved in pathological conditions. Such novel proteins could be useful in augmenting the immune system in such areas as immune recognition, antigen presentation, and immune system activation. Antibodies or antagonists directed against these proteins may be useful in reducing or eliminating disorders associated with TNF and TNF-like cytokines, such as endotoxic shock and auto-immune disorders, for example.

Summary of the Invention

The present invention provides isolated nucleic acid molecules comprising polynucleotides encoding three novel proteins that are structurally similar to a human Peptidoglycan Recognition Protein and murine Tag-7, and are believed to have similar biological effects and activities. The cytokines are named PGRP-K, PGRP-W, and PGRP-C, and the invention includes PGRP-K, PGRP-W, and PGRP-C polypeptides having at least a portion of the amino acid sequence in Figures 1A-B (SEQ ID NO:2), in Figures 2A-C (SEQ ID NO:4), and/or Figure 3 (SEQ ID NO:6) or amino acid sequence encoded by the cDNA clones deposited on December 23, 1998, assigned ATCC number 203564; December 23, 1998, assigned ATCC number 203563; and March 20, 1998, assigned ATCC number 209683, respectively. The nucleotide sequence determined by sequencing the deposited PGRP-K clone, which is shown in Figures 1A-B (SEQ ID NO:1), contains an open reading frame encoding a complete polypeptide of 243 amino acid residues including an N-terminal methionine, a predicted PGRP-like domain of about 83 amino acid residues, and a deduced molecular weight for the complete protein of about 27 kDa.

The nucleotide sequence determined by sequencing the deposited PGRP-W clone, which is shown in Figures 2A-C (SEQ ID NO:3), contains an open reading

frame encoding a complete polypeptide of 368 amino acid residues including an N-terminal methionine, a predicted PGRP-like domain of about 83 amino acid residues, and a deduced molecular weight for the complete protein of about 40 kDa.

5 The nucleotide sequence determined by sequencing the deposited PGRP-C clone, which is shown in Figure 3 (SEQ ID NO:5), contains an open reading frame encoding a complete polypeptide of 196 amino acid residues including an N-terminal methionine, a predicted PGRP-like domain of about 83 amino acid residues, and a deduced molecular weight for the complete protein of about 21 kDa.

Thus, one aspect of the invention provides isolated nucleic acid molecules
10 comprising polynucleotides having nucleotide sequences selected from the group consisting of: (a) a nucleotide sequence encoding a full-length PGRP-K, PGRP-W, or PGRP-C polypeptide having the complete amino acid sequence in Figures 1A-B (SEQ ID NO:2), in Figures 2A-C (SEQ ID NO:4), or in Figure 3 (SEQ ID NO:6), respectively, or as encoded by the cDNA clones contained in the ATCC Deposit
15 number 203564, deposited on December 23, 1998; ATCC Deposit number 203563, deposited on December 23, 1998; and ATCC Deposit number 209683, deposited March 20, 1998, respectively.; (b) a nucleotide sequence encoding the predicted PGRP-like domain of the PGRP-K polypeptide having the amino acid sequence at positions 24 to 107 in Figures 1A-B (SEQ ID NO:2), the predicted PGRP-like domain
20 of the PGRP-W polypeptide having the amino acid sequence at positions 52 to 135 in Figures 2A-C (SEQ ID NO:4), and/or the predicted PGRP-like domain of the PGRP-C polypeptide having the amino acid sequence at positions 34 to 117 in Figure 3 (SEQ ID NO:6), or as encoded by the cDNA clones contained in ATCC Numbers 203564, 203563, and 209683, respectively, deposited on December 23, 1998, and March 20,
25 1998; or as encoded by the cDNA clones contained in ATCC Numbers 203564, 203563, and 209683, respectively, deposited on December 23, 1998, and March 20, 1998; (c) a nucleotide sequence encoding a soluble PGRP-K, PGRP-W, and/or PGRP-C polypeptide having the PGRP-like domain but lacking the leader sequence; and (d) a

nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), or or (c) above.

Further embodiments of the invention include isolated nucleic acid molecules that comprise a polynucleotide having a nucleotide sequence at least 90% identical, and more preferably at least 95%, 96%, 97%, 98% or 99% identical, to any of the nucleotide sequences in (a), (b), (c), or (d) above, or a polynucleotide which hybridizes under stringent hybridization conditions to a polynucleotide in (a), (b), (c), or (d) above. This polynucleotide which hybridizes does not hybridize under stringent hybridization conditions to a polynucleotide having a nucleotide sequence consisting of only A residues or of only T residues. An additional nucleic acid embodiment of the invention relates to an isolated nucleic acid molecule comprising a polynucleotide which encodes the amino acid sequence of an epitope-bearing portion of a PGRP-K, a PGRP-W, or a PGRP-C polypeptide having an amino acid sequence in (a), (b), or above.

The present invention also relates to recombinant vectors, which include the isolated nucleic acid molecules of the present invention, and to host cells containing the recombinant vectors, as well as to methods of making such vectors and host cells and for using them for the production of PGRP-K, PGRP-W, and/or PGRP-C polypeptides or peptides by recombinant techniques.

The invention further provides isolated PGRP-K, PGRP-W, and PGRP-C polypeptides comprising amino acid sequences selected from the group consisting of: (a) the amino acid sequence of the full-length PGRP-K polypeptide having the complete amino acid sequence shown in Figures 1A-B (SEQ ID NO:2), the amino acid sequence of the full-length PGRP-W polypeptide having the complete amino acid sequence shown in Figures 2A-C (SEQ ID NO:4), the amino acid sequence of the full-length PGRP-C polypeptide having the complete amino acid sequence shown in Figure 3 (SEQ ID NO:6), or as encoded by the cDNA clones contained in ATCC Numbers 203564, 203563, and 209683, respectively, deposited on December 23, 1998, and Mach 20, 1998; (b) the amino acid sequence of the predicted PGRP-like

domain of the PGRP-K polypeptide having the amino acid sequence at positions 24 to 107 in Figures 1A-B (SEQ ID NO:2), the predicted PGRP-like domain of the PGRP-W polypeptide having the amino acid sequence at positions 52 to 135 in Figures 2A-C (SEQ ID NO:4), and/or the predicted PGRP-like domain of the PGRP-C polypeptide having the amino acid sequence at positions 34 to 117 in Figure 3 (SEQ ID NO:6), or as encoded by the cDNA clones contained in ATCC Numbers 203564, 203563, and 209683, respectively, deposited on December 23, 1998, and March 20, 1998; (c) the amino acid sequence of the soluble PGRP-K, PGRP-W, and/or PGRP-C polypeptide having the PGRP-like domain but lacking the leader sequence, wherein each of these domains is defined below.

The polypeptides of the present invention also include polypeptides having an amino acid sequence with at least 90% similarity, and more preferably at least 95% similarity to those described in (a), (b), or (c) above, as well as polypeptides having an amino acid sequence at least 80% identical, more preferably at least 90% identical, and still more preferably 95%, 96%, 97%, 98% or 99% identical to those above.

An additional embodiment of this aspect of the invention relates to a peptide or polypeptide which has the amino acid sequence of an epitope-bearing portion of a PGRP-K, a PGRP-W, or a PGRP-C polypeptide having an amino acid sequence described in (a), (b), or (c) above. Peptides or polypeptides having the amino acid sequence of an epitope-bearing portion of a PGRP-K, a PGRP-W, or a PGRP-C polypeptide of the invention include portions of such polypeptides with at least six or seven, preferably at least nine, and more preferably at least about 30 amino acids to about 50 amino acids, although epitope-bearing polypeptides of any length up to and including the entire amino acid sequence of a polypeptide of the invention described above also are included in the invention. In another embodiment, the invention provides an isolated antibody that binds specifically to a polypeptide having an amino acid sequence described in (a), (b), or (c) above.

The invention further provides methods for isolating antibodies that bind specifically to an PGRP-K, PGRP-W, or PGRP-C polypeptide having an amino acid

sequence as described herein. Such antibodies are useful diagnostically or therapeutically as described below.

The invention also provides for pharmaceutical compositions comprising soluble PGRP-K, PGRP-W, and/or PGRP-C polypeptides, particularly human
5 PGRP-K, PGRP-W, and/or PGRP-C polypeptides, which may be employed, for instance, to treat tumor and tumor metastasis, infections by bacteria, viruses and other parasites, immunodeficiencies, inflammatory diseases, regulate the apoptosis and/or proliferation of keratinocytes, epidermal cells, and epithelial cells, mediate antigen processing and presentation, mediate cell activation and proliferation, and are
10 functionally linked as primary mediators of immune recognition and immune responses.

The invention further provides compositions comprising a PGRP-K, PGRP-W, or PGRP-C polynucleotide or a PGRP-K, PGRP-W, or PGRP-C polypeptide for
administration to cells *in vitro*, to cells *ex vivo* and to cells *in vivo*, or to a multicellular
15 organism. In certain particularly preferred embodiments of this aspect of the invention, the compositions comprise a PGRP-K, PGRP-W, or PGRP-C polynucleotide for expression of a PGRP-K, PGRP-W, or PGRP-C polypeptide in a host organism for treatment of disease. Particularly preferred in this regard is
expression in a human patient for treatment of a dysfunction associated with aberrant
20 endogenous activity of a PGRP-K, PGRP-W, or PGRP-C gene.

The present invention also provides a screening method for identifying compounds capable of enhancing or inhibiting a cellular response induced by PGRP-K, PGRP-W, or PGRP-C which involves contacting cells which express PGRP-K, PGRP-W, or PGRP-C with the candidate compound, assaying a cellular response, and
25 comparing the cellular response to a standard cellular response, the standard being assayed when contact is made in absence of the candidate compound; whereby, an increased cellular response over the standard indicates that the compound is an agonist and a decreased cellular response over the standard indicates that the compound is an antagonist.

In another aspect, a method for identifying PGRP-K, PGRP-W, or PGRP-C receptors is provided, as well as a screening assay for agonists and antagonists using such receptors. This assay involves determining the effect a candidate compound has on PGRP-K, PGRP-W, or PGRP-C binding to the PGRP-K, PGRP-W, or PGRP-C receptor. In particular, the method involves contacting a PGRP-K, PGRP-W, or PGRP-C receptor with an PGRP-K, PGRP-W, or PGRP-C polypeptide and a candidate compound and determining whether PGRP-K, PGRP-W, or PGRP-C polypeptide binding to the PGRP-K, PGRP-W, or PGRP-C receptor is increased or decreased due to the presence of the candidate compound. The antagonists may be employed to prevent septic shock, inflammation, and to regulate the growth activity of keratinocytes.

The present inventors have discovered that PGRP-K, PGRP-W, and PGRP-C is expressed in keratinocytes, wound healing tissues, and chondrosarcomas, respectively. For a number of disorders of these tissues and cells, such as tumor and tumor metastasis, infection of bacteria, viruses and other parasites, immunodeficiencies, septic shock, apoptosis or proliferation of these tissues, and proper antigen processing and presentation, it is believed that significantly higher or lower levels of the PGRP-K, PGRP-W, or PGRP-C gene expression can be detected in certain tissues (e.g., keratinocytes, wound-healing tissues, and chondrosarcoma) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) taken from an individual having such a disorder, relative to a "standard" PGRP-K, PGRP-W, or PGRP-C gene expression level, i.e., the PGRP-K, PGRP-W, or PGRP-C expression level in tissue or bodily fluids from an individual not having the disorder. Thus, the invention provides a diagnostic method useful during diagnosis of a disorder, which involves: (a) assaying PGRP-K, PGRP-W, or PGRP-C gene expression levels in cells or body fluid of an individual; (b) comparing the PGRP-K, PGRP-W, or PGRP-C gene expression level with a standard PGRP-K, PGRP-W, or PGRP-C gene expression level, whereby an increase or decrease in the assayed PGRP-K, PGRP-W,

or PGRP-C gene expression level compared to the standard expression level is indicative of a disorder.

An additional aspect of the invention is related to a method for treating an individual in need of an increased level of either PGRP-K, PGRP-W, or PGRP-C activity in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an isolated PGRP-K, PGRP-W, or PGRP-C polypeptide of the invention or an agonist thereof.

A still further aspect of the invention is related to a method for treating an individual in need of a decreased level of either PGRP-K, PGRP-W, or PGRP-C activity in the body comprising, administering to such an individual a composition comprising a therapeutically effective amount of a PGRP-K, PGRP-W, or PGRP-C antagonist. Preferred antagonists for use in the present invention are either PGRP-K, PGRP-W, or PGRP-C-specific antibodies.

Brief Description of the Figures

Figures 1A-B show the nucleotide sequence (SEQ ID NO:1) and deduced amino acid sequence (SEQ ID NO:2) of the PGRP-K. The deduced complete amino acid sequence includes 243 amino acid residues and has a deduced molecular weight of about 27,000 Da. The predicted domains of the PGRP-K are: signal sequence (amino acid residues Met-1 to about Ala-17 of SEQ ID NO:2) and PGRP-like domain (amino acid residues from about Val-24 to about His-107 of SEQ ID NO:2).

Figures 2A-C show the nucleotide sequence (SEQ ID NO:3) and deduced amino acid sequence (SEQ ID NO:4) of the PGRP-W. The deduced complete amino acid sequence includes 368 amino acid residues and has a deduced molecular weight of about 40,286 Da. The predicted domains of the PGRP-W are: signal sequence (amino acid residues Met-1 to about Gly-17 of SEQ ID NO:4) and PGRP-like domain (amino acid residues from about Val-52 to about His-135 of SEQ ID NO:4).

Figures 3 show the nucleotide sequence (SEQ ID NO:5) and deduced amino acid sequence (SEQ ID NO:6) of the PGRP-C. The deduced complete amino acid

sequence includes 196 amino acid residues and has a deduced molecular weight of about 21,500 Da. The predicted domains of the PGRP-C are: signal sequence (amino acid residues Met-1 to about Ala-21 of SEQ ID NO:6) and PGRP-like domain (amino acid residues from about Val-34 to about His-117 of SEQ ID NO:6).

5 Figures 4A-B show the regions of similarity between the amino acid sequences of the PGRP-K protein of Figures 1A-B (labeled PGRP-K (HKABZ65); SEQ ID NO:2), the PGRP-W protein of Figures 2A-C (labeled PGRP-W (HWHGB15); SEQ ID NO:4), the PGRP-C protein of Figure 3 (labeled PGRP-C (HCDDP40); SEQ ID NO:6), and the Mouse Tag-7 protein (SEQ ID NO:7) (GenBank Accession Number
10 X86374), as determined by the "Megalign" routine which is part of the computer program called "DNASTar". Identical amino acid residues between these protein sequences are shaded.

Figure 5 and Table II show a structural analysis of the PGRP-K amino acid sequence of Figures 1A-B (SEQ ID NO:2), generated using the default parameters of
15 the recited computer programs. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid residues: Val-24 to Ala-35; from Gln-51 to Gln-58; from Gly-69 to Ser-72; from Leu-88 to Gly-100; from His-107 to Tyr-111; from Gly-122 to Pro-131; from Gln-146 to
20 Ile-155; from Leu-159 to His-170; from Val-172 to Pro-200; from Gly-211 to Val-223; and from Phe-230 to Tyr-242 as depicted in Figures 1A-B (SEQ ID NO:2) correspond to the shown highly antigenic regions of the PGRP-K protein.

Figure 6 and Table III show a structural analysis of the PGRP-W amino acid sequence of Figures 2A-C (SEQ ID NO:4), generated using the default parameters of
25 the recited computer programs. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid residues: Gly-17 to Gly-32; from Ile-40 to Gly-58; from Gly-82 to Gln-99; from His-104 to Val-111; from Leu-116 to Glu-125; from Gly-150 to Pro-159; from Gln-174 to

Tyr-182; from Leu-186 to Pro-207; from Val-214 to Met-225; from Thr-237 to Val-252; from Tyr-259 to Ile-268; from Gly-290 to Ala-300; from His-344 to Gln-355; and from Trp-364 to His-368 as depicted in Figures 2A-C (SEQ ID NO:4) correspond to the shown highly antigenic regions of the PGRP-W protein.

5 Figure 7 and Table IV show a structural analysis of the PGRP-C amino acid sequence of Figure 3 (SEQ ID NO:6), generated using the default parameters of the recited computer programs. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid
10 residues: Met-1 to Met-6; from Ala-20 to Cys-29; from Ile-33 to Ala-43; from Ala-63 to Asn-79; from Ile-99 to Asn-112; from Gly-133 to Arg-146; from Ala-160 to Tyr-165; from Lys-168 to Asn-181; and from Trp-190 to Pro-196 as depicted in Figure 3 (SEQ ID NO:6) correspond to the shown highly antigenic regions of the PGRP-C protein.

15

Detailed Description

The present invention provides isolated nucleic acid molecules comprising polynucleotides encoding a PGRP-K, a PGRP-W and a PGRP-C polypeptides (Figures 1A-B, 2A-C, and 3 (SEQ ID NO:1, SEQ ID NO:3, and SEQ ID NO:5, respectively), the amino acid sequences of which were determined by sequencing
20 cloned cDNAs. The PGRP-K, PGRP-W, and PGRP-C proteins shown in Figures 1A-B, 2A-C, and 3, respectively, share sequence homology with the murine Tag-7 protein (Figures 4A-B (SEQ ID NO:7)) . On December 23, 1998, and March 20, 1998, deposits of plasmid DNAs encoding PGRP-K, PGRP-W, and PGRP-C were made at the American Type Culture Collection, 10801 University Boulevard,
25 Manassas, Virginia 20110-2209, and given accession numbers 203564, 203563, and 209683, respectively. The nucleotide sequences shown in Figures 1, 2, and 3 (SEQ ID NO:1, SEQ ID NO:3, and SEQ ID NO:5, respectively) were obtained by sequencing cDNA clones (Clone ID HKABZ65, HWHGB15, and HCDDP40, respectively)

containing the same amino acid coding sequences as the clones in ATCC Accession Nos. 203564, 203563, and 209683, respectively. The deposited clone encoding PGRP-K is contained in the pCMVSPORT2.0 plasmid (Life Technologies, Rockville, MD), the deposited clone encoding PGRP-W is contained in the pCMVSPORT3.0 plasmid (Life Technologies, Rockville, MD), and the deposited clone encoding the PGRP-C is contained in the Uni-Zap XR plasmid (Stratagene, La Jolla, CA).

Nucleic Acid Molecules

Unless otherwise indicated, all nucleotide sequences determined by sequencing a DNA molecule herein were determined using an automated DNA sequencer (such as the Model 373 from Applied Biosystems, Inc., Foster City, CA), and all amino acid sequences of polypeptides encoded by DNA molecules determined herein were predicted by translation of a DNA sequence determined as above. Therefore, as is known in the art for any DNA sequence determined by this automated approach, any nucleotide sequence determined herein may contain some errors. Nucleotide sequences determined by automation are typically at least about 90% identical, more typically at least about 95% to at least about 99.9% identical to the actual nucleotide sequence of the sequenced DNA molecule. The actual sequence can be more precisely determined by other approaches including manual DNA sequencing methods well known in the art. As is also known in the art, a single insertion or deletion in a determined nucleotide sequence compared to the actual sequence will cause a frame shift in translation of the nucleotide sequence such that the predicted amino acid sequence encoded by a determined nucleotide sequence will be completely different from the amino acid sequence actually encoded by the sequenced DNA molecule, beginning at the point of such an insertion or deletion.

By "nucleotide sequence" of a nucleic acid molecule or polynucleotide is intended, for a DNA molecule or polynucleotide, a sequence of deoxyribonucleotides, and for an RNA molecule or polynucleotide, the corresponding sequence of

ribonucleotides (A, G, C and U), where each thymidine deoxyribonucleotide (T) in the specified deoxyribonucleotide sequence is replaced by the ribonucleotide uridine (U).

Using the information provided herein, such as the nucleotide sequences in Figures 1 or 2, a nucleic acid molecule of the present invention encoding a peptidoglycan recognition protein polypeptide may be obtained using standard cloning and screening procedures, such as those for cloning cDNAs using mRNA as starting material. Illustrative of the invention, the nucleic acid molecule described in Figure 1 (SEQ ID NO:1) was discovered in a cDNA library derived from Human keratinocytes, the nucleic acid molecule described in Figure 2 (SEQ ID NO:3) was discovered in cDNA libraries derived from Human keratinocytes and Human tissues undergoing wound-healing, and the nucleic acid molecule described in Figure 3 (SEQ ID NO:5) was discovered in cDNA libraries derived from Human chondrosarcoma.

The PGRP-K gene contains an open reading frame encoding a protein of about 243 amino acid residues, a PGRP-like domain of about 137 amino acids (amino acid residues from about 18 to about 155 in Figures 1A-B (SEQ ID NO:2)), and a deduced molecular weight of about 27 kDa. The PGRP-K protein shown in Figures 1A-B (SEQ ID NO: 2) is about 40% similar to the mouse Tag-7 protein which can be accessed on Genbank as Accession No. X86374.

The PGRP-W gene contains an open reading frame encoding a protein of about 368 amino acid residues, a PGRP-like domain of about 175 amino acids (amino acid residues from about 18 to about 193 in Figures 2A-C (SEQ ID NO:4)), and a deduced molecular weight of about 40.2 kDa. The PGRP-W protein shown in Figures 2A-C (SEQ ID NO:4) is about 32% similar to the mouse Tag-7 protein which can be accessed on Genbank as Accession No. X86374.

The PGRP-C gene contains an open reading frame encoding a protein of about 196 amino acid residues, a PGRP-like domain of about 98 amino acids (amino acid residues from about 23 to about 120 in Figure 3 (SEQ ID NO:6)), and a deduced molecular weight of about 21.5 kDa. The PGRP-C protein shown in Figure 3 (SEQ

ID NO:6) is about 67% similar to the mouse Tag-7 protein which can be accessed on Genbank as Accession No. X86374.

Furthermore, PGRP-W is 59% homologous to PGRP-K, PGRP-W is 42% homologous to PGRP-C, and PGRP-K is 39% homologous to PGRP-C. More importantly, PGRP-K, PGRP-W, and PGRP-C share a conserved region of homology amongst the sequences, as well as with the murine Tag-7 protein. Also, four conserved cysteines are also homologous amongst all of these sequences, as can be seen in Figures 4A-B. Based upon alignments and sequence distances, it appears as if these three PGRPs constitute a novel class of peptidoglycan binding proteins, which share a conserved domain (PGRP-like domain), which may function in binding, while also having significantly diverging regions of homology as well. Thus, it appears as each of these PGRPs is a novel member of a family of peptidoglycan binding proteins thought to play an important role in immune recognition, immune surveillance, antigen presentation, and immune system activation.

As indicated, nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically. Isolated nucleic acid molecules of the

present invention include DNA molecules comprising an open reading frame (ORF) with an initiation codon starting at position 70 of the nucleotide sequence shown in Figures 1A-B (SEQ ID NO:1), an initiation codon starting at position 106 of the nucleotide sequence shown in Figures 2A-C (SEQ ID NO:3), and an initiation codon starting at position 55 of the nucleotide sequence shown in Figure 3 (SEQ ID NO: 5). As one of ordinary skill would appreciate, due to the possibilities of sequencing errors discussed above, the actual complete PGRP-C polypeptide encoded by the deposited cDNA, which comprises about 196 amino acids, may be somewhat shorter. In particular, the determined PGRP-C coding sequence contains a second methionine codon which may serve as an alternative start codon for translation of the open reading frame, at nucleotide positions 70-72 in Figure 3 (SEQ ID NO:5). More generally, the actual open reading frame may be anywhere in the range of ± 10 amino acids, more likely in the range of ± 6 amino acids, of that predicted from either the first or second methionine codon from the N-terminus shown in Figure 3 (SEQ ID NO:5). In addition, isolated nucleic acid molecules of the invention include DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode either the PGRP-K, PGRP-W, or PGRP-C proteins. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate the degenerate variants described above. In another aspect, the invention provides isolated nucleic acid molecules encoding the PGRP-K, PGRP-W, and PGRP-C polypeptides having amino acid sequences encoded by the cDNAs contained in the plasmids deposited on December 23, 1998, and March 20, 1998.

The invention further provides an isolated nucleic acid molecule having the nucleotide sequence shown in Figures 1A-B (SEQ ID NO:1) or the nucleotide sequence of the PGRP-K cDNA contained in the above-described deposited clone, or a nucleic acid molecule having a sequence complementary to one of the above sequences. Such isolated molecules, particularly DNA molecules, are useful as probes for gene mapping, by *in situ* hybridization with chromosomes, and for detecting

expression of the PGRP-K gene in human tissue, for instance, by Northern blot analysis.

The invention further provides an isolated nucleic acid molecule having the nucleotide sequence shown in Figures 2A-C (SEQ ID NO:3) or the nucleotide
5 sequence of the PGRP-W cDNA contained in the above-described deposited clone, or a nucleic acid molecule having a sequence complementary to one of the above sequences. Such isolated molecules, particularly DNA molecules, are useful as probes for gene mapping, by *in situ* hybridization with chromosomes, and for detecting expression of the PGRP-W gene in human tissue, for instance, by Northern blot
10 analysis.

The invention further provides an isolated nucleic acid molecule having the nucleotide sequence shown in Figure 3 (SEQ ID NO:5) or the nucleotide sequence of the PGRP-C cDNA contained in the above-described deposited clone, or a nucleic acid molecule having a sequence complementary to one of the above sequences. Such
15 isolated molecules, particularly DNA molecules, are useful as probes for gene mapping, by *in situ* hybridization with chromosomes, and for detecting expression of the PGRP-C gene in human tissue, for instance, by Northern blot analysis.

The present invention is further directed to nucleic acid molecules encoding portions of the nucleotide sequences described herein as well as to fragments of the
20 isolated nucleic acid molecules described herein. In particular, the invention provides a polynucleotide having a nucleotide sequence representing the portion of SEQ ID NO:1 which consists of positions 1-534 of SEQ ID NO:1, a polynucleotide having a nucleotide sequence representing the portion of SEQ ID NO:1 which consists of positions 535-798 of SEQ ID NO:1, a polynucleotide having a nucleotide sequence
25 representing the portion of SEQ ID NO:3 which consists of positions 1-685 of SEQ ID NO:3, a polynucleotide having a nucleotide sequence representing the portion of SEQ ID NO:3 which consists of positions 686-1210 of SEQ ID NO:3, a polynucleotide having a nucleotide sequence representing the portion of SEQ ID NO:5 which consists of positions 1-414 of SEQ ID NO:5, and a polynucleotide having a

nucleotide sequence representing the portion of SEQ ID NO:5 which consists of positions 415-642 of SEQ ID NO:5.

Further, the invention includes a polynucleotide comprising a sequence at least 95% identical to any portion of at least about 30 contiguous nucleotides, preferably at least about 50 nucleotides, of the sequence from nucleotides 1 to nucleotide 1150 in Figures 1A-B (SEQ ID NO:1), from nucleotides 1 to nucleotide 1854 in Figures 2A-C (SEQ ID NO:3), and from nucleotides 1 to nucleotide 726 in Figure 3 (SEQ ID NO:5).

More generally, by a fragment of an isolated nucleic acid molecule having the nucleotide sequence of the deposited cDNA or the nucleotide sequences shown in Figures 1A-B (SEQ ID NO:1), in Figures 2A-C (SEQ ID NO:3), or in Figure 3 (SEQ ID NO:5) is intended fragments at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments 50-300 nt in length are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequence of the deposited cDNA or as shown in Figures 1A-B (SEQ ID NO:1), in Figures 2A-C (SEQ ID NO:3), and/or Figure 3 (SEQ ID NO:5). By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequences of the deposited cDNAs or the nucleotide sequences as shown in Figures 1A-B (SEQ ID NO:1), in Figures 2A-C (SEQ ID NO:3), or in Figure 3 (SEQ ID NO:5). Preferred nucleic acid fragments of the present invention include nucleic acid molecules encoding epitope-bearing portions of the PGRP-K, PGRP-W, and/or PGRP-C polypeptides as identified in Figure 5 and Table II, Figure 6 and Table III, and Figure 7 and Table IV, respectively, and described in more detail below.

In another aspect, the invention provides isolated nucleic acid molecules comprising polynucleotides which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above, for instance, the cDNA clones contained in ATCC Deposit numbers 203564,

203563, or 209683, respectively, deposited on December 23, 1998, and March 20, 1998. By "stringent hybridization conditions" is intended overnight incubation at 42° C in a solution comprising: 50% formamide, 5x SSC (750 mM NaCl, 75 mM trisodium citrate), 75 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10%
5 dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65° C.

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least
10 about 30 nt, and even more preferably about 30-70 (e.g., 50) nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the
15 reference polynucleotides (e.g., the deposited cDNA or the nucleotide sequence as shown in Figures 1A-B (SEQ ID NO:1), in Figures 2A-C (SEQ ID NO:3), and/or in Figure 3 (SEQ ID NO:5)). Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3' terminal poly(A) tract of the PGRP-K, PGRP-W, or PGRP-C cDNA shown in Figures 1A-B (SEQ ID NO:1), in Figures 2A-C (SEQ ID
20 NO:3), and/or Figure 3 (SEQ ID NO:5)), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

25 As indicated, nucleic acid molecules of the present invention which encode a PGRP-K polypeptide may include, but are not limited to those encoding the amino acid sequence of the PGRP-like domain of the polypeptide, by itself; and the coding sequence for the PGRP-like domain of the polypeptide and additional sequences, such as a pre-, or pro- or prepro- protein sequence.

As indicated, nucleic acid molecules of the present invention which encode a PGRP-W polypeptide may include, but are not limited to those encoding the amino acid sequence of the PGRP-like domain of the polypeptide, by itself; and the coding sequence for the PGRP-like domain of the polypeptide and additional sequences, such as a pre-, or pro- or prepro- protein sequence.

As indicated, nucleic acid molecules of the present invention which encode a PGRP-C polypeptide may include, but are not limited to those encoding the amino acid sequence of the PGRP-like domain of the polypeptide, by itself; and the coding sequence for the PGRP-like domain of the polypeptide and additional sequences, such as a pre-, or pro or prepro- protein sequence.

Also encoded by nucleic acids of the invention are the above protein sequences together with additional, non-coding sequences, including for example, but not limited to introns and non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription, mRNA processing, including splicing and polyadenylation signals, for example - ribosome binding and stability of mRNA; an additional coding sequence which codes for additional amino acids, such as those which provide additional functionalities.

Thus, the sequence encoding the polypeptide may be fused to a marker sequence, such as a sequence encoding a peptide which facilitates purification of the fused polypeptide. In certain preferred embodiments of this aspect of the invention, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. The "HA" tag is another peptide useful for purification which corresponds to an epitope derived from the influenza hemagglutinin protein, which has been described by Wilson et al., *Cell* 37: 767 (1984). As discussed below, other such fusion proteins include either the PGRP-K, PGRP-W, or the PGRP-C fused to Fc at the N- or C-terminus.

Variant and Mutant Polynucleotides

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of either the PGRP-K, PGRP-W, or PGRP-C proteins. Variants may occur naturally, such as a
5 natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. *Genes II*, Lewin, B., ed., John Wiley & Sons, New York (1985). Non-naturally occurring variants may be produced using art-known mutagenesis techniques.

Such variants include those produced by nucleotide substitutions, deletions or
10 additions. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding regions, non-coding regions, or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and
15 activities of either the PGRP-K, PGRP-W, and/or PGRP-C proteins or portions thereof. Also especially preferred in this regard are conservative substitutions.

Most highly preferred are nucleic acid molecules encoding the PGRP-like domain 1 of the protein having the amino acid sequence shown in Figures 1A-B (SEQ ID NO:2) or the PGRP-like domain of the PGRP-K amino acid sequence encoded by
20 the deposited cDNA clone, the nucleic acid molecules encoding the PGRP-like domain of the protein having the amino acid sequence shown in Figures 2A-C (SEQ ID NO:4) or the PGRP-like domain of the PGRP-W amino acid sequence encoded by the deposited cDNA clone, or the nucleic acid molecules encoding the PGRP-like domain of the protein having the amino acid sequence shown in Figure 3 (SEQ ID NO:6) or
25 the PGRP-like domain of the PGRP-C amino acid sequence encoded by the deposited cDNA clone. Further embodiments include an isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 90% identical, and more preferably at least 95%, 96%, 97%, 98% or 99% identical to a polynucleotide

selected from the group consisting of: (a) nucleotide sequences encoding either the PGRP-K, PGRP-W, or PGRP-C polypeptides having the complete amino acid sequences in Figures 1A-B (SEQ ID NO:2), in Figures 2A-C (SEQ ID NO:4), and/or in Figure 3 (SEQ ID NO:6); (b) a nucleotide sequence encoding the predicted PGRP-like domain of the PGRP-K, PGRP-W, and PGRP-C polypeptides having the amino acid sequences at positions 72-323 in Figures 1A-B (SEQ ID NO:2), the amino acid sequences at positions 156-407 in Figures 2A-C (SEQ ID NO:4), and the amino acid sequence at positions 102-353 in Figure 3 (SEQ ID NO:6), respectively; (c) a nucleotide sequence encoding the PGRP-K, PGRP-W, or PGRP-C polypeptides having the complete amino acid sequences encoded by the cDNA clones contained in ATCC Numbers 203564, 203563, and 209683, respectively, deposited on December 23, 1998, and March 20, 1998; (d) a nucleotide sequence encoding the PGRP-like domain of the PGRP-K, PGRP-W, or PGRP-C polypeptides having the amino acid sequences encoded by the cDNA clones contained in ATCC Numbers 203564, 203563, and 209683, respectively, deposited on December 23, 1998, and March 20, 1998; and (e) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c) or (d) above.

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding a PGRP-K, a PGRP-W, or a PGRP-C polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence(s) except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding either the PGRP-K, PGRP-W, or PGRP-C polypeptides. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or

anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least
5 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the nucleotide sequences shown in Figures 1A-B, Figures 2A-C, and/or Figure 3 or to the nucleotide sequences of the deposited cDNA clones can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park,
10 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2:482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the
15 parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The present application is directed to nucleic acid molecules at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequences shown in Figures
20 1A-B (SEQ ID NO:1), Figure 2A-C (SEQ ID NO:3), and/or Figure 3 (SEQ ID NO:5) or to the nucleic acid sequence(s) of the deposited cDNAs, irrespective of whether they encode a polypeptide having PGRP-K, PGRP-W, or PGRP-C activities, respectively. This is because even where a particular nucleic acid molecule does not encode a polypeptide having either PGRP-K, PGRP-W, or PGRP-C activities, one of
25 skill in the art would still know how to use the nucleic acid molecules, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having either PGRP-K, PGRP-W, or PGRP-C activity include, inter alia, (1) isolating the PGRP-K, PGRP-W, or PGRP-C genes or allelic variants thereof in cDNA

libraries; (2) *in situ* hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the PGRP-K, PGRP-W, or PGRP-C genes, as described in Verma et al., *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988); and Northern Blot analysis for detecting either
5 PGRP-K, PGRP-W, or PGRP-C mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequences shown in Figures 1A-B (SEQ ID NO:1), Figures 2A-C (SEQ ID NO:3), and/or Figure 3 (SEQ ID NO:5), or to the nucleic acid sequence of the deposited cDNAs which do, in fact,
10 encode a polypeptide(s) having either PGRP-K, PGRP-W, or PGRP-C protein activity. By "a polypeptide having PGRP-K, PGRP-W, or PGRP-C activity" is intended polypeptides exhibiting activity similar, but not necessarily identical, to an activity of the PGRP-K, PGRP-W, or PGRP-C proteins of the invention, as measured in a particular biological assay. For example, the PGRP-K, PGRP-W, or PGRP-C
15 proteins of the present invention bind insoluble peptidoglycan. An insoluble peptidoglycan binding assay for measuring the ability of a protein to bind to insoluble peptidoglycan can be performed by using reagents well known and commonly available in the art for detecting this binding ability. For instance, numerous such assays for peptidoglycan binding protein activities are described in the various
20 references in the Background section of this disclosure, above, as well as in example 13. Briefly, such an assay involves collecting insoluble peptidoglycan from an appropriate source (e.g., *Micrococcus luteus*), mixing the insoluble peptidoglycan with a specified amount of the protein in question, and measuring the concentration of peptidoglycan bound by the protein over a certain period of time. Such insoluble
25 peptidoglycan binding activities as can be measure in this type of assay are useful for identifying proteins that may have a immuno-modulatory effect in the body, and that may act to bind peptidoglycan and/or function in an immune recognition capacity during the infection process of certain bacterial species.

PGRP-K, PGRP-W, and/or PGRP-C proteins bind insoluble peptidoglycan, and are useful for identifying proteins that may have an immuno-modulatory effect in the body, and that may act to bind peptidoglycan and/or function in an immune recognition capacity during the infection process of certain bacterial species. Thus, "a polypeptide having PGRP-K, PGRP-W, and/or PGRP-C protein activity" includes polypeptides that also exhibit any of the same peptidoglycan binding activities in the above-described assays. Although the degree of peptidoglycan binding activity need not be identical to that of the PGRP-K, PGRP-W, and/or PGRP-C proteins, preferably, "a polypeptide having either PGRP-K, PGRP-W, and/or PGRP-C protein activity" will exhibit substantially similar peptidoglycan binding in a given activity as compared to the PGRP-K, PGRP-W, and/or PGRP-C proteins (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity relative to the reference PGRP-K, PGRP-W, and/or PGRP-C proteins). Assays for measuring such activity are known in the art. For example, see Yoshida *et al.*, JBC, 271 (23): 13854 (1996); and Kang *et al.*, PNAS (US), 95 (17): 10078 (1998).

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleic acid sequences of the deposited cDNAs or the nucleic acid sequences shown in Figures 1A-B (SEQ ID NO:1), in Figures 2A-C (SEQ ID NO:3), and/or Figure 3 (SEQ ID NO:5) will encode a polypeptide "having PGRP-K, PGRP-W, and/or PGRP-C protein activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptides, respectively, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode polypeptides having either PGRP-K, PGRP-W, and/or PGRP-C protein activity, respectively. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely

to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid), as further described below.

Vectors and Host Cells

The present invention also relates to vectors which include the isolated DNA
5 molecules of the present invention, host cells which are genetically engineered with the recombinant vectors, and the production of either PGRP-K, PGRP-W, and/or PGRP-C polypeptides or fragments thereof by recombinant techniques. The vector may be, for example, a phage, plasmid, viral or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation
10 generally will occur only in complementing host cells. The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into
15 host cells.

The DNA insert(s) should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli lac*, *trp*, *phoA* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression
20 constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the PGRP-K, PGRP-W, and/or PGRP-C transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide
25 to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin

resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, 293 and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pHE4 (HGS Inc., Provisional Number: PCT/US98/20075); pA2, PO4, and pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Selection of appropriate vectors and promoters for expression in a host cell is a well known procedure and the requisite techniques for expression vector construction, introduction of the vector into the host and expression in the host are routine skills in the art.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., PGRP-K, PGRP-W, and/or PGRP-C coding sequences), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with PGRP-K, PGRP-W, and/or PGRP-C polynucleotides of the invention, respectively, and which activates, alters, and/or amplifies endogenous PGRP-K, PGRP-W, and/or PGRP-C polynucleotides, respectively. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous PGRP-K, PGRP-W, and/or PGRP-C polynucleotide sequences via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-

438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

The host cell can be a higher eukaryotic cell, such as a mammalian cell (e.g., a human derived cell), or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. The host strain may be chosen which modulates the expression of the inserted gene sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus expression of the genetically engineered polypeptide may be controlled. Furthermore, different host cells have characteristics and specific mechanisms for the translational and post-translational processing and modification (e.g., glycosylation, phosphorylation, cleavage) of proteins. Appropriate cell lines can be chosen to ensure the desired modifications and processing of the foreign protein expressed.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986).

The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals, but also additional heterologous functional regions. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell, during purification, or during subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine techniques in the art. A preferred fusion protein comprises a heterologous region from immunoglobulin that is useful to stabilize and purify proteins. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin

molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (EP-A 0232 262). On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when Fc portion proves to be a hindrance to use in therapy and diagnosis, for example when the fusion protein is to be used as antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5 has been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett et al., *J. Molecular Recognition* 8:52-58 (1995) and K. Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995).

The PGRP-K, PGRP-W, or PGRP-C protein can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

PGRP-K, PGRP-W, and PGRP-C Polypeptides and Fragments

The invention further provides an isolated PGRP-K polypeptide having the amino acid sequence encoded by the deposited cDNA, or the amino acid sequence in Figures 1A-B (SEQ ID NO:2), or a peptide or polypeptide comprising a portion of the above polypeptides.

The invention further provides an isolated PGRP-W polypeptide having the amino acid sequence encoded by the deposited cDNA, or the amino acid sequence in Figures 2A-C (SEQ ID NO:4), or a peptide or polypeptide comprising a portion of the above polypeptides.

The invention further provides an isolated PGRP-C polypeptide having the amino acid sequence encoded by the deposited cDNA, or the amino acid sequence in Figure 3 (SEQ ID NO:6), or a peptide or polypeptide comprising a portion of the above polypeptides.

Variant and Mutant Polypeptides

To improve or alter the characteristics of either PGRP-K, PGRP-W, and/or PGRP-C polypeptides, protein engineering may be employed. Recombinant DNA technology known to those skilled in the art can be used to create novel mutant proteins or "muteins including single or multiple amino acid substitutions, deletions, additions or fusion proteins. Such modified polypeptides can show, e.g., enhanced activity or increased stability. In addition, they may be purified in higher yields and show better solubility than the corresponding natural polypeptide, at least under certain purification and storage conditions.

N-Terminal and C-Terminal Deletion Mutants

For instance, for many proteins, including the extracellular domain or the mature form(s) of a secreted protein, it is known in the art that one or more amino acids may be deleted from the N-terminus or C-terminus without substantial loss of biological function. For instance, Ron et al., *J. Biol. Chem.*, 268:2984-2988 (1993)

reported modified KGF proteins that had heparin binding activity even if 3, 8, or 27 amino-terminal amino acid residues were missing.

In the present case, since the protein of the invention is related to Tag-7, deletions of N-terminal amino acids up to the Gly (G) residue at position 191 in Figure 1 (SEQ ID NO:2) may retain some biological activity such as cytotoxicity to appropriate target cells. However, even if deletion of one or more amino acids from the N-terminus of a protein results in modification of loss of one or more biological functions of the protein, other biological activities may still be retained. Thus, the ability of the shortened protein to induce and/or bind to antibodies which recognize the complete PGRP-like domain of the protein generally will be retained when less than the majority of the residues of the complete PGRP domain of the protein are removed from the N-terminus. Whether a particular polypeptide lacking N-terminal residues of a complete protein retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art.

In one embodiment, the present invention further provides polypeptides having one or more residues deleted from the amino terminus of the amino acid sequence of the PGRP-K polypeptide depicted in Figures 1A-B (SEQ ID NO:2) or encoded by the cDNA of the deposited clone. Particularly, in one embodiment, N-terminal deletions of the PGRP-K polypeptide can be described by the general formula m to 243, where m is an integer from 2 to 243 corresponding to the position of the amino acid identified in SEQ ID NO:2 and preferably, corresponds to one of the N-terminal amino acid residues identified in the N-terminal deletions specified herein. In specific embodiments, N-terminal deletions of the PGRP-K polypeptide of the invention comprise, or alternatively, consist of, amino acid residues: G-2 to Q-243; T-3 to Q-243; L-4 to Q-243; P-5 to Q-243; W-6 to Q-243; L-7 to Q-243; L-8 to Q-243; A-9 to Q-243; F-10 to Q-243; F-11 to Q-243; I-12 to Q-243; L-13 to Q-243; G-14 to Q-243; L-15 to Q-243; Q-16 to Q-243; A-17 to Q-243; W-18 to Q-243; D-19 to Q-243; T-20 to Q-243; P-21 to Q-243; T-22 to Q-243; I-23 to Q-243; V-24 to Q-243; S-25 to Q-243; R-26 to Q-243; K-27 to Q-243; E-28 to Q-243; W-29 to Q-243; G-30 to Q-243; A-31 to Q-243; R-32 to Q-243; P-33 to Q-243; L-34 to Q-243; A-35 to Q-243; C-36 to Q-243; R-37 to Q-243; A-38 to Q-243; L-39 to Q-243; L-40 to Q-243; T-41 to Q-243; L-42 to Q-243; P-43 to Q-243; V-44 to Q-243; A-45 to Q-243; Y-46 to Q-243; I-47 to Q-243; I-48 to Q-243; T-49 to Q-243; D-50 to Q-243; Q-51 to Q-243; L-52 to Q-243; P-53 to Q-243;

G-54 to Q-243; M-55 to Q-243; Q-56 to Q-243; C-57 to Q-243; Q-58 to Q-243; Q-59 to Q-243; Q-60 to Q-243; S-61 to Q-243; V-62 to Q-243; C-63 to Q-243; S-64 to Q-243; Q-65 to Q-243; M-66 to Q-243; L-67 to Q-243; R-68 to Q-243; G-69 to Q-243; L-70 to Q-243; Q-71 to Q-243; S-72 to Q-243; H-73 to Q-243; S-74 to Q-243; V-75 to Q-243; 5 Y-76 to Q-243; T-77 to Q-243; I-78 to Q-243; G-79 to Q-243; W-80 to Q-243; C-81 to Q-243; D-82 to Q-243; V-83 to Q-243; A-84 to Q-243; Y-85 to Q-243; N-86 to Q-243; F-87 to Q-243; L-88 to Q-243; V-89 to Q-243; G-90 to Q-243; D-91 to Q-243; D-92 to Q-243; G-93 to Q-243; R-94 to Q-243; V-95 to Q-243; Y-96 to Q-243; E-97 to Q-243; G-98 to Q-243; V-99 to Q-243; G-100 to Q-243; W-101 to Q-243; N-102 to Q-243; I- 10 103 to Q-243; Q-104 to Q-243; G-105 to Q-243; L-106 to Q-243; H-107 to Q-243; T-108 to Q-243; Q-109 to Q-243; G-110 to Q-243; Y-111 to Q-243; N-112 to Q-243; N-113 to Q-243; I-114 to Q-243; S-115 to Q-243; L-116 to Q-243; G-117 to Q-243; I-118 to Q-243; A-119 to Q-243; F-120 to Q-243; F-121 to Q-243; G-122 to Q-243; N-123 to Q-243; K-124 to Q-243; I-125 to Q-243; S-126 to Q-243; S-127 to Q-243; S-128 to Q- 15 243; P-129 to Q-243; S-130 to Q-243; P-131 to Q-243; A-132 to Q-243; A-133 to Q-243; L-134 to Q-243; S-135 to Q-243; A-136 to Q-243; A-137 to Q-243; E-138 to Q-243; G-139 to Q-243; L-140 to Q-243; I-141 to Q-243; S-142 to Q-243; Y-143 to Q-243; A-144 to Q-243; I-145 to Q-243; Q-146 to Q-243; K-147 to Q-243; G-148 to Q-243; H-149 to Q-243; L-150 to Q-243; S-151 to Q-243; P-152 to Q-243; R-153 to Q- 20 243; Y-154 to Q-243; I-155 to Q-243; Q-156 to Q-243; P-157 to Q-243; L-158 to Q-243; L-159 to Q-243; L-160 to Q-243; K-161 to Q-243; E-162 to Q-243; E-163 to Q-243; T-164 to Q-243; C-165 to Q-243; L-166 to Q-243; D-167 to Q-243; P-168 to Q-243; Q-169 to Q-243; H-170 to Q-243; P-171 to Q-243; V-172 to Q-243; M-173 to Q-243; P-174 to Q-243; R-175 to Q-243; K-176 to Q-243; V-177 to Q-243; C-178 to Q- 25 243; P-179 to Q-243; N-180 to Q-243; I-181 to Q-243; I-182 to Q-243; K-183 to Q-243; R-184 to Q-243; S-185 to Q-243; A-186 to Q-243; W-187 to Q-243; E-188 to Q-243; A-189 to Q-243; R-190 to Q-243; E-191 to Q-243; T-192 to Q-243; H-193 to Q-243; C-194 to Q-243; P-195 to Q-243; K-196 to Q-243; M-197 to Q-243; N-198 to Q-243; L-199 to Q-243; P-200 to Q-243; A-201 to Q-243; K-202 to Q-243; Y-203 to Q- 30 243; V-204 to Q-243; I-205 to Q-243; I-206 to Q-243; I-207 to Q-243; H-208 to Q-243; T-209 to Q-243; A-210 to Q-243; G-211 to Q-243; T-212 to Q-243; S-213 to Q-243; C-214 to Q-243; T-215 to Q-243; V-216 to Q-243; S-217 to Q-243; T-218 to Q-243; D-219 to Q-243; C-220 to Q-243; Q-221 to Q-243; T-222 to Q-243; V-223 to Q-243; V-224 to Q-243; R-225 to Q-243; N-226 to Q-243; I-227 to Q-243; Q-228 to Q-243; S- 35 229 to Q-243; F-230 to Q-243; H-231 to Q-243; M-232 to Q-243; D-233 to Q-243; T-234 to Q-243; R-235 to Q-243; N-236 to Q-243; F-237 to Q-243; C-238 to Q-243; of SEQ ID NO:2. Polynucleotides encoding these polypeptides also are provided.

In one embodiment, the present invention further provides polypeptides having one or more residues deleted from the amino terminus of the amino acid sequence of the PGRP-W polypeptide depicted in Figures 2A-C (SEQ ID NO:4) or encoded by the cDNA of the deposited clone. Particularly, in one embodiment, N-terminal deletions of the PGRP-W polypeptide can be described by the general formula m to 368, where m is an integer from 2 to 368 corresponding to the position of the amino acid identified in SEQ ID NO:4 and preferably, corresponds to one of the N-terminal amino acid residues identified in the N-terminal deletions specified herein. In specific embodiments, N-terminal deletions of the PGRP-W polypeptide of the invention comprise, or alternatively, consist of, amino acid residues: L-2 to H-368; L-3 to H-368; W-4 to H-368; L-5 to H-368; L-6 to H-368; V-7 to H-368; F-8 to H-368; S-9 to H-368; A-10 to H-368; L-11 to H-368; G-12 to H-368; I-13 to H-368; Q-14 to H-368; A-15 to H-368; W-16 to H-368; G-17 to H-368; D-18 to H-368; S-19 to H-368; S-20 to H-368; W-21 to H-368; N-22 to H-368; K-23 to H-368; T-24 to H-368; Q-25 to H-368; A-26 to H-368; K-27 to H-368; Q-28 to H-368; V-29 to H-368; S-30 to H-368; E-31 to H-368; G-32 to H-368; L-33 to H-368; Q-34 to H-368; Y-35 to H-368; L-36 to H-368; F-37 to H-368; E-38 to H-368; N-39 to H-368; I-40 to H-368; S-41 to H-368; Q-42 to H-368; L-43 to H-368; T-44 to H-368; E-45 to H-368; K-46 to H-368; D-47 to H-368; V-48 to H-368; S-49 to H-368; T-50 to H-368; T-51 to H-368; V-52 to H-368; S-53 to H-368; R-54 to H-368; K-55 to H-368; A-56 to H-368; W-57 to H-368; G-58 to H-368; A-59 to H-368; E-60 to H-368; A-61 to H-368; V-62 to H-368; G-63 to H-368; C-64 to H-368; S-65 to H-368; I-66 to H-368; Q-67 to H-368; L-68 to H-368; T-69 to H-368; T-70 to H-368; P-71 to H-368; V-72 to H-368; N-73 to H-368; V-74 to H-368; L-75 to H-368; V-76 to H-368; I-77 to H-368; H-78 to H-368; H-79 to H-368; V-80 to H-368; P-81 to H-368; G-82 to H-368; L-83 to H-368; E-84 to H-368; C-85 to H-368; H-86 to H-368; D-87 to H-368; Q-88 to H-368; T-89 to H-368; V-90 to H-368; C-91 to H-368; S-92 to H-368; Q-93 to H-368; R-94 to H-368; L-95 to H-368; R-96 to H-368; E-97 to H-368; L-98 to H-368; Q-99 to H-368; A-100 to H-368; H-101 to H-368; H-102 to H-368; V-103 to H-368; H-104 to H-368; N-105 to H-368; N-106 to H-368; S-107 to H-368; G-108 to H-368; C-109 to H-368; D-110 to H-368; V-111 to H-368; A-112 to H-368; Y-113 to H-368; N-114 to H-368; F-115 to H-368; L-116 to H-368; V-117 to H-368; G-118 to H-368; D-119 to H-368; D-120 to H-368; G-121 to H-368; R-122 to H-368; V-123 to H-368; Y-124 to H-368; E-125 to H-368; G-126 to H-368; V-127 to H-368; G-128 to H-368; W-129 to H-368; N-130 to H-368; I-131 to H-368; Q-132 to H-368; G-133 to H-368; V-134 to H-368; H-135 to H-368; T-136 to H-368; Q-137 to H-368; G-138 to H-368; Y-139 to H-368; N-140 to H-368; N-141 to H-368; I-142 to H-368; S-143 to H-368; L-144 to H-368; G-145 to H-368; F-146 to H-

368; A-147 to H-368; F-148 to H-368; F-149 to H-368; G-150 to H-368; T-151 to H-368; K-152 to H-368; K-153 to H-368; G-154 to H-368; H-155 to H-368; S-156 to H-368; P-157 to H-368; S-158 to H-368; P-159 to H-368; A-160 to H-368; A-161 to H-368; L-162 to H-368; S-163 to H-368; A-164 to H-368; M-165 to H-368; E-166 to H-368; N-167 to H-368; L-168 to H-368; I-169 to H-368; T-170 to H-368; Y-171 to H-368; A-172 to H-368; V-173 to H-368; Q-174 to H-368; K-175 to H-368; G-176 to H-368; H-177 to H-368; L-178 to H-368; S-179 to H-368; S-180 to H-368; S-181 to H-368; Y-182 to H-368; V-183 to H-368; Q-184 to H-368; P-185 to H-368; L-186 to H-368; L-187 to H-368; G-188 to H-368; K-189 to H-368; G-190 to H-368; E-191 to H-368; N-192 to H-368; C-193 to H-368; L-194 to H-368; A-195 to H-368; P-196 to H-368; R-197 to H-368; Q-198 to H-368; K-199 to H-368; T-200 to H-368; S-201 to H-368; L-202 to H-368; K-203 to H-368; K-204 to H-368; L-205 to H-368; A-206 to H-368; P-207 to H-368; A-208 to H-368; L-209 to H-368; S-210 to H-368; H-211 to H-368; G-212 to H-368; L-213 to H-368; C-214 to H-368; G-215 to H-368; E-216 to H-368; P-217 to H-368; G-218 to H-368; R-219 to H-368; P-220 to H-368; L-221 to H-368; S-222 to H-368; R-223 to H-368; M-224 to H-368; T-225 to H-368; L-226 to H-368; P-227 to H-368; A-228 to H-368; K-229 to H-368; Y-230 to H-368; G-231 to H-368; I-232 to H-368; I-233 to H-368; I-234 to H-368; H-235 to H-368; T-236 to H-368; A-237 to H-368; G-238 to H-368; R-239 to H-368; T-240 to H-368; C-241 to H-368; N-242 to H-368; I-243 to H-368; S-244 to H-368; D-245 to H-368; E-246 to H-368; C-247 to H-368; R-248 to H-368; L-249 to H-368; L-250 to H-368; V-251 to H-368; R-252 to H-368; D-253 to H-368; I-254 to H-368; Q-255 to H-368; S-256 to H-368; F-257 to H-368; Y-258 to H-368; I-259 to H-368; D-260 to H-368; R-261 to H-368; L-262 to H-368; K-263 to H-368; S-264 to H-368; C-265 to H-368; D-266 to H-368; I-267 to H-368; G-268 to H-368; Y-269 to H-368; N-270 to H-368; F-271 to H-368; L-272 to H-368; V-273 to H-368; G-274 to H-368; Q-275 to H-368; D-276 to H-368; G-277 to H-368; A-278 to H-368; I-279 to H-368; Y-280 to H-368; E-281 to H-368; G-282 to H-368; V-283 to H-368; G-284 to H-368; W-285 to H-368; N-286 to H-368; V-287 to H-368; Q-288 to H-368; G-289 to H-368; S-290 to H-368; S-291 to H-368; T-292 to H-368; P-293 to H-368; G-294 to H-368; Y-295 to H-368; D-296 to H-368; D-297 to H-368; I-298 to H-368; A-299 to H-368; L-300 to H-368; G-301 to H-368; I-302 to H-368; T-303 to H-368; F-304 to H-368; M-305 to H-368; G-306 to H-368; T-307 to H-368; F-308 to H-368; T-309 to H-368; G-310 to H-368; I-311 to H-368; P-312 to H-368; P-313 to H-368; N-314 to H-368; A-315 to H-368; A-316 to H-368; A-317 to H-368; L-318 to H-368; E-319 to H-368; A-320 to H-368; A-321 to H-368; Q-322 to H-368; D-323 to H-368; L-324 to H-368; I-325 to H-368; Q-326 to H-368; C-327 to H-368; A-328 to H-368; M-329 to H-368; V-330 to H-368; K-331 to

H-368; G-332 to H-368; Y-333 to H-368; L-334 to H-368; T-335 to H-368; P-336 to H-368; N-337 to H-368; Y-338 to H-368; L-339 to H-368; L-340 to H-368; V-341 to H-368; G-342 to H-368; H-343 to H-368; S-344 to H-368; D-345 to H-368; V-346 to H-368; A-347 to H-368; R-348 to H-368; T-349 to H-368; L-350 to H-368; S-351 to H-368; P-352 to H-368; G-353 to H-368; Q-354 to H-368; A-355 to H-368; L-356 to H-368; Y-357 to H-368; N-358 to H-368; I-359 to H-368; I-360 to H-368; S-361 to H-368; T-362 to H-368; W-363 to H-368; of SEQ ID NO:4. Polynucleotides encoding these polypeptides also are provided.

In one embodiment, the present invention further provides polypeptides having one or more residues deleted from the amino terminus of the amino acid sequence of the PGRP-C polypeptide depicted in Figure 3 (SEQ ID NO:6) or encoded by the cDNA of the deposited clone. Particularly, in one embodiment, N-terminal deletions of the PGRP-C polypeptide can be described by the general formula m to 196, where m is an integer from 2 to 196 corresponding to the position of the amino acid identified in SEQ ID NO:6 and preferably, corresponds to one of the N-terminal amino acid residues identified in the N-terminal deletions specified herein. In specific embodiments, N-terminal deletions of the PGRP-C polypeptide of the invention comprise, or alternatively, consist of, amino acid residues: S-2 to P-196; R-3 to P-196; R-4 to P-196; S-5 to P-196; M-6 to P-196; L-7 to P-196; L-8 to P-196; A-9 to P-196; W-10 to P-196; A-11 to P-196; L-12 to P-196; P-13 to P-196; S-14 to P-196; L-15 to P-196; L-16 to P-196; R-17 to P-196; L-18 to P-196; G-19 to P-196; A-20 to P-196; A-21 to P-196; Q-22 to P-196; E-23 to P-196; T-24 to P-196; E-25 to P-196; D-26 to P-196; P-27 to P-196; A-28 to P-196; C-29 to P-196; C-30 to P-196; S-31 to P-196; P-32 to P-196; I-33 to P-196; V-34 to P-196; P-35 to P-196; R-36 to P-196; N-37 to P-196; E-38 to P-196; W-39 to P-196; K-40 to P-196; A-41 to P-196; L-42 to P-196; A-43 to P-196; S-44 to P-196; E-45 to P-196; C-46 to P-196; A-47 to P-196; Q-48 to P-196; H-49 to P-196; L-50 to P-196; S-51 to P-196; L-52 to P-196; P-53 to P-196; L-54 to P-196; R-55 to P-196; Y-56 to P-196; V-57 to P-196; V-58 to P-196; V-59 to P-196; S-60 to P-196; H-61 to P-196; T-62 to P-196; A-63 to P-196; G-64 to P-196; S-65 to P-196; S-66 to P-196; C-67 to P-196; N-68 to P-196; T-69 to P-196; P-70 to P-196; A-71 to P-196; S-72 to P-196; C-73 to P-196; Q-74 to P-196; Q-75 to P-196; Q-76 to P-196; A-77 to P-196; R-78 to P-196; N-79 to P-196; V-80 to P-196; Q-81 to P-196; H-82 to P-196; Y-83 to P-196; H-84 to P-196; M-85 to P-196; K-86 to P-196; T-87 to P-196; L-88 to P-196; G-89 to P-196; W-90 to P-196; C-91 to P-196; D-92 to P-196; V-93 to P-196; G-94 to P-196; Y-95 to P-196; N-96 to P-196; F-97 to P-196; L-98 to P-196; I-99 to P-196; G-100 to P-196; E-101 to P-196; D-102 to P-196; G-103 to P-196; L-104 to P-196; V-105 to P-196; Y-106 to P-196; E-107 to P-196; G-108 to P-196; R-109 to P-

196; G-110 to P-196; W-111 to P-196; N-112 to P-196; F-113 to P-196; T-114 to P-196; G-115 to P-196; A-116 to P-196; H-117 to P-196; S-118 to P-196; G-119 to P-196; H-120 to P-196; L-121 to P-196; W-122 to P-196; N-123 to P-196; P-124 to P-196; M-125 to P-196; S-126 to P-196; I-127 to P-196; G-128 to P-196; I-129 to P-196;
 5 S-130 to P-196; F-131 to P-196; M-132 to P-196; G-133 to P-196; N-134 to P-196; Y-135 to P-196; M-136 to P-196; D-137 to P-196; R-138 to P-196; V-139 to P-196; P-140 to P-196; T-141 to P-196; P-142 to P-196; Q-143 to P-196; A-144 to P-196; I-145 to P-196; R-146 to P-196; A-147 to P-196; A-148 to P-196; Q-149 to P-196; G-150 to P-196; L-151 to P-196; L-152 to P-196; A-153 to P-196; C-154 to P-196; G-155 to P-196;
 10 V-156 to P-196; A-157 to P-196; Q-158 to P-196; G-159 to P-196; A-160 to P-196; L-161 to P-196; R-162 to P-196; S-163 to P-196; N-164 to P-196; Y-165 to P-196; V-166 to P-196; L-167 to P-196; K-168 to P-196; G-169 to P-196; H-170 to P-196; R-171 to P-196; D-172 to P-196; V-173 to P-196; Q-174 to P-196; R-175 to P-196; T-176 to P-196; L-177 to P-196; S-178 to P-196; P-179 to P-196; G-180 to P-196;
 15 196; N-181 to P-196; Q-182 to P-196; L-183 to P-196; Y-184 to P-196; H-185 to P-196; L-186 to P-196; I-187 to P-196; Q-188 to P-196; N-189 to P-196; W-190 to P-196; P-191 to P-196; of SEQ ID NO:6. Polynucleotides encoding these polypeptides also are provided.

Further embodiments of the invention are directed to C-terminal deletions of the
 20 PGRP-K polypeptide described by the general formula 1 to n, where n is an integer from 7-242 corresponding to the position of amino acid residues identified in SEQ ID NO:2 and preferably, corresponds to one of the C-terminal amino acid residues identified in the C-terminal deletions specified herein. In specific embodiments, C terminal deletions of the PGRP-K polypeptide of the invention comprise, or
 25 alternatively, consist of, amino acid residues: M-1 to Y-242; M-1 to G-241; M-1 to I-240; M-1 to D-239; M-1 to C-238; M-1 to F-237; M-1 to N-236; M-1 to R-235; M-1 to T-234; M-1 to D-233; M-1 to M-232; M-1 to H-231; M-1 to F-230; M-1 to S-229; M-1 to Q-228; M-1 to I-227; M-1 to N-226; M-1 to R-225; M-1 to V-224; M-1 to V-223; M-1 to T-222; M-1 to Q-221; M-1 to C-220; M-1 to D-219; M-1 to T-218; M-1 to S-217;
 30 M-1 to V-216; M-1 to T-215; M-1 to C-214; M-1 to S-213; M-1 to T-212; M-1 to G-211; M-1 to A-210; M-1 to T-209; M-1 to H-208; M-1 to I-207; M-1 to I-206; M-1 to I-205; M-1 to V-204; M-1 to Y-203; M-1 to K-202; M-1 to A-201; M-1 to P-200; M-1 to L-199; M-1 to N-198; M-1 to M-197; M-1 to K-196; M-1 to P-195; M-1 to C-194; M-1 to H-193; M-1 to T-192; M-1 to E-191; M-1 to R-190; M-1 to A-189; M-1 to E-188; M-1 to W-187; M-1 to A-186; M-1 to S-185; M-1 to R-184; M-1 to K-183; M-1 to I-182;
 35 M-1 to I-181; M-1 to N-180; M-1 to P-179; M-1 to C-178; M-1 to V-177; M-1 to K-176; M-1 to R-175; M-1 to P-174; M-1 to M-173; M-1 to V-172; M-1 to P-171; M-1 to

H-170; M-1 to Q-169; M-1 to P-168; M-1 to D-167; M-1 to L-166; M-1 to C-165; M-1 to T-164; M-1 to E-163; M-1 to E-162; M-1 to K-161; M-1 to L-160; M-1 to L-159; M-1 to L-158; M-1 to P-157; M-1 to Q-156; M-1 to I-155; M-1 to Y-154; M-1 to R-153; M-1 to P-152; M-1 to S-151; M-1 to L-150; M-1 to H-149; M-1 to G-148; M-1 to K-147; M-1 to Q-146; M-1 to I-145; M-1 to A-144; M-1 to Y-143; M-1 to S-142; M-1 to I-141; M-1 to L-140; M-1 to G-139; M-1 to E-138; M-1 to A-137; M-1 to A-136; M-1 to S-135; M-1 to L-134; M-1 to A-133; M-1 to A-132; M-1 to P-131; M-1 to S-130; M-1 to P-129; M-1 to S-128; M-1 to S-127; M-1 to S-126; M-1 to I-125; M-1 to K-124; M-1 to N-123; M-1 to G-122; M-1 to F-121; M-1 to F-120; M-1 to A-119; M-1 to I-118; M-1 to G-117; M-1 to L-116; M-1 to S-115; M-1 to I-114; M-1 to N-113; M-1 to N-112; M-1 to Y-111; M-1 to G-110; M-1 to Q-109; M-1 to T-108; M-1 to H-107; M-1 to L-106; M-1 to G-105; M-1 to Q-104; M-1 to I-103; M-1 to N-102; M-1 to W-101; M-1 to G-100; M-1 to V-99; M-1 to G-98; M-1 to E-97; M-1 to Y-96; M-1 to V-95; M-1 to R-94; M-1 to G-93; M-1 to D-92; M-1 to D-91; M-1 to G-90; M-1 to V-89; M-1 to L-88; M-1 to F-87; M-1 to N-86; M-1 to Y-85; M-1 to A-84; M-1 to V-83; M-1 to D-82; M-1 to C-81; M-1 to W-80; M-1 to G-79; M-1 to I-78; M-1 to T-77; M-1 to Y-76; M-1 to V-75; M-1 to S-74; M-1 to H-73; M-1 to S-72; M-1 to Q-71; M-1 to L-70; M-1 to G-69; M-1 to R-68; M-1 to L-67; M-1 to M-66; M-1 to Q-65; M-1 to S-64; M-1 to C-63; M-1 to V-62; M-1 to S-61; M-1 to Q-60; M-1 to Q-59; M-1 to Q-58; M-1 to C-57; M-1 to Q-56; M-1 to M-55; M-1 to G-54; M-1 to P-53; M-1 to L-52; M-1 to Q-51; M-1 to D-50; M-1 to T-49; M-1 to I-48; M-1 to I-47; M-1 to Y-46; M-1 to A-45; M-1 to V-44; M-1 to P-43; M-1 to L-42; M-1 to T-41; M-1 to L-40; M-1 to L-39; M-1 to A-38; M-1 to R-37; M-1 to C-36; M-1 to A-35; M-1 to L-34; M-1 to P-33; M-1 to R-32; M-1 to A-31; M-1 to G-30; M-1 to W-29; M-1 to E-28; M-1 to K-27; M-1 to R-26; M-1 to S-25; M-1 to V-24; M-1 to I-23; M-1 to T-22; M-1 to P-21; M-1 to T-20; M-1 to D-19; M-1 to W-18; M-1 to A-17; M-1 to Q-16; M-1 to L-15; M-1 to G-14; M-1 to L-13; M-1 to I-12; M-1 to F-11; M-1 to F-10; M-1 to A-9; M-1 to L-8; M-1 to L-7; of SEQ ID NO:2. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Further embodiments of the invention are directed to C-terminal deletions of the PGRP-W polypeptide described by the general formula 1 to n, where n is an integer from 7-367 corresponding to the position of amino acid residues identified in SEQ ID NO:4 and preferably, corresponds to one of the C-terminal amino acid residues identified in the C-terminal deletions specified herein. In specific embodiments, C terminal deletions of the PGRP-W polypeptide of the invention comprise, or alternatively, consist of, amino acid residues: M-1 to K-367; M-1 to F-366; M-1 to H-365; M-1 to P-364; M-1 to W-363; M-1 to T-362; M-1 to S-361; M-1 to I-360; M-1 to

I-359; M-1 to N-358; M-1 to Y-357; M-1 to L-356; M-1 to A-355; M-1 to Q-354; M-1 to G-353; M-1 to P-352; M-1 to S-351; M-1 to L-350; M-1 to T-349; M-1 to R-348; M-1 to A-347; M-1 to V-346; M-1 to D-345; M-1 to S-344; M-1 to H-343; M-1 to G-342; M-1 to V-341; M-1 to L-340; M-1 to L-339; M-1 to Y-338; M-1 to N-337; M-1 to P-336; M-1 to T-335; M-1 to L-334; M-1 to Y-333; M-1 to G-332; M-1 to K-331; M-1 to V-330; M-1 to M-329; M-1 to A-328; M-1 to C-327; M-1 to Q-326; M-1 to I-325; M-1 to L-324; M-1 to D-323; M-1 to Q-322; M-1 to A-321; M-1 to A-320; M-1 to E-319; M-1 to L-318; M-1 to A-317; M-1 to A-316; M-1 to A-315; M-1 to N-314; M-1 to P-313; M-1 to P-312; M-1 to I-311; M-1 to G-310; M-1 to T-309; M-1 to F-308; M-1 to T-307; M-1 to G-306; M-1 to M-305; M-1 to F-304; M-1 to T-303; M-1 to I-302; M-1 to G-301; M-1 to L-300; M-1 to A-299; M-1 to I-298; M-1 to D-297; M-1 to D-296; M-1 to Y-295; M-1 to G-294; M-1 to P-293; M-1 to T-292; M-1 to S-291; M-1 to S-290; M-1 to G-289; M-1 to Q-288; M-1 to V-287; M-1 to N-286; M-1 to W-285; M-1 to G-284; M-1 to V-283; M-1 to G-282; M-1 to E-281; M-1 to Y-280; M-1 to I-279; M-1 to A-278; M-1 to G-277; M-1 to D-276; M-1 to Q-275; M-1 to G-274; M-1 to V-273; M-1 to L-272; M-1 to F-271; M-1 to N-270; M-1 to Y-269; M-1 to G-268; M-1 to I-267; M-1 to D-266; M-1 to C-265; M-1 to S-264; M-1 to K-263; M-1 to L-262; M-1 to R-261; M-1 to D-260; M-1 to I-259; M-1 to Y-258; M-1 to F-257; M-1 to S-256; M-1 to Q-255; M-1 to I-254; M-1 to D-253; M-1 to R-252; M-1 to V-251; M-1 to L-250; M-1 to L-249; M-1 to R-248; M-1 to C-247; M-1 to E-246; M-1 to D-245; M-1 to S-244; M-1 to I-243; M-1 to N-242; M-1 to C-241; M-1 to T-240; M-1 to R-239; M-1 to G-238; M-1 to A-237; M-1 to T-236; M-1 to H-235; M-1 to I-234; M-1 to I-233; M-1 to I-232; M-1 to G-231; M-1 to Y-230; M-1 to K-229; M-1 to A-228; M-1 to P-227; M-1 to L-226; M-1 to T-225; M-1 to M-224; M-1 to R-223; M-1 to S-222; M-1 to L-221; M-1 to P-220; M-1 to R-219; M-1 to G-218; M-1 to P-217; M-1 to E-216; M-1 to G-215; M-1 to C-214; M-1 to L-213; M-1 to G-212; M-1 to H-211; M-1 to S-210; M-1 to L-209; M-1 to A-208; M-1 to P-207; M-1 to A-206; M-1 to L-205; M-1 to K-204; M-1 to K-203; M-1 to L-202; M-1 to S-201; M-1 to T-200; M-1 to K-199; M-1 to Q-198; M-1 to R-197; M-1 to P-196; M-1 to A-195; M-1 to L-194; M-1 to C-193; M-1 to N-192; M-1 to E-191; M-1 to G-190; M-1 to K-189; M-1 to G-188; M-1 to L-187; M-1 to L-186; M-1 to P-185; M-1 to Q-184; M-1 to V-183; M-1 to Y-182; M-1 to S-181; M-1 to S-180; M-1 to S-179; M-1 to L-178; M-1 to H-177; M-1 to G-176; M-1 to K-175; M-1 to Q-174; M-1 to V-173; M-1 to A-172; M-1 to Y-171; M-1 to T-170; M-1 to I-169; M-1 to L-168; M-1 to N-167; M-1 to E-166; M-1 to M-165; M-1 to A-164; M-1 to S-163; M-1 to L-162; M-1 to A-161; M-1 to A-160; M-1 to P-159; M-1 to S-158; M-1 to P-157; M-1 to S-156; M-1 to H-155; M-1 to G-154; M-1 to K-153; M-1 to K-152; M-1 to T-151; M-1 to G-150; M-1 to F-149; M-1 to F-148; M-1 to A-147; M-1 to F-146; M-1 to G-

145; M-1 to L-144; M-1 to S-143; M-1 to I-142; M-1 to N-141; M-1 to N-140; M-1 to Y-139; M-1 to G-138; M-1 to Q-137; M-1 to T-136; M-1 to H-135; M-1 to V-134; M-1 to G-133; M-1 to Q-132; M-1 to I-131; M-1 to N-130; M-1 to W-129; M-1 to G-128; M-1 to V-127; M-1 to G-126; M-1 to E-125; M-1 to Y-124; M-1 to V-123; M-1 to R-122; M-1 to G-121; M-1 to D-120; M-1 to D-119; M-1 to G-118; M-1 to V-117; M-1 to L-116; M-1 to F-115; M-1 to N-114; M-1 to Y-113; M-1 to A-112; M-1 to V-111; M-1 to D-110; M-1 to C-109; M-1 to G-108; M-1 to S-107; M-1 to N-106; M-1 to N-105; M-1 to H-104; M-1 to V-103; M-1 to H-102; M-1 to H-101; M-1 to A-100; M-1 to Q-99; M-1 to L-98; M-1 to E-97; M-1 to R-96; M-1 to L-95; M-1 to R-94; M-1 to Q-93; M-1 to S-92; M-1 to C-91; M-1 to V-90; M-1 to T-89; M-1 to Q-88; M-1 to D-87; M-1 to H-86; M-1 to C-85; M-1 to E-84; M-1 to L-83; M-1 to G-82; M-1 to P-81; M-1 to V-80; M-1 to H-79; M-1 to H-78; M-1 to I-77; M-1 to V-76; M-1 to L-75; M-1 to V-74; M-1 to N-73; M-1 to V-72; M-1 to P-71; M-1 to T-70; M-1 to T-69; M-1 to L-68; M-1 to Q-67; M-1 to I-66; M-1 to S-65; M-1 to C-64; M-1 to G-63; M-1 to V-62; M-1 to A-61; M-1 to E-60; M-1 to A-59; M-1 to G-58; M-1 to W-57; M-1 to A-56; M-1 to K-55; M-1 to R-54; M-1 to S-53; M-1 to V-52; M-1 to T-51; M-1 to T-50; M-1 to S-49; M-1 to V-48; M-1 to D-47; M-1 to K-46; M-1 to E-45; M-1 to T-44; M-1 to L-43; M-1 to Q-42; M-1 to S-41; M-1 to I-40; M-1 to N-39; M-1 to E-38; M-1 to F-37; M-1 to L-36; M-1 to Y-35; M-1 to Q-34; M-1 to L-33; M-1 to G-32; M-1 to E-31; M-1 to S-30; M-1 to V-29; M-1 to Q-28; M-1 to K-27; M-1 to A-26; M-1 to Q-25; M-1 to T-24; M-1 to K-23; M-1 to N-22; M-1 to W-21; M-1 to S-20; M-1 to S-19; M-1 to D-18; M-1 to G-17; M-1 to W-16; M-1 to A-15; M-1 to Q-14; M-1 to I-13; M-1 to G-12; M-1 to L-11; M-1 to A-10; M-1 to S-9; M-1 to F-8; M-1 to V-7; of SEQ ID NO:4. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Further embodiments of the invention are directed to C-terminal deletions of the PGRP-C polypeptide described by the general formula 1 to n, where n is an integer from 7-196 corresponding to the position of amino acid residues identified in SEQ ID NO:6 and preferably, corresponds to one of the C-terminal amino acid residues identified in the C-terminal deletions specified herein. In specific embodiments, C terminal deletions of the PGRP-C polypeptide of the invention comprise, or alternatively, consist of, amino acid residues: M-1 to S-195; M-1 to R-194; M-1 to Y-193; M-1 to H-192; M-1 to P-191; M-1 to W-190; M-1 to N-189; M-1 to Q-188; M-1 to I-187; M-1 to L-186; M-1 to H-185; M-1 to Y-184; M-1 to L-183; M-1 to Q-182; M-1 to N-181; M-1 to G-180; M-1 to P-179; M-1 to S-178; M-1 to L-177; M-1 to T-176; M-1 to R-175; M-1 to Q-174; M-1 to V-173; M-1 to D-172; M-1 to R-171; M-1 to H-170; M-1 to G-169; M-1 to K-168; M-1 to L-167; M-1 to V-166; M-1 to Y-165; M-1 to N-164; M-1 to S-163; M-1 to R-162; M-1 to L-161; M-1 to A-160; M-1 to G-159; M-1 to

Q-158; M-1 to A-157; M-1 to V-156; M-1 to G-155; M-1 to C-154; M-1 to A-153; M-1 to L-152; M-1 to L-151; M-1 to G-150; M-1 to Q-149; M-1 to A-148; M-1 to A-147; M-1 to R-146; M-1 to I-145; M-1 to A-144; M-1 to Q-143; M-1 to P-142; M-1 to T-141; M-1 to P-140; M-1 to V-139; M-1 to R-138; M-1 to D-137; M-1 to M-136; M-1 to Y-135; M-1 to N-134; M-1 to G-133; M-1 to M-132; M-1 to F-131; M-1 to S-130; M-1 to I-129; M-1 to G-128; M-1 to I-127; M-1 to S-126; M-1 to M-125; M-1 to P-124; M-1 to N-123; M-1 to W-122; M-1 to L-121; M-1 to H-120; M-1 to G-119; M-1 to S-118; M-1 to H-117; M-1 to A-116; M-1 to G-115; M-1 to T-114; M-1 to F-113; M-1 to N-112; M-1 to W-111; M-1 to G-110; M-1 to R-109; M-1 to G-108; M-1 to E-107; M-1 to Y-106; M-1 to V-105; M-1 to L-104; M-1 to G-103; M-1 to D-102; M-1 to E-101; M-1 to G-100; M-1 to I-99; M-1 to L-98; M-1 to F-97; M-1 to N-96; M-1 to Y-95; M-1 to G-94; M-1 to V-93; M-1 to D-92; M-1 to C-91; M-1 to W-90; M-1 to G-89; M-1 to L-88; M-1 to T-87; M-1 to K-86; M-1 to M-85; M-1 to H-84; M-1 to Y-83; M-1 to H-82; M-1 to Q-81; M-1 to V-80; M-1 to N-79; M-1 to R-78; M-1 to A-77; M-1 to Q-76; M-1 to Q-75; M-1 to Q-74; M-1 to C-73; M-1 to S-72; M-1 to A-71; M-1 to P-70; M-1 to T-69; M-1 to N-68; M-1 to C-67; M-1 to S-66; M-1 to S-65; M-1 to G-64; M-1 to A-63; M-1 to T-62; M-1 to H-61; M-1 to S-60; M-1 to V-59; M-1 to V-58; M-1 to V-57; M-1 to Y-56; M-1 to R-55; M-1 to L-54; M-1 to P-53; M-1 to L-52; M-1 to S-51; M-1 to L-50; M-1 to H-49; M-1 to Q-48; M-1 to A-47; M-1 to C-46; M-1 to E-45; M-1 to S-44; M-1 to A-43; M-1 to L-42; M-1 to A-41; M-1 to K-40; M-1 to W-39; M-1 to E-38; M-1 to N-37; M-1 to R-36; M-1 to P-35; M-1 to V-34; M-1 to I-33; M-1 to P-32; M-1 to S-31; M-1 to C-30; M-1 to C-29; M-1 to A-28; M-1 to P-27; M-1 to D-26; M-1 to E-25; M-1 to T-24; M-1 to E-23; M-1 to Q-22; M-1 to A-21; M-1 to A-20; M-1 to G-19; M-1 to L-18; M-1 to R-17; M-1 to L-16; M-1 to L-15; M-1 to S-14; M-1 to P-13; M-1 to L-12; M-1 to A-11; M-1 to W-10; M-1 to A-9; M-1 to L-8; M-1 to L-7; of SEQ ID NO:6. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Further embodiments of the invention are directed to polypeptide fragments comprising, or alternatively, consisting of, amino acids described by the general formula m to n, where m and n are integers corresponding to any one of the amino acid

residues specified above for these symbols, respectively. Polynucleotides encoding such polypeptides are also provided.

It will be recognized in the art that some amino acid sequences of the PGRP-K, the PGRP-W, and/or the PGRP-C proteins can be varied without significant effect to the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on each protein which determine activity. Thus, the invention further includes variations of the PGRP-K protein, the PGRP-W protein, and/or the PGRP-C protein which show substantial PGRP-like activity or which include regions of either the PGRP-K protein, the PGRP-W protein, and/or the PGRP-C protein such as the polypeptide portions discussed below. Such mutants include deletions, insertions, inversions, repeats, and type substitutions. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells, *Science* 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

The resulting constructs can be routinely screened for activities or functions described throughout the specification and known in the art. Preferably, the resulting constructs have an increased and/or a decreased PGRP activity or function, while the remaining activities or functions are maintained. More preferably, the resulting constructs have more than one increased and/or decreased PGRP activity or function, while the remaining activities or functions are maintained.

Besides conservative amino acid substitution, variants of PGRP-K, PGRP-W, or PGRP-C include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as, for example, an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, PGRP-K, PGRP-W, or PGRP-C polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

The resulting constructs can be routinely screened for activities or functions described throughout the specification and known in the art. Preferably, the resulting constructs have an increased and/or decreased PGRP activity or function, while the remaining activities or functions are maintained. More preferably, the resulting constructs have more than one increased and/or decreased PGRP activity or function, while the remaining activities or functions are maintained.

Additionally, more than one amino acid (e.g., 2, 3, 4, 5, 6, 7, 8, 9 and 10) can be replaced with the substituted amino acids as described above (either conservative or nonconservative). The substituted amino acids can occur in the full length, mature, or proprotein form of the PGRP-K, PGRP-W, or PGRP-C proteins, respectively, as well as the N- and C- terminal deletion mutants, having the general formula m-n, as discussed above.

A further embodiment of the invention relates to a polypeptide which comprises the amino acid sequence of a PGRP-K, PGRP-W, or PGRP-C polypeptide having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course, in order of ever-increasing preference, it is highly preferable for a polypeptide to have an amino acid sequence which comprises the amino acid sequence of a PGRP-K, PGRP-W, or PGRP-C polypeptide, which contains at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions. In

specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of Figure 1 or fragments thereof (e.g., the mature form and/or other fragments described herein), is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

5 Thus, the fragment, derivative or analog of the polypeptides of Figures 1A-B (SEQ ID NO:2), Figures 2A-C (SEQ ID NO: 4), and/or Figure 3 (SEQ ID NO:6) or that are encoded by the deposited cDNAs, may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid
10 residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the either the PGRP-K, PGRP-W, and/or PGRP-C polypeptides are fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are
15 fused to the full length polypeptide, such as an IgG Fc fusion region peptide or leader or secretory sequence or a sequence which is employed for purification of the either the PGRP-K, PGRP-W, and/or PGRP-C polypeptides or proprotein sequences. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

20 Of particular interest are substitutions of charged amino acids with another charged amino acid and with neutral or negatively charged amino acids. The latter results in polypeptides with reduced positive charge to improve the characteristics of either the PGRP-K, PGRP-W, and/or PGRP-C polypeptides. The prevention of aggregation is highly desirable. Aggregation of polypeptides not only results in a loss
25 of activity but can also be problematic when preparing pharmaceutical formulations, because they can be immunogenic. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36:838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993)).

The replacement of amino acids can also change the selectivity of binding to

cell surface receptors. For example, Ostade et al. (Nature 361:266-268 (1993)) describes certain mutations resulting in selective binding of TNF- α to only one of the two known types of TNF receptors. Thus, the PGRP-K, PGRP-W, and/or PGRP-C proteins of the present invention may include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation.

As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table 1).

TABLE 1. Conservative Amino Acid Substitutions.

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

10

In specific embodiments, the number of substitutions, additions or deletions in the amino acid sequence of Figures 1A-B (SEQ ID NO:2), Figures 2A-C (SEQ ID NO:4), and/or Figure 3 (SEQ ID NO:6), and/or any of the polypeptide fragments described herein (e.g., PGRP-like domain) is 100, 90, 80, 75, 70, 60, 50, 40, 35, 30, 25,

20, 15, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 or 150-100, 100-50, 50-20, 20-10, 5-10, 1-5, 1-3 or 1-2.

Amino acids in the PGRP-K, PGRP-W, and/or PGRP-C polypeptides of the present invention that are essential for function can be identified by methods known
5 in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as receptor binding *in vitro*. Sites that are critical for ligand-receptor binding can also be determined by
10 structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith et al., J. Mol. Biol. 224:899-904 (1992) and de Vos et al., Science 255:306-312 (1992)).

The polypeptides of the present invention also include the polypeptides encoded by the deposited cDNAs; the polypeptide of Figures 1A-B (SEQ ID NO:2),
15 the polypeptides of Figures 2A-C (SEQ ID NO:4), and/or the polypeptides of Figure 3 (SEQ ID NO:6); the polypeptide sequence of any of the PGRP-like domains described herein; the polypeptide sequences of Figures 1A-B (SEQ ID NO:2), Figures 2A-C, and/or Figure 3 (SEQ ID NO:6), minus a portion, or all of, one or more of the PGRP-like domains described supra; and polypeptides which are at least 80%
20 identical, more preferably at least 85%, 90% or 95% identical, still more preferably at least 96%, 97%, 98% or 99% identical to the polypeptides described above, and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

By a polypeptide having an amino acid sequence at least, for example, 95%
25 "identical" to a reference amino acid sequence of either a PGRP-K, PGRP-W, and/or PGRP-C polypeptide is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acids of either the PGRP-K, PGRP-W, and/or PGRP-C proteins. In other words, to obtain

a polypeptide having an amino acid sequence at least 95% identical to one of the reference amino acid sequences, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences shown in Figures 1A-B (SEQ ID NO:2), Figures 2A-C (SEQ ID NO:4), and/or Figure 3 (SEQ ID NO:6), the amino acid sequence encoded by the deposited cDNA clones, respectively, or fragments thereof, can be determined conventionally using known computer programs such the MegAlign program, which is included in the suite of computer applications contained within the DNASTAR program. When using MegAlign or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

In a specific embodiment, the identity between a reference (query) sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, is determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245 (1990)). Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter. According to this embodiment, if the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual

correction is made to the results to take into consideration the fact that the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. A determination of whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of this embodiment. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence. For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are made for the purposes of this embodiment.

Polynucleotides encoding polypeptides that are 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to either the PGRP-K, PGRP-W, and/or PGRP-C polypeptides described herein are also provided.

Among the especially preferred fragments of the invention are fragments
5 characterized by structural or functional attributes of PGRP-K, PGRP-W, and/or PGRP-C. Such fragments include amino acid residues that comprise alpha-helix and alpha-helix forming regions ("alpha-regions"), beta-sheet and beta-sheet-forming regions ("beta-regions"), turn and turn-forming regions ("turn-regions"), coil and coil-forming regions ("coil-regions"), hydrophilic regions, hydrophobic regions, alpha
10 amphipathic regions, beta amphipathic regions, surface forming regions, and high antigenic index regions (i.e., containing four or more contiguous amino acids having an antigenic index of greater than or equal to 1.5, as identified using the default parameters of the Jameson-Wolf program) of complete (i.e., full-length) PGRP-K (SEQ ID NO:2), PGRP-W (SEQ ID NO:4), and/or PGRP-C (SEQ ID NO:6). Certain
15 preferred regions are those set out in Figure 5, Figure 6, or Figure 7, and include, but are not limited to, regions of the aforementioned types identified by analysis of the amino acid sequence depicted in Figures 1A-1B (SEQ ID NO:2), Figures 2A-2C (SEQ ID NO:4), and Figure 3 (SEQ ID NO:6), such preferred regions include; Garnier-Robson predicted alpha-regions, beta-regions, turn-regions, and coil-regions; Chou-
20 Fasman predicted alpha-regions, beta-regions, turn-regions, and coil-regions; Kyte-Doolittle predicted hydrophilic and hydrophobic regions; Eisenberg alpha and beta amphipathic regions; Emini surface-forming regions; and Jameson-Wolf high antigenic index regions, as predicted using the default parameters of these computer programs. Polynucleotides encoding these polypeptides are also encompassed by the invention.

25 In additional embodiments, the polynucleotides of the invention encode functional attributes of PGRP-K (SEQ ID NO:2), PGRP-W (SEQ ID NO:4), and/or PGRP-C (SEQ ID NO:6). Preferred embodiments of the invention in this regard include fragments that comprise alpha-helix and alpha-helix forming regions ("alpha-regions"), beta-sheet and beta-sheet forming regions ("beta-regions"), turn and

turn-forming regions ("turn-regions"), coil and coil-forming regions ("coil-regions"), hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions and high antigenic index regions of PGRP-K (SEQ ID NO:2), PGRP-W (SEQ ID NO:4), and/or PGRP-C (SEQ ID NO:6).

The data representing the structural or functional attributes of PGRP-K (SEQ ID NO:2) set forth in Figure 5 and/or Table II, as described above, was generated using the various modules and algorithms of the DNA*STAR set on default parameters. The data representing the structural or functional attributes of PGRP-W (SEQ ID NO:4) set forth in Figure 6 and/or Table III, as described above, was generated using the various modules and algorithms of the DNA*STAR set on default parameters. The data representing the structural or functional attributes of PGRP-C (SEQ ID NO:6) set forth in Figure 7 and/or Table IV, as described above, was generated using the various modules and algorithms of the DNA*STAR set on default parameters. In a preferred embodiment, the data presented in columns VIII, IX, XIII, and XIV of Tables II, III, and IV can be used to determine regions of PGRP-K, PGRP-W, and PGRP-C, respectively, which exhibit a high degree of potential for antigenicity. Regions of high antigenicity are determined from the data presented in columns VIII, IX, XIII, and/or IV by choosing values which represent regions of the polypeptide which are likely to be exposed on the surface of the polypeptide in an environment in which antigen recognition may occur in the process of initiation of an immune response.

Certain preferred regions in these regards are set out in Figures 5, 6, and 7, but may, as shown in Tables II, III, and IV, be represented or identified by using tabular representations of the data presented in Figures 5, 6, and 7, respectively. The DNA*STAR computer algorithm used to generate Figures 5, 6, and 7 (set on the original default parameters) was used to present the data in Figures 5, 6, and 7 in a tabular format (See Tables II, III, and IV). The tabular format of the data in Figures 5, 6, and 7 may be used to easily determine specific boundaries of a preferred region.

The above-mentioned preferred regions set out in Figures 5, 6, and 7, and in Tables II, III, and IV include, but are not limited to, regions of the aforementioned types identified by analysis of the amino acid sequence set out in Figures 1A-1B, Figures 2A-2C, and Figure 3. As set out in Figures 5, 6, and 7, and in Tables II, III, and IV, respectively, such preferred regions include Garnier-Robson alpha-regions, beta-regions, turn-regions, and coil-regions, Chou-Fasman alpha-regions, beta-regions, and coil-regions, Kyte-Doolittle hydrophilic regions and hydrophobic regions, Eisenberg alpha- and beta-amphipathic regions, Karplus-Schulz flexible regions, Emini surface-forming regions and Jameson-Wolf regions of high antigenic index.

Table II (PGRP-K):

	Res	Pos.	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	XIII	XIV
	Met	1	.	.	.	B	-0.33	.	.	.	-0.40	0.44
	Gly	2	.	.	.	B	-0.23	.	.	.	-0.40	0.53
5	Thr	3	.	.	.	B	B	.	.	.	-0.66	.	.	.	-0.60	0.43
	Leu	4	.	.	.	B	B	.	.	.	-1.08	.	.	.	-0.60	0.36
	Pro	5	.	.	.	B	B	.	.	.	-1.28	.	.	.	-0.60	0.30
	Trp	6	.	.	.	B	B	.	.	.	-1.38	.	.	.	-0.60	0.21
	Leu	7	.	.	.	B	B	.	.	.	-1.73	.	.	.	-0.60	0.22
10	Leu	8	.	.	.	B	B	.	.	.	-2.31	.	.	.	-0.60	0.12
	Ala	9	.	.	.	B	B	.	.	.	-2.31	.	.	.	-0.60	0.08
	Phe	10	.	.	.	B	B	.	.	.	-2.44	.	.	.	-0.60	0.08
	Phe	11	.	.	.	B	B	.	.	.	-2.97	.	.	.	-0.60	0.10
	Ile	12	.	.	.	B	B	.	.	.	-2.16	.	.	.	-0.60	0.08
15	Leu	13	.	.	.	B	B	.	.	.	-1.93	.	.	.	-0.60	0.16
	Gly	14	B	.	.	C	-1.63	.	.	.	-0.40	0.19
	Leu	15	B	.	.	C	-0.93	.	.	.	-0.40	0.28
	Gln	16	B	.	.	C	-0.54	.	.	.	-0.40	0.57
	Ala	17	B	T	.	.	0.13	.	.	.	-0.20	0.84
20	Trp	18	B	T	.	.	0.63	.	.	.	-0.05	1.57
	Asp	19	B	.	.	C	0.09	.	.	F	0.20	1.31
	Thr	20	.	.	.	B	B	.	.	.	0.04	.	.	F	-0.45	0.91
	Pro	21	.	.	.	B	B	.	.	.	-0.26	*	.	F	-0.45	0.64
	Thr	22	.	.	.	B	B	.	.	.	0.44	*	.	F	-0.15	0.51
25	Ile	23	.	.	.	B	B	.	.	.	0.78	.	.	.	-0.30	0.70
	Val	24	.	.	.	B	B	.	.	.	0.78	.	.	.	0.30	0.90
	Ser	25	.	.	A	B	0.80	.	.	F	0.90	1.08
	Arg	26	.	.	A	B	0.67	.	.	F	0.60	1.63
	Lys	27	.	.	A	B	0.39	.	*	F	0.90	2.17
30	Glu	28	.	.	A	.	.	.	T	.	1.39	.	*	F	1.30	1.63
	Trp	29	.	.	A	.	.	.	T	.	2.03	.	*	.	1.15	1.63
	Gly	30	.	.	A	.	.	.	T	.	1.52	.	*	.	1.15	1.26
	Ala	31	.	.	A	.	.	.	T	.	0.82	.	*	F	0.85	0.60
	Arg	32	.	.	A	C	0.11	.	.	F	-0.25	0.58
35	Pro	33	.	.	A	C	0.22	*	*	.	-0.10	0.31
	Leu	34	.	.	A	.	.	.	T	.	-0.08	*	.	.	0.70	0.61
	Ala	35	.	.	A	B	-0.54	.	*	.	0.30	0.31
	Cys	36	.	.	A	B	-0.77	.	*	.	-0.30	0.17
	Arg	37	.	.	A	B	B	.	.	.	-1.19	*	*	.	-0.60	0.17
40	Ala	38	.	.	A	B	B	.	.	.	-1.79	*	*	.	-0.60	0.24
	Leu	39	.	.	A	B	B	.	.	.	-1.19	*	*	.	-0.60	0.37
	Leu	40	.	.	A	B	B	.	.	.	-1.46	*	*	.	-0.60	0.29
	Thr	41	.	.	A	B	B	.	.	.	-1.38	*	*	.	-0.60	0.21
	Leu	42	.	.	A	B	B	.	.	.	-1.73	*	*	.	-0.60	0.26
45	Pro	43	.	.	A	B	B	.	.	.	-2.03	.	.	.	-0.60	0.50
	Val	44	.	.	.	B	B	.	.	.	-2.11	.	.	.	-0.60	0.24
	Ala	45	.	.	.	B	B	.	.	.	-1.61	.	.	.	-0.60	0.20
	Tyr	46	.	.	.	B	B	.	.	.	-1.30	.	.	.	-0.60	0.19
	Ile	47	.	.	.	B	B	.	.	.	-0.49	*	.	.	-0.60	0.43
50	Ile	48	.	.	.	B	B	.	.	.	-1.09	*	.	.	-0.60	0.74
	Thr	49	.	.	.	B	B	.	.	.	-0.44	*	.	.	-0.60	0.39
	Asp	50	.	.	.	B	B	.	.	.	-0.20	.	.	F	-0.11	0.86
	Gln	51	.	.	.	B	-0.56	.	.	F	0.28	1.21
	Leu	52	T	C	0.33	.	.	F	0.57	0.83
55	Pro	53	T	T	.	0.56	.	.	F	1.41	0.86
	Gly	54	T	T	.	0.87	.	.	.	0.40	0.27
	Met	55	T	T	.	0.87	.	.	.	0.36	0.56
	Gln	56	.	.	.	B	B	.	.	.	0.87	.	.	.	-0.18	0.63

	Cys	57	.	.	B	B	.	.	.	1.38	.	.	F	0.08	1.09
	Gln	58	.	.	B	B	.	.	.	0.73	.	.	F	0.04	1.48
	Gln	59	.	.	B	B	.	.	.	0.41	.	.	F	-0.15	0.63
	Gln	60	.	.	B	B	.	.	.	0.71	.	.	F	-0.15	0.63
5	Ser	61	.	.	B	B	.	.	.	0.71	.	.	F	-0.15	0.49
	Val	62	.	.	B	B	.	.	.	0.78	.	.	F	-0.15	0.49
	Cys	63	.	.	B	B	.	.	.	-0.03	*	.	.	-0.60	0.28
	Ser	64	.	.	B	B	.	.	.	0.08	*	*	.	-0.60	0.17
	Gln	65	.	.	B	B	.	.	.	-0.27	*	.	.	-0.30	0.46
10	Met	66	.	.	B	B	.	.	.	-0.78	*	.	.	-0.30	0.84
	Leu	67	.	.	B	B	.	.	.	0.08	*	.	.	-0.30	0.52
	Arg	68	.	.	B	B	.	.	.	0.44	*	.	.	-0.30	0.52
	Gly	69	.	.	B	0.71	*	.	F	0.05	0.70
	Leu	70	C	0.41	*	.	F	0.40	1.16
15	Gln	71	.	.	B	.	.	T	C	0.16	*	.	F	1.05	0.79
	Ser	72	T	C	0.72	.	.	.	0.00	0.59
	His	73	.	.	B	.	.	T	.	0.30	.	.	.	-0.05	1.13
	Ser	74	.	.	B	.	.	T	.	-0.24	.	.	.	-0.20	0.94
	Val	75	.	.	B	B	.	.	.	0.22	.	.	.	-0.60	0.49
20	Tyr	76	.	.	B	B	.	.	.	-0.07	.	.	.	-0.60	0.36
	Thr	77	.	.	B	B	.	.	.	-0.43	.	.	.	-0.60	0.28
	Ile	78	.	.	B	B	.	.	.	-0.40	.	.	.	-0.60	0.20
	Gly	79	.	.	B	B	.	.	.	-0.96	.	.	.	-0.60	0.22
	Trp	80	.	.	B	B	.	.	.	-0.69	.	.	.	-0.60	0.11
25	Cys	81	.	.	B	B	.	.	.	-0.69	.	.	.	-0.60	0.16
	Asp	82	.	.	B	B	.	.	.	-0.38	.	*	.	-0.60	0.25
	Val	83	.	.	B	B	.	.	.	-0.19	*	.	.	-0.60	0.39
	Ala	84	.	.	B	B	.	.	.	-0.66	*	*	.	-0.60	0.63
	Tyr	85	.	.	B	B	.	.	.	-1.22	*	*	.	-0.60	0.31
30	Asn	86	.	.	B	B	.	.	.	-0.90	.	*	.	-0.60	0.31
	Phe	87	.	.	B	B	.	.	.	-0.90	.	*	.	-0.29	0.30
	Leu	88	.	.	B	B	.	.	.	-0.04	.	.	.	0.02	0.32
	Val	89	.	.	B	B	.	.	.	0.20	.	*	.	1.23	0.34
	Gly	90	T	T	.	0.56	.	*	F	2.49	0.38
35	Asp	91	T	T	.	-0.30	.	*	F	3.10	0.91
	Asp	92	T	C	0.16	*	*	F	2.59	0.91
	Gly	93	T	C	0.97	*	*	F	2.43	1.44
	Arg	94	.	.	B	1.48	*	*	F	1.72	1.49
	Val	95	.	.	B	0.97	*	*	F	1.26	0.89
40	Tyr	96	.	.	B	0.62	*	*	.	0.50	0.66
	Glu	97	.	.	B	0.33	.	*	.	0.50	0.34
	Gly	98	T	.	.	0.68	*	*	.	0.00	0.48
	Val	99	T	.	.	-0.32	*	*	.	0.00	0.49
	Gly	100	T	.	.	0.53	*	*	.	0.00	0.20
45	Trp	101	C	0.43	*	*	.	-0.20	0.35
	Asn	102	.	.	B	-0.38	*	*	.	-0.40	0.46
	Ile	103	.	.	B	-0.07	.	*	.	-0.40	0.38
	Gln	104	.	.	B	0.48	.	*	.	-0.40	0.50
	Gly	105	.	.	B	0.82	.	*	F	-0.25	0.45
50	Leu	106	.	.	B	0.77	.	*	F	-0.10	1.10
	His	107	.	.	B	0.52	*	.	F	0.05	0.63
	Thr	108	T	C	1.41	*	.	F	0.15	1.00
	Gln	109	T	T	.	1.41	.	*	F	0.50	1.94
	Gly	110	T	T	.	0.87	.	*	F	0.80	2.30
55	Tyr	111	T	T	.	1.38	.	*	F	0.50	1.12
	Asn	112	C	0.60	.	*	F	-0.05	0.86
	Asn	113	.	.	B	B	.	.	.	0.57	*	*	.	-0.60	0.72
	Ile	114	.	.	B	B	.	.	.	-0.32	*	*	.	-0.60	0.45
	Ser	115	.	.	B	B	.	.	.	-0.57	.	*	.	-0.60	0.20

	Leu	116	.	.	B	B	.	.	.	-1.02	.	*	.	-0.60	0.12
	Gly	117	.	.	B	B	.	.	.	-1.72	.	*	.	-0.60	0.15
	Ile	118	.	.	B	B	.	.	.	-2.07	.	*	.	-0.60	0.10
	Ala	119	.	.	B	B	.	.	.	-1.18	.	*	.	-0.60	0.12
5	Phe	120	.	.	B	B	.	.	.	-0.83	.	*	.	-0.60	0.19
	Phe	121	.	.	B	.	.	T	.	-0.91	.	*	.	-0.20	0.55
	Gly	122	.	.	B	.	.	T	.	-0.87	*	.	F	0.19	0.38
	Asn	123	T	T	.	-0.28	*	.	F	0.83	0.59
	Lys	124	T	T	.	0.01	.	.	F	1.37	0.92
10	Ile	125	T	.	.	0.50	*	.	F	2.16	1.24
	Ser	126	T	.	.	0.90	*	*	F	2.40	1.20
	Ser	127	C	1.03	*	.	F	1.81	0.80
	Ser	128	T	C	0.44	*	.	F	1.32	1.77
	Pro	129	T	C	-0.19	.	*	F	1.68	1.33
15	Ser	130	T	C	-0.11	.	.	F	0.84	1.00
	Pro	131	T	C	-0.11	.	.	F	0.45	0.62
	Ala	132	.	A	B	-0.40	.	.	.	-0.30	0.54
	Ala	133	.	A	B	-0.69	.	.	.	-0.30	0.40
	Leu	134	.	A	B	-0.48	.	.	.	-0.60	0.26
20	Ser	135	.	A	B	-0.52	.	.	.	-0.30	0.45
	Ala	136	.	A	B	-1.12	.	.	.	-0.30	0.44
	Ala	137	.	A	B	-1.42	*	.	.	-0.30	0.44
	Glu	138	A	A	B	-1.13	*	.	.	-0.30	0.23
	Gly	139	.	.	B	-0.57	*	.	.	-0.10	0.31
25	Leu	140	.	.	B	-0.86	*	.	.	-0.40	0.48
	Ile	141	.	.	B	-1.16	*	*	.	-0.40	0.28
	Ser	142	.	.	B	-0.57	*	*	.	-0.40	0.20
	Tyr	143	.	.	B	-0.52	*	.	.	-0.40	0.41
	Ala	144	.	.	B	-0.52	*	.	.	-0.25	1.18
30	Ile	145	.	.	B	0.26	*	.	.	-0.10	0.87
	Gln	146	.	.	B	.	.	T	.	0.33	*	*	.	0.36	0.76
	Lys	147	T	T	.	0.33	.	.	F	1.17	0.62
	Gly	148	T	T	.	0.37	*	*	F	1.58	1.18
	His	149	T	T	.	1.07	*	*	F	2.44	1.05
35	Leu	150	C	1.71	*	*	F	2.60	1.03
	Ser	151	T	C	0.82	*	*	F	1.64	1.64
	Pro	152	T	T	.	0.78	*	*	F	1.13	0.84
	Arg	153	T	T	.	0.91	*	*	.	1.17	1.77
	Tyr	154	.	.	B	.	.	T	.	0.13	*	*	.	0.51	2.04
40	Ile	155	.	.	B	0.13	*	.	.	0.05	1.09
	Gln	156	.	A	B	-0.38	*	.	.	-0.60	0.46
	Pro	157	.	A	B	-0.12	*	.	.	-0.60	0.24
	Leu	158	.	A	B	-0.23	*	.	.	-0.60	0.69
	Leu	159	.	A	B	0.01	.	.	.	0.30	0.69
45	Leu	160	.	A	B	0.59	.	.	F	0.75	0.77
	Lys	161	.	A	B	-0.08	.	.	F	0.60	1.35
	Glu	162	.	A	B	-0.68	.	*	F	0.75	0.88
	Glu	163	.	A	B	0.13	.	*	F	0.75	0.88
	Thr	164	.	A	.	.	T	.	.	0.73	.	*	F	1.15	0.73
50	Cys	165	.	A	.	.	T	.	.	1.54	.	*	F	1.15	0.65
	Leu	166	.	A	.	.	T	.	.	1.47	.	*	F	0.85	0.65
	Asp	167	T	C	1.26	.	*	F	0.45	0.62
	Pro	168	T	T	.	0.40	.	*	F	0.80	1.78
	Gln	169	T	T	.	0.11	.	.	F	0.80	1.60
55	His	170	T	C	.	0.57	*	*	.	0.30	0.95
	Pro	171	.	.	B	1.49	.	*	.	-0.40	0.95
	Val	172	.	.	B	1.53	.	.	.	0.05	1.07
	Met	173	.	.	B	0.89	.	.	.	0.65	1.58
	Pro	174	.	.	B	0.22	.	.	.	0.50	0.76

	Arg	175	T	.	.	0.04	.	.	F	1.05	0.55
	Lys	176	.	.	B	0.26	*	*	F	0.65	0.85
	Val	177	.	.	B	0.22	*	.	.	0.97	0.89
	Cys	178	.	.	B	.	.	T	.	-0.07	*	.	.	1.04	0.32
5	Pro	179	.	.	B	.	.	T	.	0.19	*	.	.	0.31	0.11
	Asn	180	.	.	B	.	.	T	.	0.19	*	.	.	0.48	0.30
	Ile	181	.	.	B	.	.	T	.	-0.16	*	*	.	1.70	1.10
	Ile	182	.	.	B	0.11	*	.	.	1.18	0.95
	Lys	183	.	A	B	0.49	*	.	.	0.81	0.60
10	Arg	184	.	A	B	0.70	*	.	F	0.19	0.90
	Ser	185	.	A	C	0.11	*	*	.	1.12	2.21
	Ala	186	.	A	C	1.11	*	*	.	0.95	1.12
	Trp	187	.	A	C	2.00	*	*	.	0.95	1.12
	Glu	188	.	A	C	1.64	*	*	.	0.95	1.45
15	Ala	189	A	A	1.50	.	*	F	0.90	2.06
	Arg	190	.	A	.	.	T	.	.	1.13	.	*	F	1.30	2.67
	Glu	191	.	A	.	.	T	.	.	1.51	.	*	F	1.15	0.83
	Thr	192	.	A	.	.	T	.	.	1.84	.	*	F	1.30	1.27
	His	193	.	A	.	.	T	.	.	1.24	.	*	F	1.30	1.29
20	Cys	194	T	C	1.83	.	*	.	0.90	0.74
	Pro	195	T	T	.	0.91	.	*	.	0.50	0.82
	Lys	196	T	T	.	0.70	.	*	.	0.50	0.50
	Met	197	T	T	.	0.42	.	*	.	0.65	1.44
	Asn	198	C	0.50	.	*	.	0.10	0.94
25	Leu	199	.	.	B	0.92	.	*	.	0.50	0.94
	Pro	200	.	.	B	0.28	.	*	.	0.05	1.49
	Ala	201	.	.	.	B	T	.	.	-0.66	.	*	.	-0.20	0.69
	Lys	202	.	.	B	B	.	.	.	-0.94	.	*	.	-0.60	0.58
	Tyr	203	.	.	B	B	.	.	.	-1.83	.	*	.	-0.60	0.26
30	Val	204	.	.	B	B	.	.	.	-1.06	.	*	.	-0.60	0.18
	Ile	205	.	.	B	B	.	.	.	-1.16	.	.	.	-0.60	0.12
	Ile	206	.	.	B	B	.	.	.	-1.16	.	.	.	-0.60	0.12
	Ile	207	.	.	B	B	.	.	.	-1.54	.	.	.	-0.60	0.16
	His	208	.	.	B	B	.	.	.	-1.61	.	.	.	-0.60	0.22
35	Thr	209	.	.	B	B	.	.	.	-1.06	.	.	.	-0.60	0.46
	Ala	210	.	.	.	B	.	.	.	-0.83	.	.	F	-0.05	0.87
	Gly	211	T	T	.	-0.26	.	.	F	0.65	0.34
	Thr	212	T	T	.	-0.22	.	.	F	0.65	0.34
	Ser	213	T	T	.	-0.49	.	.	F	0.35	0.25
40	Cys	214	.	.	B	.	.	T	.	-0.49	.	.	.	0.10	0.34
	Thr	215	.	.	B	0.10	.	*	.	-0.40	0.34
	Val	216	.	.	B	-0.22	.	*	F	0.82	0.43
	Ser	217	.	.	B	.	.	T	.	0.09	.	*	F	0.59	0.43
	Thr	218	.	.	B	.	.	T	.	0.08	.	*	F	1.36	0.51
45	Asp	219	T	T	.	-0.11	.	*	F	1.93	0.99
	Cys	220	.	.	B	-0.66	*	*	F	1.70	0.55
	Gln	221	.	.	B	B	.	.	.	0.31	*	*	F	0.53	0.28
	Thr	222	.	.	B	B	.	.	.	0.61	*	*	.	0.81	0.33
	Val	223	.	.	B	B	.	.	.	0.03	*	.	.	0.04	0.99
50	Val	224	.	.	B	B	.	.	.	0.03	*	.	.	-0.13	0.40
	Arg	225	.	.	B	B	.	.	.	0.40	*	.	.	-0.30	0.48
	Asn	226	.	.	B	B	.	.	.	-0.30	*	.	.	-0.30	0.87
	Ile	227	.	.	B	B	.	.	.	-0.02	*	.	F	0.00	1.02
	Gln	228	.	.	B	B	.	.	.	0.23	*	.	F	-0.15	0.71
55	Ser	229	.	.	B	B	.	.	.	1.09	*	*	.	-0.60	0.43
	Phe	230	.	.	B	B	.	.	.	0.67	*	*	.	0.13	1.04
	His	231	.	.	B	B	.	.	.	0.78	*	*	.	0.26	0.86
	Met	232	.	.	B	B	.	.	.	1.67	*	.	.	1.29	1.26
	Asp	233	T	T	.	0.97	*	*	.	2.37	2.34

	Thr	234	T	T	.	0.60	*	*	F	2.80	1.49
	Arg	235	T	T	.	1.30	*	*	F	2.37	0.81
	Asn	236	T	T	.	0.44	*	*	.	2.24	0.81
	Phe	237	T	.	.	0.70	*	*	.	1.46	0.39
5	Cys	238	T	.	.	0.46	*	*	.	1.18	0.20
	Asp	239	T	T	.	0.77	*	*	.	0.20	0.19
	Ile	240	T	T	.	0.27	*	*	.	0.20	0.39
	Gly	241	T	T	.	-0.12	*	.	.	0.50	0.92
	Tyr	242	.	.	B	.	.	T	.	0.19	*	.	.	0.10	0.71
10	Gln	243	.	.	B	0.47	.	*	.	-0.25	1.29

15

20

25

30

Table III (PGRP-W):

	Res	Pos. I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	XIII	XIV
5	Met	1	A	.	.	B	.	.	.	-1.38	.	.	.	-0.60	0.27
	Leu	2	A	.	.	B	.	.	.	-1.80	.	.	.	-0.60	0.17
	Leu	3	A	.	.	B	.	.	.	-2.27	.	.	.	-0.60	0.11
	Trp	4	A	.	.	B	.	.	.	-2.58	.	.	.	-0.60	0.08
	Leu	5	A	.	.	B	.	.	.	-2.49	.	.	.	-0.60	0.09
10	Leu	6	A	.	.	B	.	.	.	-2.48	.	.	.	-0.60	0.14
	Val	7	A	.	.	B	.	.	.	-2.48	.	.	.	-0.60	0.14
	Phe	8	A	.	.	B	.	.	.	-2.01	.	.	.	-0.60	0.14
	Ser	9	A	.	.	B	.	.	.	-2.61	.	.	.	-0.60	0.16
	Ala	10	A	.	B	B	.	.	.	-1.80	.	.	.	-0.60	0.15
15	Leu	11	.	.	B	B	.	.	.	-1.58	.	.	.	-0.60	0.31
	Gly	12	.	.	B	B	.	.	.	-1.01	.	.	.	-0.60	0.23
	Ile	13	.	.	B	B	.	.	.	-0.66	.	*	.	-0.60	0.24
	Gln	14	.	.	B	B	.	.	.	-0.36	.	*	.	-0.60	0.29
	Ala	15	.	.	B	B	.	.	.	-0.07	.	*	.	-0.30	0.49
20	Trp	16	.	.	B	B	.	.	.	0.44	.	*	.	-0.30	0.93
	Gly	17	.	.	B	.	.	.	C	0.50	*	*	F	0.33	0.72
	Asp	18	T	T	.	1.39	*	*	F	0.91	0.75
	Ser	19	T	C	1.43	.	.	F	1.14	1.15
	Ser	20	T	T	.	1.71	.	.	F	2.52	2.32
25	Trp	21	T	T	.	2.00	.	.	F	2.80	2.01
	Asn	22	T	C	1.76	.	*	F	2.32	2.60
	Lys	23	T	C	1.80	.	.	F	1.44	1.96
	Thr	24	A	T	.	2.10	.	.	F	1.56	3.72
	Gln	25	A	T	.	1.54	*	.	F	1.84	4.01
30	Ala	26	.	.	B	1.53	*	.	F	1.62	1.49
	Lys	27	.	.	B	1.53	*	.	F	1.58	1.38
	Gln	28	.	.	B	1.14	*	*	F	2.14	1.38
	Val	29	.	.	B	.	.	T	.	0.64	*	.	F	2.60	1.35
	Ser	30	.	.	B	.	.	T	.	0.64	*	.	F	1.89	0.56
35	Glu	31	.	.	B	.	.	T	.	0.99	*	.	F	1.63	0.56
	Gly	32	.	.	B	.	.	T	.	0.13	*	.	F	0.92	1.18
	Leu	33	.	A	B	-0.57	*	*	.	-0.04	0.73
	Gln	34	A	A	0.29	*	.	.	-0.60	0.36
	Tyr	35	.	A	B	0.59	*	.	.	-0.60	0.63
40	Leu	36	.	A	B	-0.30	*	.	.	-0.45	1.24
	Phe	37	.	A	B	-0.26	*	.	.	-0.60	0.50
	Glu	38	A	A	0.56	*	.	.	-0.60	0.43
	Asn	39	A	A	-0.26	*	.	F	-0.15	0.90
	Ile	40	A	A	-0.32	*	.	F	-0.15	0.86
45	Ser	41	A	A	0.49	*	.	F	-0.15	0.71
	Gln	42	A	A	1.23	*	.	F	-0.15	0.77
	Leu	43	A	A	1.23	*	.	F	0.60	2.19
	Thr	44	A	A	0.38	*	.	F	0.90	2.73
	Glu	45	A	A	0.97	*	.	F	0.90	1.17
50	Lys	46	A	A	0.96	.	.	F	0.90	1.90
	Asp	47	.	.	B	B	.	.	.	0.64	*	*	F	0.90	1.90
	Val	48	A	.	.	B	.	.	.	0.60	.	.	F	0.90	1.59
	Ser	49	.	.	B	B	.	.	.	0.61	*	.	F	0.45	0.59
	Thr	50	.	.	B	B	.	.	.	0.72	*	.	F	-0.15	0.47
55	Thr	51	.	.	B	B	.	.	.	0.72	*	.	F	0.00	1.25
	Val	52	.	.	B	B	.	.	.	0.13	*	*	F	0.90	1.86
	Ser	53	.	A	B	0.70	*	.	F	0.90	1.30
	Arg	54	.	A	B	0.66	*	.	F	0.45	0.95
	Lys	55	.	A	C	0.38	*	.	F	0.80	1.26

	Ala	56	.	A	.	.	T	.	C	0.69	.	.	0.70	0.95
	Trp	57	A	A	0.96	.	.	0.60	0.84
	Gly	58	A	A	0.40	.	.	0.30	0.43
	Ala	59	A	A	-0.06	.	.	-0.60	0.31
5	Glu	60	A	A	-0.77	.	.	-0.60	0.29
	Ala	61	A	.	.	B	.	.	.	-0.48	.	.	-0.30	0.16
	Val	62	A	.	.	B	.	.	.	-1.08	*	.	-0.30	0.21
	Gly	63	.	.	.	B	T	.	.	-0.73	*	.	0.10	0.09
	Cys	64	.	.	B	B	.	.	.	-0.96	.	*	-0.60	0.15
10	Ser	65	.	.	B	B	.	.	.	-1.27	.	*	-0.60	0.16
	Ile	66	.	.	B	B	.	.	.	-0.99	.	*	-0.60	0.24
	Gln	67	.	.	B	B	.	.	.	-0.34	.	*	-0.60	0.64
	Leu	68	.	.	B	B	.	.	.	-0.86	.	*	-0.60	0.74
	Thr	69	.	.	B	B	.	.	.	-0.19	.	*	-0.45	0.78
15	Thr	70	.	.	B	B	.	.	.	-0.74	.	*	-0.45	0.73
	Pro	71	.	.	B	B	.	.	.	-0.67	.	*	-0.45	0.65
	Val	72	.	.	B	B	.	.	.	-1.52	.	*	-0.60	0.37
	Asn	73	.	.	B	B	.	.	.	-1.60	.	*	-0.60	0.19
	Val	74	.	.	B	B	.	.	.	-1.32	.	.	-0.60	0.09
20	Leu	75	.	.	B	B	.	.	.	-1.04	.	.	-0.60	0.16
	Val	76	.	.	B	B	.	.	.	-1.69	.	.	-0.60	0.14
	Ile	77	.	.	B	B	.	.	.	-1.04	.	.	-0.60	0.14
	His	78	.	.	B	B	.	.	.	-1.39	.	.	-0.60	0.25
	His	79	.	.	B	B	.	.	.	-1.34	.	.	-0.60	0.34
25	Val	80	.	.	.	B	.	.	C	-0.53	.	.	-0.40	0.40
	Pro	81	.	.	.	B	.	.	C	-0.34	.	.	-0.10	0.51
	Gly	82	T	.	.	0.51	.	.	0.30	0.20
	Leu	83	T	.	.	0.54	.	.	0.30	0.37
	Glu	84	A	0.58	.	.	0.50	0.39
30	Cys	85	A	T	.	1.12	.	.	0.70	0.69
	His	86	A	T	.	0.48	.	.	0.85	1.21
	Asp	87	T	T	.	0.16	.	F	1.25	0.52
	Gln	88	A	T	.	0.67	.	F	0.25	0.52
	Thr	89	.	.	B	B	.	.	.	0.67	.	F	0.45	0.51
35	Val	90	.	.	B	B	.	.	.	1.44	.	*	0.45	0.53
	Cys	91	.	.	B	B	.	.	.	0.67	.	*	0.45	0.60
	Ser	92	.	.	B	B	.	.	.	0.78	.	*	-0.15	0.34
	Gln	93	.	.	B	B	.	.	.	0.78	*	*	0.45	0.90
	Arg	94	A	.	.	B	.	.	.	0.28	*	*	0.90	2.91
40	Leu	95	A	.	.	B	.	.	.	1.13	*	.	0.90	1.79
	Arg	96	A	.	.	B	.	.	.	1.21	*	*	0.90	1.79
	Glu	97	A	A	1.48	.	*	0.60	0.93
	Leu	98	A	A	1.44	*	*	0.45	1.53
	Gln	99	A	A	0.48	.	*	0.45	1.06
45	Ala	100	A	A	1.26	.	.	-0.30	0.45
	His	101	A	A	1.14	.	.	-0.60	0.75
	His	102	A	A	1.14	.	*	-0.60	0.70
	Val	103	.	A	B	1.66	.	.	-0.45	1.11
	His	104	.	A	.	.	T	.	.	1.31	.	.	0.50	1.09
50	Asn	105	.	A	.	.	T	.	.	1.23	.	F	0.75	0.79
	Asn	106	T	T	.	1.27	.	*	1.40	0.57
	Ser	107	T	T	.	0.44	.	F	2.25	0.70
	Gly	108	T	T	.	0.71	.	F	2.50	0.32
	Cys	109	T	T	.	0.50	.	.	2.10	0.20
55	Asp	110	.	.	B	B	.	.	.	0.50	*	*	0.45	0.24
	Val	111	.	.	B	B	.	.	.	-0.20	*	*	0.20	0.39
	Ala	112	.	.	B	B	.	.	.	-0.71	*	*	-0.35	0.63
	Tyr	113	.	.	B	B	.	.	.	-1.22	*	*	-0.60	0.31
	Asn	114	.	.	B	B	.	.	.	-0.90	.	*	-0.60	0.31

	Phe	115	.	.	B	B	.	.	.	-0.90	.	*	.	-0.29	0.30
	Leu	116	.	.	B	B	.	.	.	-0.04	.	.	.	0.02	0.32
	Val	117	.	.	B	B	.	.	.	0.20	.	*	.	1.23	0.34
	Gly	118	T	T	.	0.56	.	*	F	2.49	0.38
5	Asp	119	T	T	.	-0.30	.	*	F	3.10	0.91
	Asp	120	T	C	0.16	*	*	F	2.59	0.91
	Gly	121	T	C	0.97	*	*	F	2.43	1.44
	Arg	122	.	.	B	B	.	.	.	1.48	*	*	F	1.52	1.49
	Val	123	.	.	B	B	.	.	.	0.97	*	*	F	1.06	0.89
10	Tyr	124	.	.	B	B	.	.	.	0.62	*	*	.	0.30	0.66
	Glu	125	.	.	B	B	.	.	.	0.33	.	*	.	0.30	0.34
	Gly	126	.	.	.	B	T	.	.	0.68	*	*	.	-0.20	0.48
	Val	127	.	.	.	B	T	.	.	-0.32	*	*	.	-0.20	0.49
	Gly	128	.	.	.	B	T	.	.	0.53	*	*	.	-0.20	0.20
15	Trp	129	.	.	.	B	.	.	C	0.43	*	*	.	-0.40	0.35
	Asn	130	.	.	B	B	.	.	.	-0.42	*	*	.	-0.60	0.46
	Ile	131	.	.	B	B	.	.	.	-0.11	.	*	.	-0.60	0.35
	Gln	132	.	.	B	B	.	.	.	0.43	.	*	.	-0.60	0.45
	Gly	133	.	.	B	B	.	.	.	0.78	.	*	.	-0.60	0.40
20	Val	134	.	.	B	B	.	.	.	0.72	.	*	F	-0.45	0.99
	His	135	.	.	B	B	.	.	.	0.48	*	.	F	-0.15	0.57
	Thr	136	.	.	B	.	.	T	.	1.37	*	.	F	-0.05	0.90
	Gln	137	.	.	B	.	.	T	.	1.37	.	*	F	0.10	1.94
	Gly	138	T	T	.	0.82	.	*	F	0.80	2.30
25	Tyr	139	T	T	.	1.38	.	*	F	0.50	1.12
	Asn	140	.	.	B	0.60	.	*	F	-0.25	0.86
	Asn	141	.	.	B	B	.	.	.	0.57	*	*	.	-0.60	0.72
	Ile	142	.	.	B	B	.	.	.	-0.13	*	*	.	-0.60	0.45
	Ser	143	.	.	B	B	.	.	.	-0.38	.	*	.	-0.60	0.24
30	Leu	144	.	.	B	B	.	.	.	-0.83	.	*	.	-0.60	0.15
	Gly	145	.	.	B	B	.	.	.	-1.53	.	*	.	-0.60	0.19
	Phe	146	.	.	B	B	.	.	.	-1.88	.	*	.	-0.60	0.12
	Ala	147	.	.	B	B	.	.	.	-1.30	.	*	.	-0.60	0.15
	Phe	148	A	.	.	B	.	.	.	-0.96	.	.	.	-0.60	0.21
35	Phe	149	A	.	.	B	.	.	.	-0.10	.	.	.	-0.26	0.50
	Gly	150	A	-0.10	.	.	F	1.33	0.98
	Thr	151	T	T	.	0.57	.	.	F	2.42	1.12
	Lys	152	T	T	.	0.86	.	.	F	2.76	1.76
	Lys	153	T	T	.	1.34	.	.	F	3.40	2.39
40	Gly	154	T	T	.	1.74	.	.	F	3.06	2.56
	His	155	C	1.88	.	.	F	2.32	1.71
	Ser	156	T	C	1.60	.	.	F	1.88	1.33
	Pro	157	T	C	0.97	.	.	F	0.94	1.35
	Ser	158	T	C	0.11	.	.	F	0.60	1.00
45	Pro	159	A	T	.	0.16	.	.	F	0.25	0.62
	Ala	160	A	A	-0.40	.	.	.	-0.30	0.54
	Ala	161	A	A	-0.70	.	.	.	-0.30	0.40
	Leu	162	A	A	-0.49	.	.	.	-0.60	0.26
	Ser	163	A	A	-0.19	*	.	.	-0.30	0.44
50	Ala	164	A	A	-0.79	*	.	.	-0.30	0.71
	Met	165	A	A	-1.09	*	.	.	-0.30	0.71
	Glu	166	A	A	-0.81	*	.	.	-0.30	0.37
	Asn	167	A	.	.	B	.	.	.	-0.24	*	.	.	-0.30	0.53
	Leu	168	A	.	.	B	.	.	.	-0.53	*	.	.	-0.60	0.83
55	Ile	169	A	.	.	B	.	.	.	-0.80	*	.	.	-0.60	0.49
	Thr	170	A	.	.	B	.	.	.	-0.20	*	.	.	-0.60	0.22
	Tyr	171	A	.	.	B	.	.	.	-0.16	*	.	.	-0.60	0.47
	Ala	172	A	.	.	B	.	.	.	-0.50	*	.	.	-0.45	1.35
	Val	173	A	.	.	B	.	.	.	0.28	*	.	.	-0.30	0.92

	Gln	174	.	.	B	.	.	T	.	0.36	*	*	.	0.27	0.80
	Lys	175	.	.	B	.	.	T	.	0.37	.	.	F	0.59	0.65
	Gly	176	.	.	B	.	.	T	.	0.31	.	.	F	0.91	1.18
	His	177	T	C	0.60	.	*	F	1.73	0.91
5	Leu	178	T	C	1.21	.	.	F	1.70	0.61
	Ser	179	T	C	0.36	.	*	F	0.83	0.97
	Ser	180	.	.	B	.	.	T	.	0.31	.	*	F	0.46	0.53
	Ser	181	.	.	B	.	.	T	.	0.44	*	.	F	0.44	1.11
	Tyr	182	.	.	B	.	.	T	.	-0.33	*	.	F	0.57	1.28
10	Val	183	.	.	B	B	.	.	.	-0.33	.	.	.	-0.60	0.79
	Gln	184	.	.	B	B	.	.	.	-0.38	.	.	.	-0.60	0.49
	Pro	185	.	.	B	B	.	.	.	-0.03	*	.	F	-0.45	0.31
	Leu	186	.	.	B	-0.08	*	.	F	0.30	0.83
	Leu	187	.	.	B	0.17	*	.	F	0.55	0.47
15	Gly	188	T	.	.	1.02	*	.	F	1.80	0.53
	Lys	189	T	.	.	0.36	*	*	F	2.20	1.03
	Gly	190	T	T	.	-0.24	*	*	F	2.50	0.67
	Glu	191	T	T	.	-0.02	*	*	F	2.25	0.56
	Asn	192	.	.	B	.	.	T	.	0.58	*	*	F	1.60	0.28
20	Cys	193	.	.	B	.	.	T	.	1.03	.	*	.	0.60	0.44
	Leu	194	.	.	B	0.99	.	*	.	0.75	0.50
	Ala	195	A	T	.	1.38	.	*	.	0.70	0.54
	Pro	196	A	T	.	1.07	.	*	F	1.00	2.00
	Arg	197	A	T	.	0.77	.	*	F	1.30	3.51
25	Gln	198	A	T	.	0.62	.	*	F	1.30	4.65
	Lys	199	A	A	1.48	*	*	F	0.90	2.48
	Thr	200	A	A	2.11	*	*	F	0.90	2.53
	Ser	201	A	A	1.51	*	*	F	0.90	2.92
	Leu	202	.	A	B	0.81	*	*	F	0.90	1.21
30	Lys	203	.	A	B	0.60	*	.	F	0.45	0.84
	Lys	204	.	A	B	-0.03	*	*	F	0.45	0.97
	Leu	205	A	A	-0.53	*	.	.	0.45	1.19
	Ala	206	A	A	-0.53	*	.	.	0.30	0.49
	Pro	207	A	A	0.24	*	.	.	-0.30	0.33
35	Ala	208	A	A	-0.14	*	.	.	-0.60	0.54
	Leu	209	A	T	.	-1.00	*	.	.	-0.20	0.53
	Ser	210	.	.	B	.	.	T	.	-0.86	*	.	.	-0.20	0.28
	His	211	.	.	B	.	.	T	.	-0.61	.	.	.	-0.20	0.15
	Gly	212	.	.	B	.	.	T	.	-0.40	.	.	.	-0.20	0.18
40	Leu	213	.	.	B	-0.02	.	.	.	0.84	0.23
	Cys	214	T	.	.	0.44	.	*	.	0.98	0.27
	Gly	215	T	.	.	0.86	*	.	F	2.07	0.27
	Glu	216	.	.	B	.	.	T	.	0.68	*	.	F	2.21	0.63
	Pro	217	T	T	.	0.21	*	.	F	3.40	1.82
45	Gly	218	T	T	.	0.72	*	*	F	3.06	1.52
	Arg	219	T	C	1.50	*	*	F	2.52	1.17
	Pro	220	C	1.24	*	*	F	1.98	1.49
	Leu	221	.	.	B	0.93	*	*	F	1.14	1.49
	Ser	222	.	.	B	0.33	*	*	F	0.80	1.10
50	Arg	223	.	.	B	0.47	*	*	.	-0.10	0.58
	Met	224	.	.	B	-0.23	*	*	.	0.05	1.10
	Thr	225	.	.	B	0.02	*	*	.	-0.10	0.83
	Leu	226	.	.	B	0.59	*	*	.	0.50	0.84
	Pro	227	A	0.54	*	*	.	-0.25	1.34
55	Ala	228	A	T	.	-0.46	.	*	.	0.10	0.92
	Lys	229	A	T	.	-0.74	.	*	.	-0.20	0.78
	Tyr	230	.	.	B	.	.	T	.	-1.32	.	*	.	-0.20	0.35
	Gly	231	.	.	B	.	.	T	.	-0.54	.	*	.	-0.20	0.24
	Ile	232	.	.	B	B	.	.	.	-0.64	.	*	.	-0.60	0.17

	Ile	233	.	.	B	B	.	.	.	-0.64	.	*	.	-0.60	0.15
	Ile	234	.	.	B	B	.	.	.	-1.03	*	*	.	-0.60	0.16
	His	235	.	.	B	B	.	.	.	-0.68	*	*	.	-0.60	0.22
	Thr	236	.	.	B	.	.	.	T	-0.64	*	*	.	0.10	0.62
5	Ala	237	T	T	-0.42	*	*	.	0.65	1.27
	Gly	238	T	T	0.47	*	.	F	1.25	0.50
	Arg	239	T	T	0.47	*	*	F	1.25	0.56
	Thr	240	.	.	.	B	.	T	.	0.20	*	.	F	0.25	0.39
	Cys	241	.	.	.	B	.	T	.	0.51	*	.	.	1.01	0.52
10	Asn	242	.	.	.	B	.	T	.	1.10	*	.	.	1.32	0.45
	Ile	243	.	.	B	B	.	.	.	0.78	.	*	F	1.38	0.54
	Ser	244	.	.	B	.	.	.	T	0.78	.	*	F	2.09	0.54
	Asp	245	T	T	0.28	*	*	F	3.10	0.65
	Glu	246	A	T	0.13	*	*	.	2.24	0.77
15	Cys	247	A	T	-0.72	*	*	.	1.93	0.47
	Arg	248	A	.	.	B	.	.	.	0.28	*	*	.	0.92	0.21
	Leu	249	A	.	.	B	.	.	.	0.58	*	*	.	0.61	0.24
	Leu	250	A	.	.	B	.	.	.	-0.31	*	*	.	0.30	0.74
	Val	251	A	.	.	B	.	.	.	-0.31	*	*	.	0.30	0.27
20	Arg	252	A	.	.	B	.	.	.	0.06	*	*	.	-0.30	0.56
	Asp	253	.	.	B	B	.	.	.	-0.76	*	*	F	0.45	0.91
	Ile	254	.	.	B	B	.	.	.	-0.19	*	.	F	0.60	1.06
	Gln	255	.	.	B	B	.	.	.	-0.27	*	.	F	-0.15	0.85
	Ser	256	.	.	B	B	.	.	.	0.59	*	.	.	-0.60	0.35
25	Phe	257	.	.	B	B	.	.	.	0.59	*	.	.	-0.60	0.85
	Tyr	258	.	.	B	B	.	.	.	-0.22	*	*	.	0.01	0.96
	Ile	259	.	.	B	B	.	.	.	0.71	.	*	.	0.32	0.59
	Asp	260	.	.	.	B	.	T	.	0.41	.	.	.	1.78	1.36
	Arg	261	T	.	0.04	.	.	F	2.44	1.16
30	Leu	262	T	T	0.74	.	.	F	3.10	0.89
	Lys	263	T	T	0.10	.	.	F	2.79	0.89
	Ser	264	.	.	B	.	.	.	T	0.64	.	.	F	2.08	0.32
	Cys	265	T	T	0.40	.	*	.	1.72	0.38
	Asp	266	.	.	B	.	.	.	T	0.29	.	*	.	1.01	0.30
35	Ile	267	.	.	B	.	.	.	T	0.40	*	.	.	0.10	0.36
	Gly	268	.	.	B	.	.	.	T	-0.46	*	*	.	-0.20	0.58
	Tyr	269	.	.	B	.	.	.	T	-1.01	.	*	.	-0.20	0.29
	Asn	270	.	.	B	B	.	.	.	-0.69	.	*	.	-0.60	0.30
	Phe	271	.	.	B	B	.	.	.	-0.69	.	*	.	-0.60	0.30
40	Leu	272	.	.	B	B	.	.	.	0.20	.	*	.	-0.60	0.33
	Val	273	.	.	B	B	.	.	.	0.20	.	.	.	-0.30	0.35
	Gly	274	.	.	B	.	.	.	T	-0.14	.	.	F	0.25	0.40
	Gln	275	.	.	B	.	.	.	T	-1.03	.	.	F	0.25	0.49
	Asp	276	T	-0.58	.	.	F	0.45	0.46
45	Gly	277	T	0.23	.	.	F	0.45	0.73
	Ala	278	.	.	B	B	.	.	.	0.74	.	.	.	0.30	0.73
	Ile	279	.	.	B	B	.	.	.	0.23	.	.	.	0.30	0.43
	Tyr	280	.	.	B	B	.	.	.	-0.11	.	.	.	-0.60	0.32
	Glu	281	.	.	B	B	.	.	.	-0.40	.	.	.	-0.60	0.32
50	Gly	282	.	.	.	B	.	T	.	-0.06	.	*	.	-0.20	0.48
	Val	283	.	.	B	B	.	.	.	-0.32	.	*	.	-0.60	0.49
	Gly	284	.	.	.	B	.	T	.	0.57	*	*	.	-0.20	0.21
	Trp	285	.	.	B	B	.	.	.	0.47	.	*	.	-0.60	0.37
	Asn	286	.	.	B	B	.	.	.	0.17	.	*	.	-0.60	0.49
55	Val	287	.	.	B	.	.	T	.	0.21	.	*	F	-0.05	0.66
	Gln	288	.	.	B	.	.	T	.	0.76	.	*	F	-0.05	0.84
	Gly	289	T	T	0.89	.	*	F	0.65	0.76
	Ser	290	T	T	0.83	.	*	F	1.14	1.57
	Ser	291	C	0.59	.	*	F	1.53	0.90

	Thr	292	T	C	1.44	.	*	F	1.62	1.42
	Pro	293	T	C	1.44	.	.	F	2.56	1.78
	Gly	294	T	T	.	0.90	.	.	F	3.40	2.21
	Tyr	295	.	.	B	.	.	T	.	0.61	.	.	F	2.36	1.07
5	Asp	296	.	A	B	0.10	.	.	F	1.47	0.70
	Asp	297	.	A	B	B	.	.	.	0.07	*	*	F	0.53	0.59
	Ile	298	.	A	B	B	.	.	.	-0.61	*	*	.	0.04	0.37
	Ala	299	.	A	B	B	.	.	.	-0.58	.	*	.	0.30	0.16
	Leu	300	.	A	B	B	.	.	.	-1.03	.	*	.	-0.60	0.13
10	Gly	301	.	A	B	B	.	.	.	-1.63	.	*	.	-0.60	0.17
	Ile	302	.	.	B	B	.	.	.	-1.98	.	*	.	-0.60	0.16
	Thr	303	.	.	B	B	.	.	.	-1.40	.	*	.	-0.60	0.19
	Phe	304	.	.	B	B	.	.	.	-1.51	.	*	.	-0.60	0.28
	Met	305	.	.	B	B	.	.	.	-1.01	.	*	.	-0.60	0.35
15	Gly	306	.	.	B	B	.	.	.	-1.01	.	*	.	-0.60	0.35
	Thr	307	.	.	B	B	.	.	.	-1.01	.	*	.	-0.60	0.40
	Phe	308	.	.	B	B	.	.	.	-0.91	.	.	F	-0.45	0.28
	Thr	309	.	.	.	B	T	.	.	-0.42	.	.	F	-0.05	0.44
	Gly	310	C	0.18	.	.	F	-0.05	0.48
20	Ile	311	C	-0.07	.	.	F	-0.05	0.88
	Pro	312	T	C	-0.34	.	.	F	0.45	0.62
	Pro	313	T	C	-0.23	.	.	F	0.45	0.63
	Asn	314	T	C	-0.73	.	.	.	0.00	0.91
25	Ala	315	A	T	.	-0.39	.	.	.	-0.20	0.48
	Ala	316	A	A	-0.09	.	.	.	-0.30	0.54
	Ala	317	A	A	-0.47	.	.	.	-0.30	0.34
	Leu	318	A	A	-0.26	.	.	.	-0.30	0.34
	Glu	319	A	A	-0.26	*	.	.	-0.30	0.58
	Ala	320	A	A	-0.48	*	.	.	0.30	0.97
30	Ala	321	A	A	-0.78	*	.	.	0.30	0.97
	Gln	322	A	A	-0.19	*	.	.	0.30	0.39
	Asp	323	A	A	-0.04	*	.	.	-0.30	0.67
	Leu	324	A	A	-0.63	*	.	.	-0.30	0.36
	Ile	325	A	A	-0.64	*	.	.	-0.30	0.21
35	Gln	326	A	A	-0.91	*	.	.	-0.60	0.12
	Cys	327	A	A	B	-0.87	*	.	.	-0.60	0.11
	Ala	328	.	A	B	-1.21	*	.	.	-0.60	0.32
	Met	329	.	A	B	-0.64	.	*	.	-0.30	0.18
	Val	330	.	A	B	-0.57	.	*	.	-0.60	0.53
40	Lys	331	.	A	B	-0.88	.	*	.	-0.60	0.43
	Gly	332	.	A	B	-0.42	.	*	.	-0.60	0.63
	Tyr	333	.	.	B	0.17	.	*	.	-0.25	1.31
	Leu	334	.	.	B	0.52	.	*	F	0.20	1.05
	Thr	335	.	.	B	.	.	T	.	0.57	.	*	F	0.10	1.67
45	Pro	336	.	.	B	.	.	T	.	-0.29	.	*	F	-0.05	0.88
	Asn	337	.	.	B	.	.	T	.	-0.80	.	.	.	-0.20	0.88
	Tyr	338	.	.	B	.	.	T	.	-0.90	.	.	.	-0.20	0.45
	Leu	339	.	.	B	B	.	.	.	-0.12	.	.	.	-0.60	0.29
	Leu	340	.	.	B	B	.	.	.	-0.11	.	.	.	-0.60	0.24
50	Val	341	.	.	B	B	.	.	.	0.10	.	.	.	-0.60	0.21
	Gly	342	.	.	B	B	.	.	.	-0.76	.	.	.	-0.30	0.42
	His	343	.	.	B	.	.	T	.	-1.10	*	.	.	0.10	0.38
	Ser	344	.	.	B	.	.	T	.	-0.18	*	.	.	0.10	0.52
	Asp	345	.	.	B	.	.	T	.	0.32	*	.	F	1.30	1.03
55	Val	346	.	.	B	.	.	T	.	0.37	*	.	.	0.85	1.09
	Ala	347	.	.	B	B	.	.	.	0.41	*	.	.	0.51	0.67
	Arg	348	.	.	B	B	.	.	.	0.23	*	.	F	0.87	0.54
	Thr	349	.	.	B	B	.	.	.	0.19	*	.	F	0.63	1.12
	Leu	350	.	.	B	B	.	.	.	0.19	*	.	F	1.44	1.10

	Ser	351	T	C	0.46	*	.	F	2.10	0.97
	Pro	352	T	C	0.23	*	*	F	1.29	0.68
	Gly	353	T	T	.	-0.12	*	.	F	0.98	0.68
	Gln	354	.	.	B	.	.	T	.	0.19	*	*	F	0.37	0.79
5	Ala	355	.	.	B	0.11	*	.	.	-0.19	0.82
	Leu	356	.	.	B	B	.	.	.	-0.48	*	.	.	-0.60	0.58
	Tyr	357	.	.	B	B	.	.	.	-0.57	*	.	.	-0.60	0.24
	Asn	358	.	.	B	B	.	.	.	-0.53	*	.	.	-0.60	0.31
	Ile	359	.	.	B	B	.	.	.	-0.82	*	.	.	-0.60	0.55
10	Ile	360	.	.	B	B	.	.	.	-0.44	*	.	.	-0.60	0.37
	Ser	361	.	.	B	B	.	.	.	0.33	*	.	.	-0.60	0.35
	Thr	362	.	.	B	B	.	.	.	-0.12	*	*	.	-0.60	0.69
	Trp	363	T	C	-0.08	*	*	.	0.00	0.85
	Pro	364	T	C	0.78	*	.	.	0.15	1.27
15	His	365	T	T	.	1.28	.	*	.	0.35	1.20
	Phe	366	T	T	.	1.19	.	.	.	0.35	1.45
	Lys	367	T	.	.	1.11	.	.	.	0.45	1.20
	His	368	T	.	.	1.01	.	.	.	0.45	1.13

20

25

30

35

Table IV (PGRP-C):

	Res	Pos.	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	XIII	XIV
	Met	1	.	.	.	B	0.97	*	.	.	1.64	1.91
	Ser	2	.	.	.	B	.	.	T	.	0.76	.	.	.	2.07	2.00
5	Arg	3	.	.	.	B	.	.	T	.	0.33	.	.	.	2.30	1.55
	Arg	4	.	.	.	B	.	.	T	.	-0.09	.	.	.	1.77	1.29
	Ser	5	.	.	.	B	.	.	T	.	-0.29	.	.	.	1.39	0.79
	Met	6	.	.	A	B	0.02	.	.	.	0.76	0.41
	Leu	7	.	.	A	B	-0.27	.	.	.	-0.37	0.22
10	Leu	8	.	.	A	B	-1.19	.	.	.	-0.60	0.17
	Ala	9	.	.	A	B	-1.51	.	.	.	-0.60	0.14
	Trp	10	.	.	A	B	-1.51	.	.	.	-0.60	0.26
	Ala	11	.	.	A	B	-1.72	.	.	.	-0.60	0.42
	Leu	12	.	.	.	B	.	.	T	.	-1.72	*	*	.	-0.20	0.34
15	Pro	13	.	.	.	B	.	.	T	.	-0.80	*	*	.	-0.20	0.27
	Ser	14	.	.	.	B	.	.	T	.	-1.02	*	*	.	0.10	0.52
	Leu	15	.	.	.	B	.	.	T	.	-1.08	*	*	.	-0.20	0.52
	Leu	16	.	.	A	B	-1.08	*	*	.	-0.30	0.33
	Arg	17	.	.	A	B	-0.86	*	*	.	-0.30	0.25
20	Leu	18	.	.	A	B	-0.64	.	*	.	-0.60	0.31
	Gly	19	.	.	A	C	-0.34	.	*	.	-0.10	0.65
	Ala	20	.	.	A	C	0.16	.	*	.	0.50	0.57
	Ala	21	.	.	A	C	0.97	.	*	.	0.65	1.00
	Gln	22	.	.	A	B	0.86	.	*	F	1.21	1.75
25	Glu	23	.	.	A	B	1.46	.	.	F	1.52	2.90
	Thr	24	.	.	A	.	.	T	.	.	1.21	.	.	F	2.23	4.44
	Glu	25	.	.	A	.	.	T	.	.	1.13	.	.	F	2.54	2.59
	Asp	26	T	T	.	1.06	.	.	F	3.10	0.80
	Pro	27	T	T	.	0.76	.	.	F	2.49	0.30
30	Ala	28	T	T	.	0.54	.	.	.	2.33	0.23
	Cys	29	T	T	.	-0.03	.	.	.	1.72	0.21
	Cys	30	.	.	.	B	B	.	.	.	-0.89	.	.	.	-0.29	0.10
	Ser	31	.	.	.	B	B	.	.	.	-1.10	.	.	.	-0.60	0.07
	Pro	32	.	.	.	B	B	.	.	.	-0.78	.	.	.	-0.60	0.20
35	Ile	33	.	.	.	B	B	.	.	.	-0.19	.	.	.	0.00	0.75
	Val	34	.	.	.	B	.	.	T	.	0.48	.	*	F	1.45	0.90
	Pro	35	.	.	.	B	.	.	T	.	0.86	.	*	F	1.90	1.01
	Arg	36	T	T	.	1.20	.	*	F	2.00	1.51
	Asn	37	T	C	0.82	*	*	F	3.00	4.07
40	Glu	38	.	.	A	.	.	T	.	.	0.90	*	*	F	2.50	2.66
	Trp	39	.	.	A	.	.	T	.	.	1.17	*	*	F	2.20	1.12
	Lys	40	.	.	A	C	1.08	*	*	.	1.10	0.70
	Ala	41	.	.	A	C	0.97	*	*	.	0.80	0.54
	Leu	42	.	.	A	C	0.30	*	.	.	0.50	0.90
45	Ala	43	A	A	A	-0.29	*	*	.	0.30	0.24
	Ser	44	A	A	A	0.00	*	.	.	-0.30	0.24
	Glu	45	A	A	A	-0.08	*	.	.	-0.30	0.50
	Cys	46	A	A	A	-0.30	*	.	.	0.30	0.68
	Ala	47	A	A	A	0.21	.	.	.	-0.30	0.42
50	Gln	48	.	.	A	B	-0.01	.	.	.	-0.30	0.32
	His	49	.	.	A	B	0.08	*	*	.	-0.60	0.50
	Leu	50	.	.	A	B	-0.73	*	*	.	-0.60	0.76
	Ser	51	.	.	A	B	0.04	*	*	.	-0.60	0.36
	Leu	52	.	.	.	B	0.39	*	*	.	-0.10	0.52
55	Pro	53	.	.	.	B	B	.	.	.	-0.47	*	*	.	-0.60	0.99
	Leu	54	.	.	.	B	B	.	.	.	-1.29	.	*	.	-0.60	0.55
	Arg	55	.	.	.	B	B	.	.	.	-1.33	.	*	.	-0.60	0.49

	Tyr	56	.	.	B	B	.	.	.	-1.33	.	*	.	-0.60	0.24
	Val	57	.	.	B	B	.	.	.	-0.56	.	*	.	-0.60	0.39
	Val	58	.	.	B	B	.	.	.	-0.66	.	*	.	-0.60	0.27
	Val	59	.	.	B	B	.	.	.	-0.43	.	*	.	-0.60	0.25
5	Ser	60	.	.	B	-0.89	.	.	.	-0.40	0.34
	His	61	.	.	B	-0.94	.	.	.	-0.40	0.45
	Thr	62	.	.	B	-0.39	.	.	.	-0.10	0.81
	Ala	63	T	.	.	-0.20	.	.	F	0.45	0.81
	Gly	64	T	T	.	0.66	.	.	F	0.65	0.32
10	Ser	65	T	T	.	0.64	.	.	F	0.65	0.35
	Ser	66	T	T	.	0.47	.	.	F	0.65	0.51
	Cys	67	T	T	.	0.19	.	.	F	0.65	0.79
	Asn	68	T	.	.	0.48	.	.	F	0.45	0.60
	Thr	69	C	0.16	.	.	F	0.25	0.60
15	Pro	70	T	T	.	0.46	.	.	F	0.65	0.60
	Ala	71	T	T	.	0.76	.	*	F	0.65	0.64
	Ser	72	.	.	B	.	.	T	.	1.42	.	*	F	0.25	0.77
	Cys	73	.	.	B	.	.	T	.	0.83	*	*	F	0.25	0.86
	Gln	74	.	A	B	1.26	.	*	F	-0.15	0.86
20	Gln	75	.	A	B	1.47	*	*	F	0.60	1.26
	Gln	76	.	A	B	1.20	*	*	F	0.60	3.79
	Ala	77	.	A	B	1.50	*	.	F	0.60	1.62
	Arg	78	.	A	B	2.13	*	.	F	0.60	1.62
	Asn	79	.	A	B	1.89	*	.	.	0.45	1.28
25	Val	80	.	A	B	1.86	*	.	.	-0.15	1.98
	Gln	81	.	A	B	1.26	*	*	.	-0.15	1.37
	His	82	.	A	B	1.89	*	*	.	-0.60	0.85
	Tyr	83	.	.	B	1.47	*	.	.	-0.25	2.28
	His	84	.	.	B	0.66	.	.	.	0.05	1.90
30	Met	85	.	.	B	B	.	.	.	1.17	.	.	.	-0.45	1.15
	Lys	86	.	.	B	B	.	.	.	0.88	.	.	.	-0.60	0.73
	Thr	87	.	.	.	B	T	.	.	0.24	.	.	.	-0.20	0.56
	Leu	88	.	.	.	B	T	.	.	0.49	.	*	.	-0.20	0.30
	Gly	89	.	.	.	B	T	.	.	-0.33	.	.	.	0.10	0.25
35	Trp	90	.	.	B	B	.	.	.	-0.08	.	.	.	-0.60	0.13
	Cys	91	.	.	B	B	.	.	.	-0.37	.	.	.	-0.60	0.16
	Asp	92	.	.	B	.	.	T	.	-0.06	.	*	.	-0.20	0.25
	Val	93	.	.	B	.	.	T	.	0.06	*	.	.	-0.20	0.38
	Gly	94	.	.	B	.	.	T	.	-0.41	.	*	.	-0.20	0.61
40	Tyr	95	.	.	B	.	.	T	.	-1.01	.	*	.	-0.20	0.30
	Asn	96	.	.	B	B	.	.	.	-0.69	.	*	.	-0.60	0.29
	Phe	97	.	.	B	B	.	.	.	-0.69	.	*	.	-0.60	0.29
	Leu	98	.	.	B	B	.	.	.	0.17	.	*	.	-0.60	0.32
	Ile	99	.	.	B	B	.	.	.	0.17	.	.	.	0.30	0.33
45	Gly	100	.	.	B	.	.	T	.	-0.40	.	.	.	0.10	0.38
	Glu	101	.	.	B	.	.	T	.	-1.26	.	.	F	0.85	0.38
	Asp	102	T	T	.	-0.80	.	.	F	1.25	0.40
	Gly	103	T	C	0.01	.	.	F	1.05	0.63
	Leu	104	.	.	B	0.56	*	*	.	0.50	0.63
50	Val	105	.	.	B	1.01	*	*	.	0.78	0.37
	Tyr	106	.	.	B	0.67	*	*	.	1.06	0.74
	Glu	107	.	.	B	0.38	*	.	F	1.49	0.89
	Gly	108	T	T	.	0.72	.	.	F	1.92	1.25
	Arg	109	T	T	.	0.83	.	*	F	2.80	1.29
55	Gly	110	T	T	.	1.38	.	.	F	2.37	0.64
	Trp	111	T	T	.	1.28	.	*	.	1.04	0.94
	Asn	112	C	0.69	.	.	.	0.36	0.47
	Phe	113	.	.	B	1.00	.	.	.	-0.12	0.48
	Thr	114	C	0.59	.	.	.	-0.20	0.63

	Gly	115	C	0.59	.	*	.	-0.20	0.52
	Ala	116	C	0.84	.	*	.	-0.20	0.60
	His	117	T	C	0.03	.	.	.	0.30	0.56
	Ser	118	T	C	0.44	.	.	.	0.00	0.47
5	Gly	119	T	C	0.76	.	.	.	0.00	0.49
	His	120	T	C	0.89	.	.	.	0.00	0.58
	Leu	121	T	.	.	0.88	.	.	.	0.00	0.67
	Trp	122	C	0.61	.	.	.	-0.20	0.67
	Asn	123	C	0.02	.	.	.	-0.20	0.66
10	Pro	124	.	.	B	B	.	.	.	0.02	.	*	.	-0.60	0.56
	Met	125	.	.	.	B	T	.	.	-0.83	*	*	.	-0.20	0.52
	Ser	126	.	.	B	B	.	.	.	-0.32	.	*	.	-0.60	0.23
	Ile	127	.	.	B	B	.	.	.	-0.73	.	*	.	-0.60	0.20
	Gly	128	.	.	B	B	.	.	.	-1.33	.	*	.	-0.60	0.17
15	Ile	129	.	.	B	B	.	.	.	-1.47	.	*	.	-0.60	0.13
	Ser	130	.	.	B	B	.	.	.	-0.87	.	*	.	-0.60	0.18
	Phe	131	.	.	B	B	.	.	.	-0.81	.	*	.	-0.60	0.29
	Met	132	.	.	B	.	.	T	.	-0.52	.	*	.	-0.20	0.66
	Gly	133	T	T	.	-0.18	*	*	.	0.20	0.48
20	Asn	134	T	T	.	0.82	*	*	.	0.20	0.93
	Tyr	135	T	T	.	0.27	*	*	.	1.25	1.85
	Met	136	T	.	.	0.76	*	*	.	1.31	1.39
	Asp	137	T	.	.	1.04	*	*	.	1.57	1.33
	Arg	138	.	.	B	1.18	*	*	F	1.58	1.23
25	Val	139	.	.	B	.	.	T	.	1.18	*	.	F	2.04	1.92
	Pro	140	.	.	B	.	.	T	.	0.83	.	.	F	2.60	1.99
	Thr	141	T	C	0.54	.	*	F	2.24	1.03
	Pro	142	.	.	B	.	.	T	.	0.66	.	*	F	0.73	0.97
	Gln	143	.	A	B	-0.04	*	*	F	1.12	1.23
30	Ala	144	.	A	B	0.22	*	.	.	0.56	0.86
	Ile	145	.	A	B	0.43	*	.	.	-0.30	0.56
	Arg	146	.	A	B	0.40	*	.	.	0.30	0.56
	Ala	147	.	A	B	-0.20	*	.	.	-0.30	0.55
	Ala	148	.	A	B	-1.01	*	.	.	-0.30	0.65
35	Gln	149	.	A	B	-1.01	*	*	.	-0.30	0.27
	Gly	150	.	A	B	-0.79	*	*	.	-0.60	0.27
	Leu	151	.	A	B	-1.24	*	.	.	-0.60	0.14
	Leu	152	.	A	B	-1.51	.	.	.	-0.60	0.08
	Ala	153	.	A	B	-1.51	.	.	.	-0.60	0.06
40	Cys	154	.	A	B	-1.51	.	.	.	-0.60	0.08
	Gly	155	.	A	B	-1.51	.	.	.	-0.60	0.16
	Val	156	.	A	B	-1.29	.	.	.	-0.60	0.16
	Ala	157	.	A	B	-1.29	*	*	.	-0.60	0.29
	Gln	158	.	A	B	-0.59	*	*	.	-0.60	0.25
45	Gly	159	.	A	B	-0.22	*	*	.	-0.30	0.65
	Ala	160	.	A	B	0.12	*	*	F	0.45	0.86
	Leu	161	.	A	B	0.73	*	*	F	0.45	0.80
	Arg	162	.	.	B	.	.	T	.	0.47	*	*	F	0.40	1.26
	Ser	163	.	.	B	.	.	T	.	-0.34	*	*	F	0.25	0.93
50	Asn	164	.	.	B	.	.	T	.	0.04	.	*	F	-0.05	0.93
	Tyr	165	.	.	B	.	.	T	.	0.29	.	*	.	0.70	0.95
	Val	166	.	.	B	B	.	.	.	1.07	.	*	.	-0.12	0.70
	Leu	167	.	.	B	B	.	.	.	1.07	*	*	.	-0.24	0.59
	Lys	168	.	.	B	B	.	.	.	1.37	*	.	F	0.39	0.74
55	Gly	169	.	.	B	0.51	*	*	F	1.82	1.67
	His	170	.	.	B	B	.	.	.	0.76	*	.	F	1.80	1.50
	Arg	171	.	.	B	B	.	.	.	1.72	*	.	F	1.62	1.30
	Asp	172	.	.	B	B	.	.	.	2.22	*	.	F	1.44	2.57
	Val	173	.	.	B	B	.	.	.	1.37	*	.	F	1.26	2.72

	Gln	174	.	.	B	B	.	.	.	1.41	*	.	F	1.08	1.15
	Arg	175	.	.	B	B	.	.	.	1.23	*	.	F	0.57	0.92
	Thr	176	.	.	B	B	.	.	.	0.78	*	.	F	0.24	1.92
	Leu	177	.	.	.	B	.	.	C	0.78	*	.	F	1.16	1.10
5	Ser	178	T	C	1.63	*	.	F	1.53	0.90
	Pro	179	T	C	0.82	*	.	F	1.20	1.08
	Gly	180	T	T	.	0.47	*	.	F	0.98	1.08
	Asn	181	T	T	.	0.74	.	.	F	0.86	1.26
	Gln	182	.	A	B	0.74	*	.	F	-0.06	1.11
10	Leu	183	.	A	B	0.16	*	.	.	-0.48	0.93
	Tyr	184	.	A	B	0.37	*	.	.	-0.60	0.40
	His	185	.	A	B	0.71	*	.	.	-0.60	0.40
	Leu	186	.	A	B	0.42	*	.	.	-0.60	0.79
	Ile	187	.	A	B	0.21	*	.	.	-0.60	0.53
15	Gln	188	.	A	B	0.99	*	.	.	-0.60	0.60
	Asn	189	.	A	.	.	T	.	.	0.99	.	*	.	-0.20	0.99
	Trp	190	T	C	1.13	.	*	.	0.15	2.21
	Pro	191	T	C	1.64	.	*	.	0.45	2.50
	His	192	T	T	.	2.32	.	*	.	0.86	2.09
20	Tyr	193	T	T	.	1.93	.	.	.	0.77	3.07
	Arg	194	T	.	.	1.54	.	.	.	1.68	2.54
	Ser	195	C	1.44	.	.	.	1.69	2.38
	Pro	196	T	.	.	1.27	.	*	.	2.10	1.94

25

Among highly preferred fragments in this regard are those that comprise regions of PGRP-K (SEQ ID NO:2), PGRP-W (SEQ ID NO:4), AND PGRP-C (SEQ ID NO:6) that combine several structural features, such as several of the features set out above.

The polypeptides of the present invention could be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art.

As described in detail below, the polypeptides of the present invention can also be used to raise polyclonal and monoclonal antibodies, which are useful in assays for detecting PGRP-K, PGRP-W, and/or PGRP-C protein expression as described below or as agonists and antagonists capable of enhancing or inhibiting PGRP-K, PGRP-W, and/or PGRP-C protein function. Further, such polypeptides can be used in the yeast two-hybrid system to "capture" PGRP-K, PGRP-W, and/or PGRP-C protein binding proteins which are also candidate agonists and antagonists according to the present invention. The yeast two hybrid system is described in Fields and Song, Nature 340:245-246 (1989).

Epitope-Bearing Portions

In another aspect, the invention provides a peptide or polypeptide comprising an epitope-bearing portion of a polypeptide of the invention. The epitope of this polypeptide portion is an immunogenic or antigenic epitope of a polypeptide of the invention. The term
5 "epitopes," as used herein, refers to portions of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. In a preferred embodiment, the present invention encompasses a polypeptide comprising an epitope, as well as the polynucleotide encoding this polypeptide. An "immunogenic epitope," as used herein, is defined as a portion of a protein that elicits an antibody response
10 in an animal, as determined by any method known in the art, for example, by the methods for generating antibodies described infra. (See, for example, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998- 4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as determined by any method well known in the art, for example, by the immunoassays
15 described herein. Immunospecific binding excludes non-specific binding but does not necessarily exclude cross- reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic.

Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985), further described in
20 U.S. Patent No. 4,631,211).

In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, and, most preferably, between about 15 to about 30 amino acids. Preferred
25 polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Additional non-exclusive preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as portions thereof. Antigenic epitopes are useful, for example, to raise

antibodies, including monoclonal antibodies, that specifically bind the epitope. Preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these antigenic epitopes. Antigenic epitopes can be used as the target molecules in immunoassays. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983)).

Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985). Preferred immunogenic epitopes include the immunogenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these immunogenic epitopes. The polypeptides comprising one or more immunogenic epitopes may be presented for eliciting an antibody response together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse), or, if the polypeptide is of sufficient length (at least about 25 amino acids), the polypeptide may be presented without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting).

Antigenic epitope-bearing peptides and polypeptides of the invention preferably contain a sequence of at least seven, more preferably at least nine and most preferably between about 15 to about 30 amino acids contained within the amino acid sequences of the polypeptides of the invention. Non-limiting examples of antigenic polypeptides or peptides that can be used to generate PGRP-K specific antibodies include: a polypeptide comprising amino acid residues from about Val-24 to about Ala-35 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Gln-51 to about Gln-58 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Gly-69 to about Ser-72 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Leu-88 to about Gly-100 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about His-107 to about Tyr-111 in

Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Gly-122 to about Pro-131 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Gln-146 to about Ile-155 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Leu-159 to about His-170 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Val-172 to about Pro-200 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Gly-211 to about Val-223 in Figures 1A-B (SEQ ID NO:2); a polypeptide comprising amino acid residues from about Phe-230 to about Tyr-242 in Figures 1A-B (SEQ ID NO:2).

Non-limiting examples of antigenic polypeptides or peptides that can be used to generate PGRP-W specific antibodies include: a polypeptide comprising amino acid residues from about Gly-17 to about Gly-32 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Ile-40 to about Gly-58 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Gly-82 to about Gln-99 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about His-104 to about Val-111 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Leu-116 to about Glu-125 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Gly-150 to about Pro-159 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Gln-174 to about Tyr-182 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Leu-186 to about Pro-207 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Val-214 to about Met-225 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Thr-237 to about Val-252 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Tyr-259 to about Ile-268 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Gly-290 to about Ala-300 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about His-

344 to about Gln-355 in Figures 2A-C (SEQ ID NO:4); a polypeptide comprising amino acid residues from about Trp-364 to about His-369 in Figures 2A-C (SEQ ID NO:4).

Non-limiting examples of antigenic polypeptides or peptides that can be used to generate PGRP-C specific antibodies include: a polypeptide comprising amino acid residues from about Met-1 to about Met-6 in Figure 3 (SEQ ID NO:6); a polypeptide comprising amino acid residues from about Ala-20 to about Cys-29 in Figure 3 (SEQ ID NO:6); a polypeptide comprising amino acid residues from about Ile-33 to about Ala-43 in Figure 3 (SEQ ID NO:6); a polypeptide comprising amino acid residues from about Ala-63 to about Asn-79 in Figure 3 (SEQ ID NO:6); a polypeptide comprising amino acid residues from about Ile-99 to about Asn-112 in Figure 3 (SEQ ID NO:6); a polypeptide comprising amino acid residues from about Gly-133 to about Arg-146 in Figure 3 (SEQ ID NO:6); a polypeptide comprising amino acid residues from about Ala-160 to about Tyr-165 in Figure 3 (SEQ ID NO:6); a polypeptide comprising amino acid residues from about Lys-168 to about Asn-181 in Figure 3 (SEQ ID NO:6); and a polypeptide comprising amino acid residues from about Trp-190 to about Pro-196 in Figure 3 (SEQ ID NO:6). These polypeptide fragments have been determined to bear antigenic epitopes of the PGRP-K, PGRP-W, and PGRP-C proteins, respectively, by the analysis of the Jameson-Wolf antigenic index, as shown in Figures 6, 8, and 10, above.

Epitope-bearing polypeptides of the present invention may be used to induce antibodies according to methods well known in the art including, but not limited to, in vivo immunization, in vitro immunization, and phage display methods. See, e.g., Sutcliffe et al., supra; Wilson et al., supra, and Bittle et al., J. Gen. Virol., 66:2347-2354 (1985). If in vivo immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as maleimidobenzoyl-N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are

immunized with either free or carrier- coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg of peptide or carrier protein and Freund's adjuvant or any other adjuvant known for stimulating an immune response. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention comprising an immunogenic or antigenic epitope can be fused to other polypeptide sequences. For example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof) resulting in chimeric polypeptides. Such fusion proteins may facilitate purification and may increase half-life in vivo. This has been shown for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion disulfide bonds have also been found to be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*, 270:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin ("HA") tag or flag tag) to aid in detection and purification of the expressed polypeptide. For example, a system described by Janknecht et al. allows for the ready

purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972- 897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. The tag
5 serves as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to
10 as "DNA shuffling"). DNA shuffling may be employed to modulate the activities of polypeptides of the invention, such methods can be used to generate polypeptides with altered activity, as well as agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten et al., Curr. Opinion Biotechnol. 8:724-33 (1997); Harayama, Trends Biotechnol. 16(2):76-82
15 (1998); Hansson, et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo and Blasco, Biotechniques 24(2):308- 13 (1998) (each of these patents and publications are hereby incorporated by reference in its entirety). In one embodiment, alteration of polynucleotides corresponding to SEQ ID NO:X and the polypeptides encoded by these polynucleotides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA
20 segments by homologous or site-specific recombination to generate variation in the polynucleotide sequence. In another embodiment, polynucleotides of the invention, or the encoded polypeptides, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of a
25 polynucleotide encoding a polypeptide of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

Fusion Proteins

As one of skill in the art will appreciate, PGRP-K, PGRP-W, and/or PGRP-C polypeptides of the present invention and the epitope-bearing fragments thereof described above can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. This has been shown, e.g., for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins (EP A 394,827; Traunecker et al., Nature 331:84-86 (1988)). Fusion proteins that have a disulfide-linked dimeric structure due to the IgG part can also be more efficient in binding and neutralizing other molecules than the monomeric PGRP-K, PGRP-W, and/or PGRP-C proteins or protein fragments alone (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995)).

Detection of Disease States

Cells which express either the PGRP-K, PGRP-W and/or PGRP-C polypeptides are believed to have a potent cellular response to infection include, for example, keratinocyte cells, wound-healing tissues, vascular tissues, endothelial tissues, and tissues of the immune, nervous, and endocrine systems. Furthermore, it is believed that cells which express either the PGRP-K, PGRP-W and/or PGRP-C polypeptides regulate apoptosis and/or the proliferation of keratinocytes, epidermal cells, and epithelial cells, as well as mediating the recognition of pathogens (e.g., bacteria) and the processing and presentation of antigens to the immune system. By "a potent cellular response to infection" is intended any genotypic, phenotypic, and/or morphologic change to a cell, cell line, tissue, tissue culture or patient that is induced by infection from bacterial (Gram positive and negative), viral, fungal, parasitic, etc.. As indicated, such cellular responses include not only normal physiological responses infection (e.g., antigenic processing and presentation, immune response), but also diseases associated with aberrant immune system recognition, aberrant antigen processing and presentation in the immune system, aberrant immune system responses to infection, activation, survival, migration and differentiation of immune cells, as well as infections of immuno-compromised individuals, and aberrant regulation of the proliferation/apoptosis of keratinocytes and/or other cells in the body (e.g., immune system cells).

Thus, it is believed that certain tissues in mammals with certain diseases and infections (e.g., bacterial infection of immuno-compromised individuals), diseases associated with increased or decreased cell survival, secretion, activation, migration, differentiation, and proliferation; diseases associated with the defects of wound healing, keratinocyte and cartilage proliferation, cellular immunity, immune dysfunction, and endocrine dysfunction; express significantly altered (e.g., enhanced or decreased) levels of either the PGRP-K, PGRP-W and/or PGRP-C polypeptides and mRNAs encoding the PGRP-K, PGRP-W and/or PGRP-C polypeptides when compared to a corresponding "standard" mammal, i.e., a mammal of the same species not having the disease. Diseases associated with defects in the proliferation of keratinocytes or cartilagenous tissues, include, for example, skin or cartilagenous cancers (such as Chondrosarcomas, basal cell carcinomas, squamous cell carcinomas, melanomas, Chondromatosis, Dyschondroplasia). Diseases associated with immune dysfunction and decreased cellular immunity include, for example, bacterial infections (e.g., cutaneous infection due to *Mycobacterium gordonae* in an AIDS patient) and diseases associated with bacterial infection of the skin (e.g., boils, cellulitis, erysipelas, impetigo).

Further, it is believed that altered levels of either the PGRP-K, PGRP-W and/or PGRP-C polypeptide can be detected in certain body fluids (e.g., lymph, sera, plasma, urine, and

spinal fluid) from mammals with the disorder when compared to sera from mammals of the same species not having the disorder. Thus, the invention provides a diagnostic method useful during diagnosis, which involves assaying the expression level of the gene encoding the PGRP-K, PGRP-W and/or PGRP-C polypeptide(s) in mammalian cells or body fluid and
5 comparing the gene expression level with a standard PGRP-K, PGRP-W and/or PGRP-C gene expression level, whereby an increase or decrease in the gene expression level over the standard is indicative of the disease.

By "assaying" the expression level of the gene encoding either the "PGRP-K, PGRP-W and/or PGRP-C polypeptide" is intended qualitatively or quantitatively measuring or
10 estimating the level of the PGRP-K, PGRP-W and/or PGRP-C polypeptide(s) or the level of the mRNA encoding either the PGRP-K, PGRP-W and/or PGRP-C polypeptide(s) in a first biological sample either directly (e.g., by determining or estimating absolute polypeptide or mRNA level) or relatively (e.g., by comparing to either the PGRP-K, PGRP-W and/or PGRP-C polypeptide(s) level or mRNA level in a second biological sample). Preferably, the PGRP-K, PGRP-W and/or PGRP-C protein level or mRNA level in the first biological sample is
15 measured or estimated and compared to a standard PGRP-K, PGRP-W and/or PGRP-C protein level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the disease state. As will be appreciated in the art, once a standard PGRP-K, PGRP-W and/or PGRP-C protein level or mRNA level is known, it can
20 be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source which contains PGRP-K, PGRP-W and/or PGRP-C protein or mRNA. Biological samples include mammalian body fluids (such as lymph, sera, plasma, urine, synovial fluid and spinal fluid), and keratinocytes, wound-healing tissues,
25 human chondrosarcoma, and other tissues. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source. Where a diagnosis has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting altered PGRP-K, PGRP-W and/or PGRP-C gene expression will
30 experience a worse clinical outcome relative to patients expressing the gene at a normal level.

Nucleic acids for diagnosis may be obtained from a biological sample of a subject, such as from blood, urine, saliva, tissue biopsy or autopsy material, using techniques known in the art. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be
35 used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by

hybridizing amplified DNA to labeled PGRP-K, PGRP-W, and/or PGRP-C nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, e.g., Myers et al., Science 230:1242 (1985)). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton et al., Proc. Natl. Acad. Sci. USA 85:4397-4401 (1985)). In another embodiment, an array of oligonucleotide probes comprising either PGRP-K, PGRP-W, and/or PGRP-C polynucleotide sequences or fragments thereof, can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example, Chee et al., Science 274:610-613 (1996)). The diagnostic assays offer a process for diagnosing or determining a susceptibility to specific diseases through detection of mutations in the PGRP-K, PGRP-W and/or PGRP-C genes by the methods described herein or otherwise known in the art.

In addition, specific diseases can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of PGRP-K, PGRP-W and/or PGRP-C polypeptides or mRNAs. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art, which include, but are not limited to, Northern blot analysis, (Harada et al., Cell 63:303-312 (1990)), S1 nuclease mapping (Fijita et al., Cell 49:357-367 (1987)), RNase protection, the polymerase chain reaction (PCR), reverse transcription in combination with the polymerase chain reaction (RT-PCR) (Makino et al., Technique 2:295-301 (1990)), reverse transcription in combination with the ligase chain reaction (RT-LCR) and other hybridization methods.

Assaying PGRP-K, PGRP-W, and/or PGRP-C polypeptide levels in a biological sample can be by any techniques known in the art, which include, but are not limited to, radioimmunoassays, competitive-binding assays, Western Blot analysis and enzyme linked immunosorbent assays (ELISAs) and other antibody-based techniques. For example, PGRP-K, PGRP-W, and/or PGRP-C polypeptide expression in tissues can be studied with classical immunohistological methods (Jalkanen et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen et al., J. Cell. Biol. 105:3087-3096 (1987)). Suitable labels are known in the art and include enzyme labels, such as, Glucose oxidase, and radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium ($^{99\text{m}}\text{Tc}$), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

Antibodies

Further polypeptides of the invention relate to antibodies and T-cell antigen receptors
5 (TCR) which immunospecifically bind a polypeptide, polypeptide fragment, or variant of
SEQ ID NOs: 2, 4, and/or 6, and/or an epitope, of the present invention (as determined by
immunoassays well known in the art for assaying specific antibody-antigen binding).
Antibodies of the invention include, but are not limited to, polyclonal, monoclonal,
multispecific, human, humanized or chimeric antibodies, single chain antibodies, Fab
10 fragments, F(ab') fragments, fragments produced by a Fab expression library, anti-idiotypic
(anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and
epitope-binding fragments of any of the above. The term "antibody," as used herein, refers to
immunoglobulin molecules and immunologically active portions of immunoglobulin
molecules, i.e., molecules that contain an antigen binding site that immunospecifically binds an
15 antigen. The immunoglobulin molecules of the invention can be of any type (e.g., IgG, IgE,
IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of
immunoglobulin molecule.

Most preferably the antibodies are human antigen-binding antibody fragments of the
present invention and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain
20 Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising
either a VL or VH domain. Antigen-binding antibody fragments, including single-chain
antibodies, may comprise the variable region(s) alone or in combination with the entirety or a
portion of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the
invention are antigen-binding fragments also comprising any combination of variable region(s)
25 with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be
from any animal origin including birds and mammals. Preferably, the antibodies are human,
murine (e.g., mouse and rat), donkey, sheep rabbit, goat, guinea pig, camel, horse, or chicken.
As used herein, "human" antibodies include antibodies having the amino acid sequence of a

human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described infra and, for example in, U.S. Patent No. 5,939,598 by Kucherlapati et al.

5 The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO
10 91/00360; WO 92/05793; Tutt, et al., J. Immunol. 147:60-69 (1991); U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny et al., J. Immunol. 148:1547-1553 (1992).

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which they recognize or
15 specifically bind. The epitope(s) or polypeptide portion(s) may be specified as described herein, e.g., by N-terminal and C-terminal positions, by size in contiguous amino acid residues, or listed in the Tables and Figures. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention,
20 and allows for the exclusion of the same.

Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of a polypeptide of the present invention are included. Antibodies that bind polypeptides with at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%,
25 at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In specific embodiments, antibodies of the present invention cross-react with murine, rat and/or rabbit homologs of human proteins and the corresponding epitopes

thereof. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of 2, 3, 4, 5, or more of the specific antigenic and/or immunogenic polypeptides disclosed herein. Further included in the present invention are antibodies which bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide of the invention. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, or 10^{-15} M.

The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for determining competitive binding, for example, the immunoassays described herein. In preferred embodiments, the antibody competitively inhibits binding to the epitope by at least 95%, at least 90%, at least 85 %, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50%.

Antibodies of the present invention may act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes antibodies which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Preferably, antibodies of the present invention bind an antigenic epitope disclosed herein, or a portion thereof. The invention features both receptor-specific antibodies and ligand-specific antibodies. The invention also features receptor-

specific antibodies which do not prevent ligand binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. For example, receptor activation can be determined by detecting the phosphorylation (e.g., tyrosine or serine/threonine) of the receptor or its substrate by immunoprecipitation followed by western blot analysis (for example, as described supra). In specific embodiments, antibodies are provided that inhibit ligand activity or receptor activity by at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50% of the activity in absence of the antibody.

The invention also features receptor-specific antibodies which both prevent ligand binding and receptor activation as well as antibodies that recognize the receptor-ligand complex, and, preferably, do not specifically recognize the unbound receptor or the unbound ligand. Likewise, included in the invention are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but do not prevent the ligand from binding the receptor. Further included in the invention are antibodies which activate the receptor. These antibodies may act as receptor agonists, i.e., potentiate or activate either all or a subset of the biological activities of the ligand-mediated receptor activation, for example, by inducing dimerization of the receptor. The antibodies may be specified as agonists, antagonists or inverse agonists for biological activities comprising the specific biological activities of the peptides of the invention disclosed herein. The above antibody agonists can be made using methods known in the art. See, e.g., PCT publication WO 96/40281; U.S. Patent No. 5,811,097; Deng et al., *Blood* 92(6):1981-1988 (1998); Chen et al., *Cancer Res.* 58(16):3668-3678 (1998); Harrop et al., *J. Immunol.* 161(4):1786-1794 (1998); Zhu et al., *Cancer Res.* 58(15):3209-3214 (1998); Yoon et al., *J. Immunol.* 160(7):3170-3179 (1998); Prat et al., *J. Cell. Sci.* 111(Pt2):237-247 (1998); Pitard et al., *J. Immunol. Methods* 205(2):177-190 (1997); Liautard et al., *Cytokine* 9(4):233-241 (1997); Carlson et al., *J. Biol. Chem.* 272(17):11295-11301 (1997); Taryman et al., *Neuron* 14(4):755-762 (1995); Muller et al.,

Structure 6(9):1153-1167 (1998); Bartunek et al., Cytokine 8(1):14-20 (1996) (which are all incorporated by reference herein in their entireties).

Antibodies of the present invention may be used, for example, but not limited to, to purify, detect, and target the polypeptides of the present invention, including both in vitro and in vivo diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g., Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference herein in its entirety).

As discussed in more detail below, the antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO 92/08495; WO 91/14438; WO 89/12624; U.S. Patent No. 5,314,995; and EP 396,387.

The antibodies of the invention include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

The antibodies of the present invention may be generated by any suitable method known in the art. Polyclonal antibodies to an antigen-of-interest can be produced by various procedures well known in the art. For example, a polypeptide of the invention can be administered to various host animals including, but not limited to, rabbits, mice, rats, etc. to induce the production of sera containing polyclonal antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants are also well known in the art.

Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art and are discussed in detail in the Examples (e.g., Example 16). In a non-limiting example, mice can be immunized with a polypeptide of the invention or a cell expressing such peptide. Once an immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known

techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are selected and cloned by limited dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the invention. Ascites fluid, which generally contains high levels of antibodies, can be generated by immunizing mice with positive hybridoma clones.

Accordingly, the present invention provides methods of generating monoclonal antibodies as well as antibodies produced by the method comprising culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by fusing splenocytes isolated from a mouse immunized with an antigen of the invention with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody able to bind a polypeptide of the invention.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')₂ fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain.

For example, the antibodies of the present invention can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such phage can be utilized to display antigen binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., J.

Immunol. Methods 182:41-50 (1995); Ames et al., J. Immunol. Methods 184:177-186 (1995); Kettleborough et al., Eur. J. Immunol. 24:952-958 (1994); Persic et al., Gene 187 9-18 (1997); Burton et al., Advances in Immunology 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., BioTechniques 12(6):864-869 (1992); and Sawai et al., AJRI 34:26-34 (1995); and Better et al., Science 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., Methods in Enzymology 203:46-88 (1991); Shu et al., PNAS 90:7995-7999 (1993); and Skerra et al., Science 240:1038-1040 (1988). For some uses, including in vivo use of antibodies in humans and in vitro detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies et al., (1989) J. Immunol. Methods 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated

herein by reference in their entirety. Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and a framework regions from a human immunoglobulin molecule. Often, framework residues in the human
5 framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See,
10 e.g., Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., Nature 332:323 (1988), which are incorporated herein by reference in their entirety.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, Molecular Immunology
15 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994); Roguska et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Patent No. 5,565,332).

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human
20 immunoglobulin sequences. See also, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

Human antibodies can also be produced using transgenic mice which are incapable of
25 expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and

diversity region may be introduced into mouse embryonic stem cells in addition to the human heavy and light chain genes. The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce chimeric mice. The chimeric mice are then bred to produce homozygous offspring which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, *Int. Rev. Immunol.* 13:65-93 (1995). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al., *Bio/technology* 12:899-903 (1988)).

Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, FASEB J. 7(5):437-444; (1989) and Nissinoff, J. Immunol. 147(8):2429-2438 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand. For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligands/receptors, and thereby block its biological activity.

Polynucleotides Encoding Antibodies

The invention further provides polynucleotides comprising a nucleotide sequence encoding an antibody of the invention and fragments thereof. The invention also encompasses polynucleotides that hybridize under stringent or lower stringency hybridization conditions, e.g., as defined supra, to polynucleotides that encode an antibody, preferably, that specifically binds to a polypeptide of the invention, preferably, an antibody that binds to a polypeptide having the amino acid sequence of SEQ ID NOs: 2, 4, and/or 6.

The polynucleotides may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier et al., BioTechniques 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.

Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g.,
5 an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone
10 from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

Once the nucleotide sequence and corresponding amino acid sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA
15 techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel et al., eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example
20 to create amino acid substitutions, deletions, and/or insertions.

In a specific embodiment, the amino acid sequence of the heavy and/or light chain variable domains may be inspected to identify the sequences of the complementarity determining regions (CDRs) by methods that are well known in the art, e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the
25 regions of sequence hypervariability. Using routine recombinant DNA techniques, one or more of the CDRs may be inserted within framework regions, e.g., into human framework regions to humanize a non-human antibody, as described supra. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework

regions (see, e.g., Chothia et al., J. Mol. Biol. 278: 457-479 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a polypeptide of the invention. Preferably, as discussed supra, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., Proc. Natl. Acad. Sci. 81:851-855 (1984); Neuberger et al., Nature 312:604-608 (1984); Takeda et al., Nature 314:452-454 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described supra, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, Science 242:423-42 (1988); Huston et al., Proc. Natl. Acad. Sci. USA 85:5879-5883 (1988); and Ward et al., Nature 334:544-54 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in E. coli may also be used (Skerra et al., Science 242:1038-1041 (1988)).

Methods of Producing Antibodies

The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques.

Recombinant expression of an antibody of the invention, or fragment, derivative or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably containing the heavy or light chain variable domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, or a heavy or light chain thereof, or a heavy or light chain variable domain, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention, or a heavy or light chain thereof, or a single chain antibody of the invention, operably linked to a heterologous promoter. In preferred embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains

may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent
5 cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors
10 containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic-virus, CaMV; tobacco mosaic virus, TMV) or transformed
15 with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such
20 as *Escherichia coli*, and more preferably, eukaryotic cells, especially for the expression of whole recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., *Gene*
25 45:101 (1986); Cockett et al., *Bio/Technology* 8:2 (1990)).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical

compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., EMBO J. 2:1791 (1983)), in which the antibody coding sequence may be ligated individually into the vector in
5 frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, Nucleic Acids Res. 13:3101-3109 (1985); Van Heeke & Schuster, J. Biol. Chem. 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to
10 matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells.
15 The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence
20 of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts. (e.g., see Logan & Shenk,
25 Proc. Natl. Acad. Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the

entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., *Methods in Enzymol.* 153:51-544 (1987)).

5 In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification
10 of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS,
15 MDCK, 293, 3T3, WI38, and in particular, breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell line such as, for example, CRL7030 and Hs578Bst.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the antibody molecule may be
20 engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are
25 switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody

molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., Cell 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy et al., Cell 22:817 (1980)) genes can be employed in tk-, hgp^rt- or ap^rt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., Natl. Acad. Sci. USA 77:357 (1980); O'Hare et al., Proc. Natl. Acad. Sci. USA 78:1527 (1981)); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 Clinical Pharmacy 12:488-505; Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, 1993, TIB TECH 11(5):155-215; and hyg^r, which confers resistance to hygromycin (Santerre et al., Gene 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli et al. (eds), Current Protocols in Human Genetics, John Wiley & Sons, NY (1994); Colberre-Garapin et al., J. Mol. Biol. 150:1 (1981), which are incorporated by reference herein in their entireties.

The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the

number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., Mol. Cell. Biol. 3:257 (1983)).

The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, Nature 322:52 (1986); Kohler, Proc. Natl. Acad. Sci. USA 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once an antibody molecule of the invention has been produced by an animal, chemically synthesized, or recombinantly expressed, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art, to facilitate purification.

The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences. The antibodies may be specific for antigens other than polypeptides (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to

particular cell types, either in vitro or in vivo, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in in vitro immunoassays and purification methods using methods known in the art. See e.g., Harbor et al., supra, and PCT publication WO 93/21232; EP 439,095; Naramura et al., Immunol. Lett. 39:91-99 (1994); U.S. Patent 5,474,981; Gillies et al., PNAS 89:1428-1432 (1992); Fell et al., J. Immunol. 146:2446-2452(1991), which are incorporated by reference in their entireties.

The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the constant region, hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. See, e.g., U.S. Patent Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,112,946; EP 307,434; EP 367,166; PCT publications WO 96/04388; WO 91/06570; Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 (1991); Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., Proc. Natl. Acad. Sci. USA 89:11337-11341(1992) (said references incorporated by reference in their entireties).

As discussed, supra, the polypeptides corresponding to a polypeptide, polypeptide fragment, or a variant of SEQ ID NOs: 2, 4, and/or 6 may be fused or conjugated to the above antibody portions to increase the in vivo half life of the polypeptides or for use in immunoassays using methods known in the art. Further, the polypeptides corresponding to SEQ ID NOs: 2, 4, and/or 6 may be fused or conjugated to the above antibody portions to

facilitate purification. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP 394,827; Traunecker et al., *Nature* 331:84-86 (1988)). The polypeptides of the present invention
5 fused or conjugated to an antibody having disulfide-linked dimeric structures (due to the IgG) may also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., *J. Biochem.* 270:3958-3964 (1995)). In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP A 232,262).
10 Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995)).

Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE
20 vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein
25 (Wilson et al., *Cell* 37:767 (1984)) and the "flag" tag.

The present invention further encompasses antibodies or fragments thereof conjugated to a diagnostic or therapeutic agent. The antibodies can be used diagnostically to, for example, monitor the development or progression of a tumor as part of a clinical testing

procedure to, e.g., determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the antibody (or fragment thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{111}In or ^{99}Tc .

Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ^{213}Bi . A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU),

cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

The conjugates of the invention can be used for modifying a given biological response, the therapeutic agent or drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNF- α , TNF- β , AIM I (See, International Publication No. WO 97/33899), AIM II (See, International Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *Int. Immunol.*, 6:1567-1574 (1994)), VEGF (See, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.* (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987);

Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies '84: Biological And Clinical Applications, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in Monoclonal Antibodies For Cancer Detection
5 And Therapy, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", Immunol. Rev. 62:119-58 (1982).

Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated
10 herein by reference in its entirety.

An antibody, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can be used as a therapeutic.

Immunophenotyping

15 The antibodies of the invention may be utilized for immunophenotyping of cell lines and biological samples. The translation product of the gene of the present invention may be useful as a cell specific marker, or more specifically as a cellular marker that is differentially expressed at various stages of differentiation and/or maturation of particular cell types. Monoclonal antibodies directed against a specific epitope, or combination of epitopes, will
20 allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Patent 5,985,660; and Morrison *et al.*, *Cell*, 96:737-49 (1999)).

25 These techniques allow for the screening of particular populations of cells, such as might be found with hematological malignancies (i.e. minimal residual disease (MRD) in acute leukemic patients) and "non-self" cells in transplantations to prevent Graft-versus-Host Disease (GVHD). Alternatively, these techniques allow for the screening of hematopoietic

stem and progenitor cells capable of undergoing proliferation and/or differentiation, as might be found in human umbilical cord blood.

Assays For Antibody Binding

5 The antibodies of the invention may be assayed for immunospecific binding by any method known in the art. The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin
10 reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described
15 briefly below (but are not intended by way of limitation).

Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasylol) supplemented with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium
20 vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (e.g., 1-4 hours) at 4°C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 4°C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One
25 of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding

immunoprecipitation protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (e.g., PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), blocking the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) or radioactive molecule (e.g., ³²P or ¹²⁵I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the

art. For further discussion regarding ELISAs see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 11.2.1.

The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g., ³H or ¹²⁵I) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the antibody bound to the labeled antigen. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of interest conjugated to a labeled compound (e.g., ³H or ¹²⁵I) in the presence of increasing amounts of an unlabeled second antibody.

Therapeutic Uses

The present invention is further directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more of the disclosed diseases, disorders, or conditions. Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the invention, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders

or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

The antibodies of the invention may be administered alone or in combination with other types of treatments (e.g., radiation therapy, chemotherapy, hormonal therapy, immunotherapy and anti-tumor agents). Generally, administration of products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment, human antibodies, fragments derivatives, analogs, or nucleic acids, are administered to a human patient for therapy or prophylaxis.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides of the invention, including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8}

M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, and 10^{-15} M.

Gene Therapy

5 In a specific embodiment, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or prevent a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the
10 nucleic acids produce their encoded protein that mediates a therapeutic effect.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

For general reviews of the methods of gene therapy, see Goldspiel et al., Clinical Pharmacy 12:488-505 (1993); Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann.
15 Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, TIBTECH 11(5):155-215 (1993). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); and Kriegler, Gene Transfer and Expression, A Laboratory
20 Manual, Stockton Press, NY (1990).

In a preferred aspect, the compound comprises nucleic acid sequences encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody
25 coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for

intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989). In specific embodiments, the expressed antibody molecule is a single chain antibody; alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids in vitro, then transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

In a specific embodiment, the nucleic acid sequences are directly administered in vivo, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated retrovirals or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by

homologous recombination (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989)).

In a specific embodiment, viral vectors that contains nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller et al., *Meth. Enzymol.* 217:581-599 (1993)). These retroviral vectors contain the components
5 necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., *Biotherapy* 6:291-302 (1994),
10 which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., *J. Clin. Invest.* 93:644-651 (1994); Kiem et al., *Blood* 83:1467-1473 (1994); Salmons and Gunzberg, *Human Gene Therapy* 4:129-141 (1993); and Grossman and Wilson, *Curr. Opin. in Genetics and*
15 *Devel.* 3:110-114 (1993).

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and
20 muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current Opinion in Genetics and Development* 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., *Human Gene Therapy* 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can
25 be found in Rosenfeld et al., *Science* 252:431-434 (1991); Rosenfeld et al., *Cell* 68:143- 155 (1992); Mastrangeli et al., *J. Clin. Invest.* 91:225-234 (1993); PCT Publication WO94/12649; and Wang, et al., *Gene Therapy* 2:775-783 (1995). In a preferred embodiment, adenovirus vectors are used.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., Proc. Soc. Exp. Biol. Med. 204:289-300 (1993); U.S. Patent No. 5,436,146).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler and Behr, Meth. Enzymol. 217:599-618 (1993); Cohen et al., Meth. Enzymol. 217:618-644 (1993); Cline, Pharmac. Ther. 29:69-92m (1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as Tlymphocytes, Blymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic

stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

In an embodiment in which recombinant cells are used in gene therapy, nucleic acid
5 sequences encoding an antibody are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained in vitro can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO
10 94/08598; Stemple and Anderson, Cell 71:973-985 (1992); Rheinwald, Meth. Cell Bio. 21A:229 (1980); and Pittelkow and Scott, Mayo Clinic Proc. 61:771 (1986)).

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the
15 appropriate inducer of transcription. Demonstration of Therapeutic or Prophylactic Activity

The compounds or pharmaceutical compositions of the invention are preferably tested in vitro, and then in vivo for the desired therapeutic or prophylactic activity, prior to use in humans. For example, in vitro assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition include, the effect of a compound on a cell line
20 or a patient tissue sample. The effect of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance with the invention, in vitro assays which can be used to determine whether administration of a specific compound is indicated, include in vitro cell culture assays in which a patient tissue
25 sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

Therapeutic/Prophylactic Administration and Composition

The invention provides methods of treatment, inhibition and prophylaxis by administration to a subject of an effective amount of a compound or pharmaceutical composition of the invention, preferably an antibody of the invention. In a preferred aspect, the compound is substantially purified (e.g., substantially free from substances that limit its effect or produce undesired side-effects). The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.

Formulations and methods of administration that can be employed when the compound comprises a nucleic acid or an immunoglobulin are described above; additional appropriate formulations and routes of administration can be selected from among those described herein below.

Various delivery systems are known and can be used to administer a compound of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compounds or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by
5 means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

In another embodiment, the compound or composition can be delivered in a vesicle, in
10 particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

In yet another embodiment, the compound or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*;
15 Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability; Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J.,
20 *Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy et al., *Science* 228:190 (1985); During et al., *Ann. Neurol.* 25:351 (1989); Howard et al., *J. Neurosurg.* 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-
25 138 (1984)).

Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

In a specific embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered in vivo to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see 5 U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox- like peptide which is known to enter the nucleus (see e.g., Joliot et al., Proc. Natl. Acad. Sci. USA 88:1864-1868 (1991)), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated 10 within host cell DNA for expression, by homologous recombination.

The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a compound, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. 15 Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a 20 preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, 25 ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with

traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The compounds of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of the compound of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage

ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

5 For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign
10 polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the antibodies by modifications such as, for example, lipidation.

The invention also provides a pharmaceutical pack or kit comprising one or more
15 containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. **Diagnosis and Imaging**

20 Labeled antibodies, and derivatives and analogs thereof, which specifically bind to a polypeptide of interest can be used for diagnostic purposes to detect, diagnose, or monitor diseases and/or disorders associated with the aberrant expression and/or activity of a polypeptide of the invention. The invention provides for the detection of aberrant expression of a polypeptide of interest, comprising (a) assaying the expression of the polypeptide of
25 interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide of interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of aberrant expression.

The invention provides a diagnostic assay for diagnosing a disorder, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide of interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a particular disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Antibodies of the invention can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell. Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (125I, 121I), carbon (14C), sulfur (35S), tritium (3H), indium (112In), and technetium (99Tc); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

One aspect of the invention is the detection and diagnosis of a disease or disorder associated with aberrant expression of a polypeptide of interest in an animal, preferably a mammal and most preferably a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled molecule which specifically binds to the polypeptide of interest; b) waiting for a time interval following the administering for permitting the labeled molecule to preferentially concentrate at sites in the subject where the polypeptide is expressed (and for unbound labeled molecule to be cleared to background level); c) determining

background level; and d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with aberrant expression of the polypeptide of interest. Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).

Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

In an embodiment, monitoring of the disease or disorder is carried out by repeating the method for diagnosing the disease or disease, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

Presence of the labeled molecule can be detected in the patient using methods known in the art for in vivo scanning. These methods depend upon the type of label used. Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but

are not limited to, computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI). Kits

The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises an antibody of the invention, preferably a purified antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated polypeptide comprising an epitope which is specifically immunoreactive with an antibody included in the kit. Preferably, the kits of the present invention further comprise a control antibody which does not react with the polypeptide of interest. In another specific embodiment, the kits of the present invention contain a means for detecting the binding of an antibody to a polypeptide of interest (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate).

In another specific embodiment of the present invention, the kit is a diagnostic kit for use in screening serum containing antibodies specific against proliferative and/or cancerous polynucleotides and polypeptides. Such a kit may include a control antibody that does not react with the polypeptide of interest. Such a kit may include a substantially isolated polypeptide antigen comprising an epitope which is specifically immunoreactive with at least one anti-polypeptide antigen antibody. Further, such a kit includes means for detecting the binding of said antibody to the antigen (e.g., the antibody may be conjugated to a fluorescent

compound such as fluorescein or rhodamine which can be detected by flow cytometry). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized polypeptide antigen. The polypeptide antigen of the kit may also be attached to a solid support.

5 In a more specific embodiment the detecting means of the above-described kit includes a solid support to which said polypeptide antigen is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to the polypeptide antigen can be detected by binding of the said reporter-labeled antibody.

10 In an additional embodiment, the invention includes a diagnostic kit for use in screening serum containing antigens of the polypeptide of the invention. The diagnostic kit includes a substantially isolated antibody specifically immunoreactive with polypeptide or polynucleotide antigens, and means for detecting the binding of the polynucleotide or polypeptide antigen to the antibody. In one embodiment, the antibody is attached to a solid
15 support. In a specific embodiment, the antibody may be a monoclonal antibody. The detecting means of the kit may include a second, labeled monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

 In one diagnostic configuration, test serum is reacted with a solid phase reagent having a surface-bound antigen obtained by the methods of the present invention. After binding with
20 specific antigen antibody to the reagent and removing unbound serum components by washing, the reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-antigen antibody on the solid support. The reagent is again washed to remove unbound labeled antibody, and the amount of reporter associated with the reagent is determined. Typically, the reporter is an enzyme which is
25 detected by incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate (Sigma, St. Louis, MO).

 The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip sticks, 96-

well plate or filter material. These attachment methods generally include non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group. Alternatively, streptavidin coated plates can
5 be used in conjunction with biotinylated antigen(s).

Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound recombinant antigens, and a reporter-labeled anti-human antibody for detecting surface-bound anti-antigen antibody.

10 *Formulations*

The PGRP-K, PGRP-W, or PGRP-C polypeptide compositions (preferably containing a polypeptide which is a soluble form of the extracellular domain), respectively,
15 will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with either PGRP-K, PGRP-W, or PGRP-C polypeptide alone), the site of delivery of the PGRP-K, PGRP-W, and/or PGRP-C polypeptide composition, the method of administration, the scheduling of administration, and other factors known to practitioners.
20 The "effective amount" of PGRP-K, PGRP-W, and/or PGRP-C polypeptide for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of PGRP-K, PGRP-W, and/or PGRP-C polypeptide administered parenterally per dose will be in the range of about 1 $\mu\text{g/kg/day}$ to 10 mg/kg/day of patient body weight, although, as noted above,
25 this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day , and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the PGRP-K, PGRP-W, or PGRP-C polypeptide is typically administered at a dose rate of about 1 $\mu\text{g/kg/hour}$ to about 50 $\mu\text{g/kg/hour}$, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-

pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing the PGRP-K, PGRP-W, or PGRP-C of the invention may be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray. By "pharmaceutically acceptable carrier" is meant a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The PGRP-K, PGRP-W and PGRP-C polypeptides are also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., *Biopolymers* 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (R. Langer et al., *J. Biomed. Mater. Res.* 15:167-277 (1981), and R. Langer, *Chem. Tech.* 12:98-105 (1982)), ethylene vinyl acetate (R. Langer et al., Id.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release PGRP-K, PGRP-W, and PGRP-C polypeptide compositions also include liposomally entrapped PGRP-K, PGRP-W, and PGRP-C polypeptides. Liposomes containing PGRP-K, PGRP-W, and/or PGRP-C polypeptides are prepared by methods known per se: DE 3,218,121; Epstein et al., *Proc. Natl. Acad. Sci. (USA)* 82:3688-3692 (1985); Hwang et al., *Proc. Natl. Acad. Sci. (USA)* 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent

cholesterol, the selected proportion being adjusted for the optimal PGRP-K, PGRP-W, or PGRP-C polypeptide therapy.

For parenteral administration, in one embodiment, the PGRP-K, PGRP-W, and PGRP-C polypeptides are formulated generally by mixing them, respectively, at the desired
5 degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

10 Generally, the formulations are prepared by contacting either the PGRP-K, PGRP-W, or PGRP-C polypeptide uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's
15 solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate,
20 succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose
25 or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

The PGRP-K, PGRP-W, or PGRP-C polypeptide is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of PGRP-K, PGRP-W, or PGRP-C polypeptide salts.

PGRP-K, PGRP-W, or PGRP-C polypeptide to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic PGRP-K, PGRP-W, or PGRP-C polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

PGRP-K, PGRP-W, or PGRP-C polypeptides ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous PGRP-K, PGRP-W, or PGRP-C polypeptide solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized PGRP-K, PGRP-W, or PGRP-C polypeptide using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the polypeptides of the present invention may be employed in conjunction with other therapeutic compounds.

Agonists and Antagonists - Assays and Molecules

The invention also provides a method of screening compounds to identify those which enhance or block the action of PGRP-K, PGRP-W, or PGRP-C on cells, such as its interaction with either PGRP-K, PGRP-W, or PGRP-C binding molecules such as receptor
5 molecules. An agonist is a compound which increases the natural biological functions of PGRP-K, PGRP-W, or PGRP-C or which functions in a manner similar to PGRP-K, PGRP-W, or PGRP-C while antagonists decrease or eliminate such functions.

In another aspect of this embodiment the invention provides a method for identifying a receptor protein or other ligand-binding protein which binds specifically to a PGRP-K,
10 PGRP-W, or PGRP-C polypeptide. For example, a cellular compartment, such as a membrane or a preparation thereof, may be prepared from a cell that expresses a molecule that binds PGRP-K, PGRP-W, or PGRP-C. The preparation is incubated with labeled PGRP-K, PGRP-W, or PGRP-C and complexes of PGRP-K, PGRP-W, or PGRP-C, respectively, bound to the receptor or other binding proteins are isolated and characterized
15 according to routine methods known in the art. Alternatively, the PGRP-K, PGRP-W, or PGRP-C polypeptide may be bound to a solid support so that binding molecules solubilized from cells are bound to the column and then eluted and characterized according to routine methods.

In the assay of the invention for agonists or antagonists, a cellular compartment, such
20 as a membrane or a preparation thereof, may be prepared from a cell that expresses a molecule that binds PGRP-K, PGRP-W, or PGRP-C such as a molecule of the immune system, such as a macrophage or a monocyte. The preparation is incubated with labeled PGRP-K, PGRP-W, or PGRP-C in the absence or the presence of a candidate molecule which may be a PGRP-K, PGRP-W, or PGRP-C agonist or antagonist. The ability of the candidate molecule to bind the
25 binding molecule is reflected in decreased binding of the labeled ligand. Molecules which bind gratuitously, i.e., without inducing the effects of PGRP-K, PGRP-W, or PGRP-C on binding the PGRP-K, PGRP-W, or PGRP-C binding molecule, are most likely to be good antagonists.

Molecules that bind well and elicit effects that are the same as or closely related to PGRP-K, PGRP-W, or PGRP-C are agonists.

PGRP-K, PGRP-W, or PGRP-C-like effects of potential agonists and antagonists may be measured, for instance, by determining activity of a second messenger system following interaction of the candidate molecule with a cell or appropriate cell preparation, and comparing the effect with that of PGRP-K, PGRP-W, or PGRP-C or molecules that elicit the same effects as PGRP-K, PGRP-W, or PGRP-C. Second messenger systems that may be useful in this regard include but are not limited to AMP guanylate cyclase, ion channel or phosphoinositide hydrolysis second messenger systems.

Another example of an assay for PGRP-K, PGRP-W, or PGRP-C antagonists is a competitive assay that combines PGRP-K, PGRP-W, or PGRP-C and a potential antagonist with membrane-bound receptor molecules or recombinant PGRP-K, PGRP-W, or PGRP-C receptor molecules under appropriate conditions for a competitive inhibition assay. PGRP-K, PGRP-W, or PGRP-C can be labeled, such as by radioactivity, such that the number of PGRP-K, PGRP-W, or PGRP-C molecules bound to a receptor molecule can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a receptor molecule, without inducing PGRP-K, PGRP-W, or PGRP-C induced activities, thereby preventing the action of PGRP-K, PGRP-W, or PGRP-C by excluding PGRP-K, PGRP-W, or PGRP-C from binding.

Other potential antagonists include antisense molecules. Antisense technology can be used to control gene expression through antisense DNA or RNA or through triple-helix formation. Antisense techniques are discussed, for example, in Okano, J. *Neurochem.* 56: 560 (1991); "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance Lee et al., *Nucleic*

Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA. For example, the 5' coding portion of a polynucleotide that encodes the PGRP-like domain of one of the polypeptides of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of PGRP-K, PGRP-W, or PGRP-C, respectively. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into either the PGRP-K, PGRP-W, or PGRP-C polypeptide. The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of PGRP-K, PGRP-W, or PGRP-C.

The agonists and antagonists may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as described above.

The antagonists may be employed for instance to inhibit PGRP-K, PGRP-W, or PGRP-C chemotaxis and activation of macrophages and their precursors, and of neutrophils, monocytes, basophils, B lymphocytes and some T-cell subsets, e.g., activated and CD8 cytotoxic T cells and natural killer cells, in certain auto-immune and chronic inflammatory and infective diseases. Examples of auto-immune diseases include multiple sclerosis, and insulin-dependent diabetes. The antagonists may also be employed to treat infectious diseases including silicosis, sarcoidosis, idiopathic pulmonary fibrosis by preventing the recruitment and activation of mononuclear phagocytes. They may also be employed to treat idiopathic hyper-eosinophilic syndrome by preventing eosinophil production and migration. Endotoxic shock may also be treated by the antagonists by preventing the migration of macrophages and their production of the human chemokine polypeptides of the present invention. The antagonists may also be employed for treating atherosclerosis, by preventing monocyte infiltration in the artery wall. The antagonists may also be employed to treat histamine-mediated allergic reactions and immunological disorders including late phase allergic reactions,

chronic urticaria, and atopic dermatitis by inhibiting chemokine-induced mast cell and basophil degranulation and release of histamine. IgE-mediated allergic reactions such as allergic asthma, rhinitis, and eczema may also be treated. The antagonists may also be employed to treat chronic and acute inflammation by preventing the attraction of monocytes to a wound area. They may also be employed to regulate normal pulmonary macrophage populations, since chronic and acute inflammatory pulmonary diseases are associated with sequestration of mononuclear phagocytes in the lung. Antagonists may also be employed to treat rheumatoid arthritis by preventing the attraction of monocytes into synovial fluid in the joints of patients. Monocyte influx and activation plays a significant role in the pathogenesis of both degenerative and inflammatory arthropathies. The antagonists may be employed to interfere with the deleterious cascades attributed primarily to IL-1 and TNF, which prevents the biosynthesis of other inflammatory cytokines. In this way, the antagonists may be employed to prevent inflammation. The antagonists may also be employed to inhibit prostaglandin-independent fever induced by chemokines. The antagonists may also be employed to treat cases of bone marrow failure, for example, aplastic anemia and myelodysplastic syndrome. The antagonists may also be employed to treat asthma and allergy by preventing eosinophil accumulation in the lung. The antagonists may also be employed to treat subepithelial basement membrane fibrosis which is a prominent feature of the asthmatic lung.

Antibodies against PGRP-K, PGRP-W, or PGRP-C may be employed to bind to and inhibit PGRP-K, PGRP-W, or PGRP-C activity to treat ARDS, by preventing infiltration of neutrophils into the lung after injury. The antagonists may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as hereinafter described.

Prophylactic and Therapeutic Methods

It is to be understood that although the following discussion is specifically directed to human patients, the teachings are also applicable to any animal that expresses PGRP-K, PGRP-W, or PGRP-C.

PGRP-K, PGRP-W, or PGRP-C polypeptides or polynucleotides (including PGRP-K, PGRP-W, or PGRP-C fragments, variants, derivatives, and analogs, and PGRP-K, PGRP-W, or PGRP-C agonists and antagonists as described herein) are useful to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, PGRP-K, PGRP-W, or PGRP-C polypeptides or polynucleotides and/or PGRP-K, PGRP-W, or PGRP-C agonists or antagonists may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

PGRP-K, PGRP-W, or PGRP-C polypeptides or polynucleotides (including PGRP-K, PGRP-W, or PGRP-C fragments, variants, derivatives, and analogs, and PGRP-K, PGRP-W, or PGRP-C agonists and antagonists as described herein) are useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, PGRP-K, PGRP-W, or PGRP-C polynucleotides or polypeptides can be used as a marker or detector of a particular immune system disease or disorder.

Similarly, PGRP-K, PGRP-W, or PGRP-C polypeptides or polynucleotides (including PGRP-K, PGRP-W, or PGRP-C fragments, variants, derivatives, and analogs, and PGRP-K, PGRP-W, or PGRP-C agonists and PGRP-K, PGRP-W, or PGRP-C antagonists as described herein) are useful to modulate inflammation. For example, PGRP-K, PGRP-W, or PGRP-C polypeptides or polynucleotides and/or PGRP-K, PGRP-W, or PGRP-C agonists and antagonists of the invention may inhibit the proliferation and differentiation of cells involved in an inflammatory response or alternatively may be involved in killing of hematopoietic cells during processes of inflammation or tissue injury. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including ischemia-reperfusion injury, arthritis, and/or nephritis. Additionally, these molecules may be used to treat or prevent

killing of hematopoietic cells and/or other cells during processes of inflammation or tissue injury.

PGRP-K, PGRP-W, or PGRP-C polypeptides or polynucleotides (including PGRP-K, PGRP-W, or PGRP-C fragments, variants, derivatives, and analogs, and PGRP-K, PGRP-W, or PGRP-C agonists and antagonists as described herein) are useful to treat or detect hyperproliferative disorders, including neoplasms. PGRP-K, PGRP-W, or PGRP-C polypeptides or polynucleotides and/or PGRP-K, PGRP-W, or PGRP-C agonists or antagonists, may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, PGRP-K, PGRP-W, or PGRP-C polypeptides or polynucleotides and/or PGRP-K, PGRP-W, or PGRP-C agonists or antagonists may proliferate other cells which can inhibit the hyperproliferative disorder. For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

Given the activities modulated by PGRP-K, PGRP-W, or PGRP-C, it is readily apparent that a substantially altered (increased or decreased) level of expression of PGRP-K, PGRP-W, or PGRP-C in an individual compared to the standard or "normal" level produces pathological conditions such as those described above. It will also be appreciated by one of ordinary skill that the PGRP-K, PGRP-W, or PGRP-C agonists of the invention will exert modulating activities on any of its target cells. Therefore, it will be appreciated that conditions caused by a decrease in the standard or normal level of PGRP-K, PGRP-W, or PGRP-C mediated activity in an individual, can be treated by administration of PGRP-K, PGRP-W, or PGRP-C polypeptide or an agonist thereof.

Thus, in one embodiment, the present invention is directed to a method for enhancing (i.e., increasing) PGRP-K, PGRP-W, or PGRP-C mediated activity (e.g., immunity) which involves administering to an individual in need of an increased level of PGRP-K, PGRP-W, or PGRP-C mediated activity, a therapeutically effective amount of PGRP-K, PGRP-W, or PGRP-C polypeptide, fragment, variant, derivative, or analog, or an agonist capable of increasing PGRP-K, PGRP-W, or PGRP-C mediated activity. In specific embodiments,

PGRP-K, PGRP-W, or PGRP-C mediated signaling is increased to treat a disease or condition wherein decreased cell survival, secretion, proliferation, migration, and/or differentiation is exhibited.

In another embodiment, the present invention is directed to a method for suppressing (i.e., decreasing) PGRP-K, PGRP-W, or PGRP-C mediated activity (e.g., inflammation), which involves administering to an individual in need of a decreased level of PGRP-K, PGRP-W, or PGRP-C mediated activity, a therapeutically effective amount of PGRP-K, PGRP-W, or PGRP-C polypeptide, fragment, variant, derivative, or analog or an antagonist capable of decreasing PGRP-K, PGRP-W, or PGRP-C mediated activity. In specific embodiments, PGRP-K, PGRP-W, or PGRP-C mediated signaling is decreased to treat a disease or condition wherein increased cell survival, secretion, proliferation, migration and/or differentiation is exhibited.

In addition to treating diseases associated with elevated or decreased levels of PGRP-K, PGRP-W, or PGRP-C mediated activity, the invention encompasses methods of administering PGRP-K, PGRP-W, or PGRP-C agonists or antagonists to elevate or reduce PGRP-K, PGRP-W, or PGRP-C mediated biological activity, respectively.

For treating abnormal conditions related to an under-expression of PGRP-K, PGRP-W, or PGRP-C and its activity, or in which elevated or decreased levels of PGRP-K, PGRP-W, or PGRP-C are desired, several approaches are available. One approach comprises administering to an individual in need of an increased level of PGRP-K, PGRP-W, or PGRP-C mediated activity in the body, a therapeutically effective amount of an isolated PGRP-K, PGRP-W, or PGRP-C polypeptide, fragment, variant, derivative or analog of the invention, or a compound which activates PGRP-K, PGRP-W, or PGRP-C, i.e., an agonist as described above, optionally in combination with a pharmaceutically acceptable carrier. Alternatively, gene therapy may be employed to effect the endogenous production of PGRP-K, PGRP-W, or PGRP-C by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector using techniques known in the art. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral

particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For an overview of gene therapy, see Chapter 20, Gene Therapy and other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

Further, treatment can be administered, for example, in the form of gene replacement therapy. Specifically, one or more copies of a PGRP-K, PGRP-W, or PGRP-C nucleotide sequence of the invention that directs the production of a PGRP-K, PGRP-W, or PGRP-C gene product, respectively, exhibiting normal function, may be inserted into the appropriate cells within a patient or animal subject, using vectors which include, but are not limited to, adenovirus, adeno-associated virus, retrovirus and herpesvirus vectors, in addition to other particles that introduce DNA into cells, such as liposomes and gene activated matrices. Because the PGRP-K, PGRP-W, or PGRP-C gene is expressed in epithelial and endothelial tissues, such gene replacement techniques should be capable of delivering PGRP-K, PGRP-W, or PGRP-C gene sequence to these cells within patients, or, alternatively, should involve direct administration of such PGRP-K, PGRP-W, or PGRP-C polynucleotide sequences to the site of the cells in which the PGRP-K, PGRP-W, or PGRP-C gene sequences are to be expressed. Alternatively, targeted homologous recombination can be utilized to correct the defective endogenous PGRP-K, PGRP-W, or PGRP-C gene and/or regulatory sequences thereof (e.g., promoter and enhancer sequences), or alternatively, to "turn on" other dormant PGRP-K, PGRP-W, or PGRP-C activity in the appropriate tissue or cell type.

Additional methods which may be utilized to increase the overall levels of PGRP-K, PGRP-W, or PGRP-C expression and/or PGRP-K, PGRP-W, or PGRP-C activity include the introduction of appropriate PGRP-K, PGRP-W, or PGRP-C-expressing cells, preferably autologous cells, into a patient at positions and in numbers which are sufficient to ameliorate the symptoms of abnormalities in cell growth regulation, cell signaling, and other PGRP-K, PGRP-W, or PGRP-C mediated activities. Such cells may be either recombinant or non-recombinant. Among the cells which can be administered to increase the overall levels of PGRP-K, PGRP-W, or PGRP-C gene expression in a patient are normal cells, which express the PGRP-K, PGRP-W, or PGRP-C gene. Cell-based gene therapy techniques are well

known to those skilled in the art, see, e.g., Anderson et al., U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959.

Thus, one embodiment of the invention comprises administering to an individual in need of an increased level of PGRP-K, PGRP-W, or PGRP-C mediated activity a compound that stimulates PGRP-K, PGRP-W, or PGRP-C mediated activity (agonist), such as for example, an antibody or PGRP-K, PGRP-W, or PGRP-C fragment, variant, derivative or analog of the invention, along with a pharmaceutically acceptable carrier in an amount effective to enhance (i.e., increase) PGRP-K, PGRP-W, or PGRP-C mediated activity.

If the activity of PGRP-K, PGRP-W, or PGRP-C is in excess, several approaches are available to reduce or inhibit PGRP-K, PGRP-W, or PGRP-C activity using molecules derived from the polypeptide and polynucleotide sequences described above. Accordingly, a further aspect of the invention is related to a method for treating an individual in need of a decreased level of PGRP-K, PGRP-W, or PGRP-C mediated activity in the body comprising, administering to such an individual a composition comprising a therapeutically effective amount of a PGRP-K, PGRP-W, or PGRP-C polypeptide, fragment, variant, derivative or analog of the invention which acts as a PGRP-K, PGRP-W, or PGRP-C antagonist or PGRP-K, PGRP-W, or PGRP-C antagonist identified using the methods described herein, optionally, in combination with a pharmaceutically acceptable carrier. Preferably, PGRP-K, PGRP-W, or PGRP-C activity is decreased to treat a disease wherein increased cell survival, secretion, proliferation, migration, and/or differentiation is exhibited. Polypeptides, derivatives, variants and analogs of the invention and other compounds which function as antagonists of PGRP-K, PGRP-W, or PGRP-C can routinely be identified using the assays described *infra* and other techniques known in the art. Preferred antagonists for use in the present invention are PGRP-K, PGRP-W, or PGRP-C-specific antibodies.

In another approach, PGRP-K, PGRP-W, or PGRP-C activity can be reduced or inhibited by decreasing the level of PGRP-K, PGRP-W, or PGRP-C gene expression, respectively. In one embodiment, this is accomplished through the use of antisense sequences, either internally generated or separately administered (see, for example, O'Connor, J. Neurochem. (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Antisense technology can be used to control gene expression through antisense DNA or RNA or through triple-helix formation. Antisense

techniques are discussed, for example, in Okano, J. *Neurochem.* 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., *Nucleic Acids Research* 6:3073 (1979); Cooney et al., *Science* 241:456 (1988); and Dervan et al., *Science* 251:1360 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA. For example, the 5' coding portion of a polynucleotide that encodes a PGRP-K, PGRP-W, or PGRP-C polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the PGRP-K, PGRP-W, or PGRP-C polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into polypeptide.

In one embodiment, the PGRP-K, PGRP-W, or PGRP-C antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the PGRP-K, PGRP-W, or PGRP-C antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding PGRP-K, PGRP-W, or PGRP-C, or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, *Nature* 29:304-310 (1981)), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell* 22:787-797 (1980)), the herpes thymidine promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445 (1981)), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature* 296:39-42 (1982)), etc.

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a PGRP-K, PGRP-W, or PGRP-C gene. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double stranded PGRP-K, PGRP-W, or PGRP-C antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to

hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a PGRP-K, PGRP-W, or PGRP-C RNA it may contain and still form a stable duplex (or triplex as the case may be), respectively. One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Potential PGRP-K, PGRP-W, or PGRP-C antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al., Science 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy PGRP-K, PGRP-W, or PGRP-C mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, Nature 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within the nucleotide sequences of PGRP-K, PGRP-W, or PGRP-C (Figures 1A-C; SEQ ID NO:1). Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of either the PGRP-K, PGRP-W, or PGRP-C mRNA; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts. DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. Since ribozymes, unlike antisense molecules are catalytic, a lower intracellular concentration is required for efficiency.

Endogenous PGRP-K, PGRP-W, or PGRP-C gene expression can also be reduced by inactivating or "knocking out" the PGRP-K, PGRP-W, or PGRP-C gene, respectively, or its promoter using targeted homologous recombination (e.g., see Smithies et al., Nature 317:330-234 (1985); Thomas et al., Cell 51:503-512 (1987); Thompson et al., Cell 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). Such approach can be adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors.

Alternatively, endogenous PGRP-K, PGRP-W, or PGRP-C gene expression can be reduced by targeted deoxyribonucleotide sequences complementary to the regulatory region of the PGRP-K, PGRP-W, or PGRP-C gene (i.e., the PGRP-K, PGRP-W, or PGRP-C promoters and/or enhancers), respectively, to form triple helical structures that prevent transcription of the PGRP-K, PGRP-W, or PGRP-C genes in target cells in the body, see

generally, Helene et al., Ann. N.Y. Acad. Sci. 660:27-36 (1992); Helene, C., Anticancer Drug Des., 6(6):569-584 (1991); and Maher, L.J., Bioassays 14(12):807-815 (1992)).

Thus, one embodiment of the invention comprises administering to an individual in need of a decreased level of PGRP-K, PGRP-W, or PGRP-C mediated activity, a PGRP-K, PGRP-W, or PGRP-C inhibitor compound (antagonist), such as for example, an antibody or PGRP-K, PGRP-W, or PGRP-C fragment, variant, derivative or analog of the invention, along with a pharmaceutically acceptable carrier in an amount effective to suppress (i.e., lower) PGRP-K, PGRP-W, or PGRP-C mediated activity.

Chromosome Assays

The nucleic acid molecules of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

In certain preferred embodiments in this regard, the cDNA herein disclosed is used to clone genomic DNA of a PGRP-K, PGRP-W, and/or PGRP-C protein gene. This can be accomplished using a variety of well known techniques and libraries, which generally are available commercially. The genomic DNA then is used for *in situ* chromosome mapping using well known techniques for this purpose.

In addition, in some cases, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the 3' untranslated region of the gene is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Fluorescence *in situ* hybridization ("FISH") of a cDNA clone to a metaphase

chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with probes from the cDNA as short as 50 or 60 bp. For a review of this technique, see Verma *et al.*, *Human Chromosomes: A Manual Of Basic Techniques*, Pergamon Press, New York (1988).

5 Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, *Mendelian Inheritance In Man*, available on-line through Johns Hopkins University, Welch Medical Library. The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified
10 through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

15 Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

20 Examples

Example 1: Isolation of PGRP-K, PGRP-W, and/or PGRP-C cDNA Clone(s) From the Deposited Sample(s)

25 The cDNA for PGRP-K (ATCC Accession No: 203564) is inserted into the Sal I and Not I multiple cloning site of pCMVSPORT 2.0 (Life Technologies). pCMVSPORT 2.0 contains an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. (See, for instance, Gruber, C. E., et al., *Focus* 15:59- (1993).)

30 The cDNA for PGRP-W (ATCC Accession No: 203563) is inserted into the Sal I and Not I multiple cloning site of pCMVSPORT 3.0 (Life Technologies). pCMVSPORT 3.0 contains an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. (See, for instance, Gruber, C. E., et al., *Focus* 15:59- (1993).)

The cDNA for PGRP-C (ATCC Accession No: 209683) is inserted into the EcoRI and Xho I multiple cloning site of Uni-Zap XR (Stratagene). Uni-Zap XR contains an ampicillin

resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. (See, for instance, Gruber, C. E., et al., *Focus* 15:59- (1993).)

Two approaches can be used to isolate either PGRP-K, PGRP-W, and/or PGRP-C from the deposited sample. First, a specific polynucleotide of SEQ ID NO:1, SEQ ID NO:3, and
5 SEQ ID NO:5, respectively, with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ³²P-g-ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid mixture is transformed into a suitable
10 host (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening
15 (e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

Alternatively, two primers of 17-20 nucleotides derived from both ends of either the SEQ ID NO:1, SEQ ID NO:3, or SEQ ID NO:5, respectively, (i.e., within the region of SEQ ID
20 NO:1 bounded by the 5' NT and the 3' NT of the clone) are synthesized and used to amplify the PGRP-K, PGRP-W, and/or PGRP-C cDNAs using the deposited cDNA plasmids as template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25 ul of reaction mixture with 0.5 ug of the above cDNA template. A convenient reaction mixture is 1.5-5 mM MgCl₂, 0.01% (w/v) gelatin, 20 uM each of dATP, dCTP, dGTP, dTTP,
25 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94 degree C for 1 min; annealing at 55 degree C for 1 min; elongation at 72 degree C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected
30 sequence by subcloning and sequencing the DNA product.

Several methods are available for the identification of the 5' or 3' non-coding portions of either the PGRP-K, PGRP-W, or PGRP-C genes which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are
35 well known in the art. For instance, a method similar to 5' RACE is available for generating

the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer
5 specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of either the PGRP-K, PGRP-W, or PGRP-C gene of interest is used to PCR amplify the 5' portion of the PGRP-K, PGRP-W, or PGRP-C full-length gene, respectively. This amplified product may then be sequenced and used to generate the full length gene.

This above method starts with total RNA isolated from the desired source, although poly-
10 A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved
15 RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The
20 resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the PGRP-K, PGRP-W, or PGRP-C gene, respectively.

Alternatively, a genomic clone comprising the human PGRP-K, PGRP-W, or PGRP-C coding exons can be isolated by screening a human genomic library as discussed *infra*. Once positive clones have been identified, the DNA inserts contained in the genomic clone can be
25 isolated, and the DNA sequenced. Once the DNA sequence has been determined, the utilization of a number of computer-based DNA sequence analysis programs, such as, for example, BLAST and GRAIL, will allow the identification of the coding exons and the non-coding introns associated with either the PGRP-K, PGRP-W, or PGRP-C gene, respectively, and hence the identification of any 5' portion of the PGRP-K, PGRP-W, or PGRP-C full-length
30 gene which may not have been previously present in the deposited clone.

Example 2: Isolation of PGRP-K, PGRP-W, or PGRP-C Genomic Clones

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using
35 primers selected for the cDNA sequences corresponding to SEQ ID NO:1, SEQ ID NO:3, or SEQ ID NO:5, according to the method described in Example 1. (See also, Sambrook.)

Example 3: Chromosomal Mapping of PGRP-K, PGRP-W, or PGRP-C

An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:1, SEQ ID NO:3, or SEQ ID NO:5. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95 degree C; 1 minute, 56 degree C; 1 minute, 70 degree C. This cycle is repeated 32 times followed by one 5 minute cycle at 70 degree C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

Example 4: Bacterial Expression of PGRP-K, PGRP-W, or PGRP-C

PGRP-K, PGRP-W, or PGRP-C polynucleotides encoding PGRP-K, PGRP-W, or PGRP-C polypeptides, respectively, of the invention are amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as Sal I and Not I, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, Sal I and Not I correspond to the restriction enzyme sites on the bacterial expression vector pQE-9 (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

To clone either the PGRP-K, PGRP-W, or PGRP-C polypeptide in a bacterial vector, a 5' primer including a restriction site shared by the bacterial vector of interest, and including a number of nucleotides of the amino terminal coding sequence of the sequence of interest, is designed and synthesized. Likewise, a 3' primer including a restriction site shared by the bacterial vector of interest, and including nucleotides complementary to the 3' end of the coding sequence of the sequence of interest, is designed and synthesized. It would be obvious to one skilled in the art as to how to design the primers of interest. The primers are synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported.

For example, specifically, to clone the PGRP-K polypeptide in a bacterial vector, the 5' primer is easily designed by one skilled in the art to clone the PGRP-K polypeptide in a bacterial vector, the 5' primer has the sequence 5'

GCAGCACATATGGATTCTCCTGGAACAAAACACAAGCTAAAC 3' (SEQ ID NO: 8) containing the underlined NdeI restriction site followed by a number of nucleotides of the amino terminal coding sequence of the full-length PGRP-K sequence in SEQ ID NO:1. One of ordinary skill in the art would appreciate, of course, that the point in the protein coding sequence where the 5' primer begins may be varied to amplify a DNA segment encoding any desired portion of the complete PGRP-K protein shorter or longer than the full-length form of the protein. The 3' primer has the sequence 5' GCAGCAGGTACCTTAGTGTTTGAATGAGGCCAGGTGCTGATGATG 3' (SEQ ID NO: 9) containing the underlined Asp718 restriction site followed by a number of nucleotides complementary to the 3' end of the coding sequence of the PGRP-K DNA sequence of SEQ ID NO:1. PGRP-W and PGRP-C can also be cloned into a bacterial vector using primers and restriction sites specific to those proteins.

The pQE-9 vector is digested with NdeI and Asp718 and the amplified fragment (of either PGRP-K, PGRP-W, or PGRP-C) is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the *E. coli* strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalactopyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg).

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements operatively linked to a PGRP-K, PGRP-W, or PGRP-C polynucleotide, called pHE4a. (ATCC Accession Number 209645, deposited February 25, 1998.) This vector contains: 1) a neomycinphosphotransferase gene as a selection marker, 2) an *E. coli* origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (lacIq). The origin of replication (oriC) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHE4a by restricting the vector with NdeI and KpnI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

Example 5: Cloning and Expression of PGRP-K, PGRP-W, or PGRP-C in a Baculovirus Expression System

In this example, the plasmid shuttle vector pA2 is used to insert either the PGRP-K, PGRP-W, or PGRP-C polynucleotide into a baculovirus to express PGRP-K, PGRP-W, or PGRP-C, respectively. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that expresses the cloned PGRP-K, PGRP-W, or PGRP-C polynucleotide, respectively.

Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

Specifically, the PGRP-K, PGRP-W, or PGRP-C cDNA sequence contained in the deposited clones, including the AUG initiation codon and any naturally associated leader sequence, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

To clone either the PGRP-K, PGRP-W, or PGRP-C polypeptide in the baculovirus vector of interest, a 5' primer including a restriction site shared by the baculovirus vector of interest, an efficient signal for initiation of translation in eukaryotic cells (Kozak, M., *J. Mol. Biol.* 196:947-950 (1987)), and including a number of nucleotides of the amino terminal coding sequence of the sequence of interest, is designed and synthesized. Likewise, a 3' primer including a restriction site shared by the bacterial vector of interest, and including nucleotides complementary to the 3' sequence of the sequence of interest, is designed and synthesized. It would be obvious to one skilled in the art as to how to design the primers of interest. The primers are synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported.

For example, specifically, the cDNA sequence encoding the PGRP-K protein in the deposited clone shown in SEQ ID NO:1, is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer has the sequence 5' GCAGCAGGATCCGCCATCATGGGGACGCTGCCATGGCTTCTTGCCTTC 3' (SEQ ID NO: 10) containing the BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells (Kozak, M., *J. Mol. Biol.* 196:947-950 (1987)), followed by a number of nucleotides of the sequence of the PGRP-K protein shown in Figures 1A-B. The 3' primer has the sequence 5' GCAGCAGGTACCTTATTGATATCCAATGTCACAAAAGTTCCGTGTG 3' (SEQ ID NO: 11) containing the KpnI restriction site followed by a number of nucleotides complementary to the 3' sequence in Figures 1A-B.

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five ug of a plasmid containing the polynucleotide is co-transfected with 1.0 ug of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA",

Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One ug of BaculoGold™ virus DNA and 5 ug of the plasmid are mixed in a sterile well of a microtiter plate containing 50 ul of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 ul Lipofectin plus 90 ul Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27 degrees C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27 degrees C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 ul of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4 degree C.

To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 uCi of ³⁵S-methionine and 5 uCi ³⁵S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced PGRP-K, PGRP-W, or PGRP-C polypeptide.

Example 6: Expression of PGRP-K, PGRP-W, or PGRP-C in Mammalian Cells

PGRP-K, PGRP-W, or PGRP-C polypeptides can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV, HIV and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2DHFR (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human HeLa, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, PGRP-K, PGRP-W, or PGRP-C polypeptides can be expressed in stable cell lines containing either the PGRP-K, PGRP-W, or PGRP-C polynucleotide integrated into a chromosome, respectively. The co-transfection with a selectable marker such as DHFR, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected PGRP-K, PGRP-W, or PGRP-C gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

Derivatives of the plasmid pSV2-DHFR (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of either PGRP-K, PGRP-W, or PGRP-C.

The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC4 is digested with BamHI and KpnI and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector
5 is then isolated from a 1% agarose gel.

The cDNA sequence encoding either the PGRP-K, PGRP-W, or PGRP-C protein in the respective deposited clone is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer sequence contains an appropriate restriction site, an efficient signal for initiation of translation in eukaryotic cells (Kozak, M., *J. Mol. Biol.*
10 196:947-950 (1987)), followed by a number of nucleotides of the sequence of the PGRP-K, PGRP-W, or PGRP-C sequence. The 3' primer sequence contains an appropriate restriction site followed by a number of nucleotides complementary to the 3' sequence of the PGRP-K, PGRP-W, or PGRP-C sequence.

For example, specifically, the PGRP-K 5' primer has the sequence
15 5' GCAGCAGGATCCGCCATCATGGGGACGCTGCCATGGCTTCTTGCCTTC 3' (SEQ ID NO: 12) containing the BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells (Kozak, M., *J. Mol. Biol.* 196:947-950 (1987)), followed by a number of nucleotides of the sequence of the PGRP-K protein shown in Figures 1A-B. In this embodiment, the 3' primer has the sequence 5'
20 GCAGCAGGTACCTTATTGATATCCAATGTCACAAAAGTTCCGTGTG 3' (SEQ ID NO: 13) containing the KpnI restriction site followed by a number of nucleotides complementary to the 3' sequence in Figures 1A-B. It would be obvious to one skilled in the art as to how to design the primers of interest. The primers are synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported.

25 If a naturally occurring signal sequence is used to produce a secreted protein, the vector does not need a second signal peptide. Alternatively, if a naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence in an effort to secrete the protein from the cell. (See, e.g., WO 96/34891.)

The amplified fragment is then digested with the BamHI and KpnI and purified on a 1%
30 agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five
35 µg of the expression plasmid pC4 is cotransfected with 0.5 µg of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable

marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 uM, 2 uM, 5 uM, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 uM. Expression of EITHER PGRP-K, PGRP-W, OR PGRP-C is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

Example 7: Construction of N-Terminal and/or C-Terminal Deletion Mutants

The following general approach may be used to clone a N-terminal or C-terminal deletion PGRP-K, PGRP-W, or PGRP-C deletion mutant. Generally, two oligonucleotide primers of about 15-25 nucleotides are derived from the desired 5' and 3' positions of a polynucleotide of SEQ ID NO:1. The 5' and 3' positions of the primers are determined based on the desired PGRP-K, PGRP-W, or PGRP-C polynucleotide fragment. An initiation and stop codon are added to the 5' and 3' primers respectively, if necessary, to express the PGRP-K, PGRP-W, or PGRP-C polypeptide fragment encoded by the polynucleotide fragment. Preferred PGRP-K, PGRP-W, or PGRP-C polynucleotide fragments are those encoding the N-terminal and C-terminal deletion mutants disclosed above in the "Polynucleotide and Polypeptide Fragments" section of the Specification.

Additional nucleotides containing restriction sites to facilitate cloning of the PGRP-K, PGRP-W, or PGRP-C polynucleotide fragment in a desired vector may also be added to the 5' and 3' primer sequences. The PGRP-K, PGRP-W, or PGRP-C polynucleotide fragment is amplified from genomic DNA or from the deposited cDNA clone using the appropriate PCR oligonucleotide primers and conditions discussed herein or known in the art. The PGRP-K, PGRP-W, or PGRP-C polypeptide fragments encoded by the PGRP-K, PGRP-W, or PGRP-C polynucleotide fragments, respectively, of the present invention may be expressed and purified in the same general manner as the full length polypeptides, although routine modifications may be necessary due to the differences in chemical and physical properties between a particular fragment and full length polypeptide.

As a means of exemplifying but not limiting the present invention, the polynucleotide encoding the PGRP-K polypeptide fragment Met-1 to Ile-155 is amplified and cloned as

follows: A 5' primer is generated comprising a restriction enzyme site followed by an initiation codon in frame with the polynucleotide sequence encoding the N-terminal portion of the polypeptide fragment beginning with Met-1. A complementary 3' primer is generated comprising a restriction enzyme site followed by a stop codon in frame with the polynucleotide sequence encoding C-terminal portion of the PGRP-K polypeptide fragment ending with Ile-155.

The amplified polynucleotide fragment and the expression vector are digested with restriction enzymes which recognize the sites in the primers. The digested polynucleotides are then ligated together. The PGRP-K polynucleotide fragment is inserted into the restricted expression vector, preferably in a manner which places the PGRP-K polypeptide fragment coding region downstream from the promoter. The ligation mixture is transformed into competent *E. coli* cells using standard procedures and as described in the Examples herein. Plasmid DNA is isolated from resistant colonies and the identity of the cloned DNA confirmed by restriction analysis, PCR and DNA sequencing.

Example 8: Protein Fusions of PGRP-K, PGRP-W, or PGRP-C

PGRP-K, PGRP-W, or PGRP-C polypeptides are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of PGRP-K, PGRP-W, or PGRP-C polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the half-life time in vivo. Nuclear localization signals fused to PGRP-K, PGRP-W, or PGRP-C polypeptides can target the protein(s) to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule..

Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector,

and the PGRP-K, PGRP-W, or PGRP-C polynucleotide, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

Human IgG Fc region:

10 GGGATCCGGAGCCCAAATCTTCTGACAAAACCTCACACATGCCCACCGTGCCCAGCA
CCTGAATTCGAGGGTGCACCGTCAGTCTTCTCTCCCCCAAACCCAAGGACAC
CCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGTGGTGGACGTAAGCCAC
GAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATG
CCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCG
15 TCCTCACCGTCTCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGT
CTCCAACAAAGCCCTCCCAACCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGC
AGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCAA
GAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAAGCGACATCGCC
GTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCC
20 GTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAG
CAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCAC
AACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAATGAGTGCGACGGC
CGCGACTCTAGAGGAT (SEQ ID NO:14)

25 Example 9: Production of an Antibody

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) For example, cells expressing PGRP-K, PGRP-W, or PGRP-C will be administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of PGRP-K, PGRP-W, or PGRP-C protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology. (Köhler et al., Nature 256:495 (1975); Köhler et al., Eur. J.

Immunol. 6:511 (1976); Köhler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures involve immunizing an animal (preferably a mouse) with the PGRP-K, PGRP-W, or PGRP-C polypeptide or, more preferably, with a secreted PGRP-K, PGRP-W, or PGRP-C polypeptide-expressing cell. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56 degree C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 ug/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the PGRP-K, PGRP-W, or PGRP-C polypeptide, respectively.

Alternatively, additional antibodies capable of binding to the PGRP-K, PGRP-W, or PGRP-C polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the PGRP-K, PGRP-W, or PGRP-C specific antibody can be blocked by PGRP-K, PGRP-W, or PGRP-C, respectively. Such antibodies comprise anti-idiotypic antibodies to the PGRP-K, PGRP-W, or PGRP-C specific antibody and can be used to immunize an animal to induce formation of further PGRP-K, PGRP-W, or PGRP-C specific antibodies, respectively.

It will be appreciated that Fab and F(ab')₂ and other fragments of the antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). Alternatively, secreted PGRP-K, PGRP-W, or PGRP-C protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic chemistry.

For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs

derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

Example 10: Method of Detecting Abnormal Levels of PGRP-K, PGRP-W, or PGRP-C in a Biological Sample

PGRP-K, PGRP-W, or PGRP-C polypeptides can be detected in a biological sample, and if an increased or decreased level of PGRP-K, PGRP-W, or PGRP-C is detected, the respective polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

For example, antibody-sandwich ELISAs are used to detect PGRP-K, PGRP-W, or PGRP-C in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies to PGRP-K, PGRP-W, or PGRP-C, respectively, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 11. The wells are blocked so that non-specific binding of PGRP-K, PGRP-W, or PGRP-C to their respective well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing PGRP-K, PGRP-W, or PGRP-C. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded PGRP-K, PGRP-W, or PGRP-C.

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot the PGRP-K, PGRP-W, or PGRP-C polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of PGRP-K, PGRP-W, or PGRP-C in the sample using the standard curve.

Example 11: Formulating a Polypeptide

The PGRP-K, PGRP-W, or PGRP-C composition will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with either the PGRP-K, PGRP-W, or PGRP-C polypeptide alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of PGRP-K, PGRP-W, or PGRP-C administered parenterally per dose will be in the range of about 1 µg/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, PGRP-K, PGRP-W, or PGRP-C is typically administered at a dose rate of about 1 µg/kg/hour to about 50 µg/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing PGRP-K, PGRP-W, or PGRP-C are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

PGRP-K, PGRP-W, or PGRP-C is also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., Biopolymers 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (R. Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and R. Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (R. Langer et al.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also include liposomally entrapped PGRP-K, PGRP-W, or PGRP-C polypeptides. Liposomes containing either the PGRP-K, PGRP-W, or PGRP-C are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. USA 82:3688-

3692 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal secreted polypeptide therapy.

For parenteral administration, in one embodiment, PGRP-K, PGRP-W, or PGRP-C is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

Generally, the formulations are prepared by contacting either PGRP-K, PGRP-W, or PGRP-C uniformly and intimately with liquid carriers or finely divided solid carriers, or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

PGRP-K, PGRP-W, or PGRP-C is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

PGRP-K, PGRP-W, or PGRP-C used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2

micron membranes). Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

PGRP-K, PGRP-W, or PGRP-C polypeptides ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous PGRP-K, PGRP-W, or PGRP-C polypeptide solution, respectively, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized PGRP-K, PGRP-W, or PGRP-C polypeptide using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, PGRP-K, PGRP-W, or PGRP-C may be employed in conjunction with other therapeutic compounds.

Example 12: Method of Treatment Using Gene Therapy - In Vivo

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) PGRP-K, PGRP-W, or PGRP-C sequences into an animal to increase or decrease the expression of the PGRP-K, PGRP-W, or PGRP-C polypeptide, respectively. The PGRP-K, PGRP-W, or PGRP-C polynucleotide may be operatively linked to a promoter or any other genetic elements necessary for the expression of either the PGRP-K, PGRP-W, or PGRP-C polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151, 5580859; Tabata H. et al. (1997) Cardiovasc. Res. 35(3):470-479, Chao J et al. (1997) Pharmacol. Res. 35(6):517-522, Wolff J.A. (1997) Neuromuscul. Disord. 7(5):314-318, Schwartz B. et al. (1996) Gene Ther. 3(5):405-411, Tsurumi Y. et al. (1996) Circulation 94(12):3281-3290 (incorporated herein by reference).

The PGRP-K, PGRP-W, or PGRP-C polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The PGRP-

K, PGRP-W, or PGRP-C polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the PGRP-K, PGRP-W, or PGRP-C polynucleotides may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The PGRP-K, PGRP-W, or PGRP-C polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The PGRP-K, PGRP-W, or PGRP-C polynucleotide constructs can be delivered to the interstitial space of tissues within the an animal, including of pancreas, kidney, muscle, skeletal muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked PGRP-K, PGRP-W, or PGRP-C polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about

20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked PGRP-K, PGRP-W, or PGRP-C polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected PGRP-K, PGRP-W, or PGRP-C polynucleotide in muscle *in vivo* is determined as follows. Suitable PGRP-K, PGRP-W, or PGRP-C template DNA for production of mRNA coding for PGRP-K, PGRP-W, or PGRP-C polypeptides, respectively, is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The PGRP-K, PGRP-W, or PGRP-C template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 μ m cross-section of the individual quadriceps muscles is histochemically stained for PGRP-K, PGRP-W, or PGRP-C protein expression. A time course for PGRP-K, PGRP-W, or PGRP-C protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of PGRP-K, PGRP-W, or PGRP-C DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice will be used to extrapolate proper dosages and other treatment parameters in humans and other animals using PGRP-K, PGRP-W, or PGRP-C naked DNA.

Example 13: Peptidoglycan Binding Assay

Insoluble peptidoglycan is prepared from *Micrococcus luteus* as has been described in the art (Araki, Y., Nakatani, T., Nakayama, K. and Ito, E., 1972, J. Biol. Chem., 247:6312-632).

The peptidoglycan binding assay is performed by incubating 0.32 mgs of peptidoglycan in 280 uls of 10mM maleate buffer, pH 6.5/0.15 M NaCl with 3-6 ugs of a PGRP in 40 uls of 1 M imidazole/0.5 M NaCl/20 mM Tris-HCl, pH 7.9, for 30 min. at 4°C. One-sixteenth of the supernatant and of the pellet was removed for analysis. The protein was separated for the peptidoglycan by boiling in 2% SDS/PAGE loading buffer, subjected to electrophoresis on an SDS/15% polyacrylamide gel, and stained with Coomassie brilliant blue.

It can be appreciated by those skilled in the art that the above assay may be altered and/or refined to a degree to enhance visualization of the binding, while essentially maintaining the general scheme of the assay. Further peptidoglycan binding assays are described by Yoshida *et al.*, JBC, 271 (23): 13854 (1996), which is incorporated in its entirety by reference herein by reference.

Example 14: Measurement of Apoptosis ability of PGRP-K, PGRP-W, or PGRP-C

In a first incubation step, anti-histone antibody is fixed adsorptively on the wall of a microtiter plate module. Subsequently, non-specific binding sites on the wall are saturated by treatment with incubation buffer (e.g., blocking solution). During the second incubation step, the nucleosomes contained in the appropriate cell (e.g., WEHI 164 cells) sample treated with the PGRP-K, PGRP-W, or PGRP-C bind via their histone components to the immobilized anti-histone antibody. In the third incubation step, anti-DNA-peroxidase reacts with the DNA-part of the nucleosomes. After removal of all unbound peroxidase conjugate by a washing step, the amount of peroxidase retained in the immunocomplex is determined photometrically with ABTS (2,2'-azino-di-[3-ethylbenzthiazoline sulfonate]), as a substrate.

Anti-histone antibody reacts with the histones H1, H2A, H2B, H3 and H4 from the sample. Anti-DNA POD antibody binds to single- and double-stranded DNA. Therefore, the ELISA allows the detection on mono- and oligonucleosomes and may be applied to measure apoptotic cell death. The level of cell death is measured by the amount of cytoplasmic histone-associated DNA fragments which are indicated as the absorbance A405nm/A490. (See 5 Boehringer mannheim Catalogue, 0990 C 93 2 1541170).

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within 10 the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference.

Moreover, the sequence submitted herewith in paper and computer readable form are 15 herein incorporated by reference in their entireties.

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>5</u> , line <u>21</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 23 December 1998	Accession Number 203563
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") 	
<input checked="" type="checkbox"/> For receiving Office use only This sheet was received with the international application Authorized officer MELVIN L. BROOKS SR. INTERNATIONAL DIVISION 703-305-5163 msj	<input type="checkbox"/> For International Bureau use only This sheet was received by the International Bureau on: Authorized officer

ATCC Deposit No. 203563

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No. 203563

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>5</u> , line <u>21</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 23 December 1998	Accession Number 203564
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<p style="text-align: center;">For receiving Office use only</p> <p><input checked="" type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer: MELVIN S. BEJAN INTERNATIONAL DIVISION 703 305 5160</p>	<p style="text-align: center;">For International Bureau use only</p> <p><input type="checkbox"/> This sheet was received by the International Bureau on:</p> <p>Authorized officer:</p>
--	---

ATCC Deposit No. 203564

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No. 203564

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>5</u> , line <u>21</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 March 1998	Accession Number 209683
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<input checked="" type="checkbox"/> For receiving Office use only This sheet was received with the international application
Authorized officer MELVIN A. BROOKS SR. INTERNATIONAL DIVISION 703-305-5163 M.S.T.

<input type="checkbox"/> For International Bureau use only This sheet was received by the International Bureau on:
Authorized officer

ATCC Deposit No. 209683

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No. 209683

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

(a) a polynucleotide fragment of SEQ ID NO:1, SEQ ID NO:3, and/or SEQ ID NO:5 or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit Numbers: 203564, 203563, and 209683, respectively;

(b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:2, SEQ ID NO:4, and/or SEQ ID NO:6, or a polypeptide encoded by the cDNA sequence included in ATCC Deposit Numbers: 203564, 203563, and/or 209683, respectively;

(c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:2, SEQ ID NO:4, and/or SEQ ID NO:6, or a polypeptide domain encoded by the cDNA sequence included in ATCC Numbers: 203564, 203563, and/or 209683, respectively;

(d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:2, SEQ ID NO:4, and/or SEQ ID NO:6, or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit Numbers: 203564, 203563, and/or 209683, respectively;

(e) a polynucleotide encoding a polypeptide of SEQ ID NO:2, SEQ ID NO:4, and/or SEQ ID NO:6, or a polypeptide encoded by the cDNA sequence included in ATCC Deposit Numbers: 203564, 203563, and/or 209683, respectively, having biological activity; and

(f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding PGRP-K (SEQ ID NO:2), PGRP-W (SEQ ID NO:4), or PGRP-C (SEQ ID NO:6).

3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the polypeptides encoded by the cDNA sequences included in ATCC Deposit Numbers: 203564, 203563, or 209683, respectively.

4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or SEQ ID NO:5, or the cDNA sequence included in ATCC Deposit Numbers: 203564, 203563, or 209683, respectively.

5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions of nucleotides encoding either the C-terminus or the N-terminus.

6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions of nucleotides encoding either the C-terminus or the N-terminus.

7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.

8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.

9. A recombinant host cell produced by the method of claim 8.

10. The recombinant host cell of claim 9 comprising vector sequences.

11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

(a) a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:6, or the encoded sequence included in ATCC Deposit Numbers: 203564, 203563, or 209683, respectively.

(b) a polypeptide fragment of SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:6, or the encoded sequence included in ATCC Deposit Numbers: 203564, 203563, or 209683, respectively, having biological activity;

(c) a polypeptide domain of SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:6, or the encoded sequence included in ATCC Deposit Numbers: 203564, 203563, or 209683, respectively; and

(d) a polypeptide epitope of SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:6, or the encoded sequence included in ATCC Deposit Numbers: 203564, 203563, or 209683, respectively.

12. The isolated polypeptide of claim 11, wherein the polypeptide fragment comprises sequential amino acid deletions from either the C-terminus or the N-terminus.

13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.

14. A recombinant host cell that expresses the isolated polypeptide of claim 11.

15. A method of making an isolated polypeptide comprising:

(a) culturing the recombinant host cell of claim 14 under conditions such that said polypeptide is expressed; and

(b) recovering said polypeptide.

16. The polypeptide produced by claim 15.

17. A method for preventing, treating, or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of the polypeptide of claim 11 or of the polynucleotide of claim 1.

18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to expression or activity of PGRP-K, PGRP-W, or PGRP-C comprising:

(a) determining the presence or absence of a mutation in the polynucleotide of claim 1;

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

19. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to expression or activity of PGRP-K, PGRP-W, or PGRP-C comprising:

(a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample;

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

20. A method of identifying a binding partner to the polypeptide of claim 11 comprising:

- (a) contacting the polypeptide of claim 11 with a plurality of compounds; and
- (b) identifying compounds that bind the polypeptide.

21. A method of identifying compounds capable of enhancing or inhibiting a cellular response induced by PGRP-K, PGRP-W, or PGRP-C comprising:

- (a) contacting cells which express the polypeptide of claim 11, with a candidate compound; and
- (b) assaying a cellular response.

22. The method of claim 21 wherein the cellular response is the binding of insoluble peptidoglycan.

23. The method of claim 21 wherein the binding of insoluble peptidoglycan is assayed.

1/10

Figure 1A

1 CTGGGCTGGAACAGCACAGAACCCACAGGGCTGCCGTCCACACTCTCCCGGTCAGAGTCC 60
61 TGGGACCACATGGGGACGCTGCCATGGCTTCTTGCCCTTCTTCATTCTGGGTCTCCAGGCT 120
1 M G T L P W L L A F F I L G L Q A 17
121 TGGGATACTCCACCATCGTCTCCCGCAAGGAGTGGGGGGCAAGACCGCTCGCCTGCAGG 180
18 W D T P T I V S R K E W G A R P L A C R 37
181 GCCCTGCTGACCTTGCCTGTGGCCTACATCATCACAGACCAGCTCCCAGGGATGCAGTGC 240
38 A L L T L P V A Y I I T D Q L P G M Q C 57
241 CAGCAGCAGAGCGTTTGCAGCCAGATGCTGCCGGGGTTGCAGTCCCATTCGGTCTACACC 300
58 Q Q Q S V C S Q M L R G L Q S H S V Y T 77
301 ATAGGCTGGTGGCAGCTGGCGTACAACCTTCCTGGTTGGGGATGATGGCAGGGTGTATGAA 360
78 I G W C D V A Y N F L V G D D G R V Y E 97
361 GGTGTGGCTGGAACATCCAAGGCTTGCAACCCAGGGCTACAACAACATTTCCTGGGC 420
98 G V G W N I Q G L H T Q G Y N N I S L G 117
421 ATGCCCTTCTTTGGCAATAAGATAAGCAGCAGTCCCAGCCCTGCTGCCCTTATCAGCTGCA 480
118 I A F F G N K I S S S P S P A A L S A A 137
481 GAGGGTCTGATCTCTATGCCATCCAGAAGGGTCACCTGTGCCCCAGGTATATTGAGCCA 540
138 E G L I S Y A I Q K G H L S P R Y I Q P 157
541 CTTCTTCTGAAAGAAGAGACCTGCCCTGGACCCCTCAACATCCAGTGATGCCAGGAAGGTT 600
158 L L L K E E T C L D P Q H P V M P R K V 177
601 TGCCCCAACATCATCAACGATCTGCTTGGGAAGCCAGAGAGACACTGCCCTAAAATG 660
178 C P N I I K R S A W E A R E T H C P K M 197
661 AACCTCCCAGCCAAATATGTATCATCATCCACACCGCTGGCACAAGCTGCACTGTATCC 720
198 N L P A K Y V I I I H T A G T S C T V S 217
721 ACAGACTGCCAGACTGTGCTCCGAAACATACAGTCCCTTTCACATGGACACACGGAACCTTT 780
218 T D C Q T V V R N I Q S F H M D T R N F 237

2/10

Figure 1B

781	TGTGACATTGGATATCAATAAGGCCAGGCGTGGCGGCGATTACGTCTGTAATCCCAGGAC	840
238	C D I G Y Q	243
841	TTTGGGAGGCCAAGGCGGGCAGATCACTTCAGGCCAGGAATTCAAGAGCAGCCTGGCCAA	900
901	TATGGCGAACTCTGTCTCTACTGAAAACAAACAAACAAACAAACAAACAAAGAAA	960
961	CAACAAAAATTAGCCGGGTGTGGTGGCACACGCCTGTAGTCCCAGCTACTCAGGAGGCTG	1020
1021	AGGCATAAGAATTGCTTGAACCCCTGGAGGCGGAGGTTGCAGTGAGCTGAGATTGGGCCAC	1080
1081	CGCACTCCAGTCTGGGAGACAGAGTGAGACTGTCTCAAACAAACAACAAAAAATCCCTA	1140
1141	ACATAATCTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	1182

3/10

Figure 2A

1	GGGCAAGCTGACTGCACCCCTGACCTGCTGGGCTGGGACAGCACAGGACCCACAGATATCT	60
61	GCTGCCATCCACACTCTCCAGATTGGTGTCTGGGACCACTGGGGATGCTGCTGTGGCT	120
1	M L L W L	5
121	TCTTGCTCTCTGCTCTGGGTATCCAGGCCTGGGGTGATTCTCTCTGGAACAAACACA	180
6	L V F S A L G I Q A W G D S S W N K T Q	25
181	AGCTAAACAGGTATCAGAGGGGCTCCAGTACCTATTTGAGAACATCTCCAGCTCACTGA	240
26	A K Q V S E G L Q Y L F E N I S Q L T E	45
241	AAAAGATGTCTCCACCACGGTCTCTCGCAAGGCATGGGGGCAGAAGCTGTTGGCTGCAG	300
46	K D V S T T V S R K A W G A E A V G C S	65
301	TATTCAGCTGACCACGCCAGTGAATGTCCTTGTTATACACCATGTCCCTGGACTGGAGTG	360
66	I Q L T T P V N V L V I H H V P G L E C	85
361	TCACGACCAGACAGTCTGCAGCCAGAGACTGCGGGAAGTGCAGGCCCATGTGCCACAA	420
86	H D Q T V C S Q R L R E L Q A H H V H N	105
421	CAACAGTGGGTGTGATGTGGCCTACAACCTCCTGGTTGGGGATGATGGCAGGGTGATGA	480
106	N S G C D V A Y N F L V G D D G R V Y E	125
481	AGGTGTTGGCTGGAATATCCAAGGAGTGCACACCCAAGGCTACAACAACATCTCCCTGGG	540
126	G V G W N I Q G V H T Q G Y N N I S L G	145
541	CTTTGCCCTCTTCGGCACTAAGAAAGGCCACAGTCCCAGCCCTGCTGCCCTGTGGCCAT	600
146	F A F F G T K K G H S P S P A A L S A M	165
601	GGAAAACCTAATCACCTATGCTGTCCAGAAGGGCCACCTGTTCATCCAGTTATGTTTCAGCC	660
166	E N L I T Y A V Q K G H L S S S Y V Q P	185
661	ACTTCTTGGGAAAGGCGAGAAGTGCCTGGGCCCTCGGCAGAAGACAAGCCTGAAGAAGCT	720
186	L L G K G E N C L A P R Q K T S L K K L	205
721	TGCCCCGGCAATTGTCCACGGTCTGTGTGGGGAGCCAGGGAGACCACTGTCCAGGATGAC	780
206	A P A L S H G L C G E P G R P L S R M T	225

4/10

Figure 2B

781 TCTCCCAGCGAAGTATGGCATCATTATCCACACTGCCGGGAGGACCTGCAACATTTCCTGA 840
226 L P A K Y G I I I H T A G R T C N I S D 245

841 TGAGTGCCGCCTGCTGGTCCGGGACATCCAGTCTTTCTACATAGACAGGCTCAAGTCATG 900
246 E C R L L V R D I Q S F Y I D R L K S C 265

901 CGACATTGGTTATAACTTCCTGGTGGGCCAGGATGGCGCCATTATGAAGGGGTGGGCTG 960
266 D I G Y N F L V G Q D G A I Y E G V G W 285

961 GAATGTCCAAGGCTCCTCCACCCCTGGCTACGATGACATTGCCCTGGGCATTACCTTCAT 1020
286 N V Q G S S T P G Y D D I A L G I T F M 305

1021 GGGCACCTTCACAGGTATACCAACCAATGCTGCAGCACTAGAGGCAGCCCAAGACCTGAT 1080
306 G T F T G I P P N A A A L E A A Q D L I 325

1081 CCAGTGTGCCATGGTCAAAGGGTACCTGACTCCCAACTACCTGCTGGTGGGCCACAGTGA 1140
326 Q C A M V K G Y L T P N Y L L V G H S D 345

1141 TGTGGCCCGAACCTTGTCTCCTGGGCAGGCTTTGTACAACATCATCAGCACCTGGCCTCA 1200
346 V A R T L S P G Q A L Y N I I S T W P H 365

1201 TTTCAAACACTGAGAGAAGCCCCAGGTCTCTGAGACTGCTTTCCCTCCCTGTCAGGT 1260
366 F K H 368

1261 CTCTCTGTCTCTAACCATCCAGCTTGGCTCAACACCTTTTGCCCTCCTCCCTGCCACAC 1320

1321 AGTCCTGTGCTCTCTTTTTCAGGTGGGATGATCATGCCTCTCCTGCCAACATCCTCCAA 1380

1381 GGGCTCCAAACTCATAGCTGGACATTACAGCCCTCTGAGTCTGAGTCCAGATTTCCTTC 1440

1441 TCTCCTTACTTCTCTCCTTGGAAACCAACTCCTCAGCCAGGTGAGACAATGGGCTGG 1500

1501 TTCTTGTTTCATTTCCTCTCTCTCTCCATTCCTCTGCTGGTGAGCCTTCCCTGGT 1560

1561 GTCTGCCTGGCAGCCCCACCACCCACCTATCACCCCTCACCCATAACTCAGGTCAACGT 1620

5/10

Figure 2C

1621 GACCAACCTTCCTTGCTTACACATAAACTTGTATATATTGGATGTAGCCCTTATTTAAT 1680
1681 GGCTGTCATTATTTATAGATATGTCTATCCTTGCTACTTGGTTGTGAGTTTCTCCAGGGG 1740
1741 AGGAACTGTGTTTTATTCATCTCTATGTCTCTGTTTCTCAGCAGTGTCTGAAATTTAAT 1800
1801 GGGTTCCTACTGATGTTTATTAGAGAAATGGATGAATAAATGAATGAAGAGATCCAAAAA 1860
1861 AAAAAAAAAAAAAA 1876


6/10

Figure 3

1	GATCCCCCGGGCTGCAGGAATTCGGCAGCAGCCGGACCCTGCCGCCCTGCCACTATGTCC	60
1		2
	M S	
61	CGCCGCTCTATGCTGCTTGCCCTGGGCTCTCCCCAGCCTCCTTCGACTCGGAGCGGCTCAG	120
3	R R S M L L A W A L P S L L R L G A A Q	22
121	GAGACAGAAGACCCGGCCTGCTGCAGCCCCATAGTCCCCCGGAACGAGTGAAGGCCCTG	180
23	E T E D P A C C S P I V P R N E W K A L	42
181	GCA TCAGAGTGGCCCGCAGCACCTGAGCCTGCCCTTACGCTATGTGGTGGTATCGCACACG	240
43	A S E C A Q H L S L P L R Y V V V S H T	62
241	GCGGGCAGCAGCTGCAACACCCCGCCTCGTGCCAGCAGCAGGCCCGGAATGTGCAGCAC	300
63	A G S S C N T P A S C Q Q Q A R N V Q H	82
301	TACCACATGAAGACACTGGGCTGGTGGCAGCTGGGCTACAACCTCCTGATGGAGAAGAC	360
83	Y H M K T L G W C D V G Y N F L I G E D	102
361	GGGCTCGTATACGAGGGCGTGGCTGGAACCTTCAGGGTGCCCACTCAGGTCACTTATGG	420
103	G L V Y E G R G W N F T G A H S G H L W	122
421	AACCCCATGTCCATTGGCATCAGCTTCATGGGCAACTACATGGATCGGGTGCCACACCC	480
123	N P M S I G I S F M G N Y M D R V P T P	142
481	CAGGCCATCCGGGCAGCCAGGGTCTACTGGCCTGCCGTGTGGCTCAGGGAGCCCTGAGG	540
143	Q A I R A A Q G L L A C G V A Q G A L R	162
541	TCCAACATATGTGCTCAAAGGACACCGGATGTGCAGCGTACACTCTCTCCAGGCAACCAG	600
163	S N Y V L K G H R D V Q R T L S P G N Q	182
601	CTCTACCACCTCATCCAGAATTGGCCCACTACCGCTCCCCCTGAGGCCCTGTGATCCG	660
183	L Y H L I Q N W P H Y R S P	196
661	CACCCCATTCCTCCCTCCCATGGCCAAAAACCCCACTGTCTCCTTCTCCAATAAAGATG	720
721	TAGCTCAAAAAAAAAAAAAAAAAAAAAA	749


8/10

Figure 5




9/10

Figure 6



10/10

Figure 7



SEQUENCE LISTING

<110> Human Genome Sciences, Inc.

<120> Peptidoglycan Recognition Proteins

<130> PF492.PCT

<140> Unassigned

<141> 1999-12-20

<150> 60/113,809

<151> 1998-12-23

<160> 14

<170> PatentIn Ver. 2.0

<210> 1

<211> 1182

<212> DNA

<213> Homo sapiens

<400> 1

```

ctgggctgga acagcacaga acccacaggg ctgccgtcca cactctcccg gtcagagtcc 60
tgggaccaca tggggacgct gccatggctt ctgccttctt tcattctggg tctccaggct 120
tgggatactc ccaccatcgt ctcccgaag gaggggggg caagaccgct cgctgcagg 180
gccctgctga ccctgcctgt ggctacatc atcacagacc agtcccagg gatgcagtgc 240
cagcagcaga gcgtttgcag ccagatgctg cgggggttgc agtcccattc cgtctacacc 300
ataggctggt gcgacgtggc gtacaacttc ctggttgggg atgatggcag ggtgtatgaa 360
ggtgttggct ggaacatcca aggcttgcac acccagggct acaacaacat ttccctgggc 420
atcgcttctt ttggcaataa gataagcagc agtcccagcc ctgctgcctt atcagctgca 480
gagggtctga tctcctatgc catccagaag ggtcacctgt cgcccaggta tattcagcca 540
cttcttctga aagaagagac ctgcctggac cctcaacatc cagtgatgcc caggaagggt 600
tgccccaaca tcatcaaagc atctgcttgg gaagccagag agacacactg ccctaaaatg 660
aacctcccag ccaaatatgt catcatcatc cacaccgctg gcacaagctg cactgtatcc 720
acagactgcc agactgtcgt ccgaaacata cagtcctttc acatggacac acggaacttt 780
tgtgacattg gatatcaata aggccaggcg tggcggcgat tacgtctgta atcccaggac 840
tttgggaggg caaggcgggc agatcacttc aggccaggaa ttcaagagca gcctggccaa 900
tatggcgaaa ctctgtctct actgaaaaca aacaaacaaa caacaaagaaa 960
caacaaaaat tagccgggtg tgggtggcaca cgctgtagt cccagctact caggaggctg 1020
aggcataaga attgcttgaa ccctggaggc ggagggttga gtgagctgag attgggccac 1080
cgcaactccag tctgggagac agagtgagac tgtctcaaaa caacaacaaa aaaatcccta 1140
acataatctc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa 1182

```

<210> 2

<211> 243

<212> PRT

<213> Homo sapiens

<400> 2

```

Met Gly Thr Leu Pro Trp Leu Leu Ala Phe Phe Ile Leu Gly Leu Gln
  1              5              10              15

```

```

Ala Trp Asp Thr Pro Thr Ile Val Ser Arg Lys Glu Trp Gly Ala Arg
      20              25              30

```

```

Pro Leu Ala Cys Arg Ala Leu Leu Thr Leu Pro Val Ala Tyr Ile Ile

```

35 40 45
 Thr Asp Gln Leu Pro Gly Met Gln Cys Gln Gln Gln Ser Val Cys Ser
 50 55 60
 Gln Met Leu Arg Gly Leu Gln Ser His Ser Val Tyr Thr Ile Gly Trp
 65 70 75 80
 Cys Asp Val Ala Tyr Asn Phe Leu Val Gly Asp Asp Gly Arg Val Tyr
 85 90 95
 Glu Gly Val Gly Trp Asn Ile Gln Gly Leu His Thr Gln Gly Tyr Asn
 100 105 110
 Asn Ile Ser Leu Gly Ile Ala Phe Phe Gly Asn Lys Ile Ser Ser Ser
 115 120 125
 Pro Ser Pro Ala Ala Leu Ser Ala Ala Glu Gly Leu Ile Ser Tyr Ala
 130 135 140
 Ile Gln Lys Gly His Leu Ser Pro Arg Tyr Ile Gln Pro Leu Leu Leu
 145 150 155 160
 Lys Glu Glu Thr Cys Leu Asp Pro Gln His Pro Val Met Pro Arg Lys
 165 170 175
 Val Cys Pro Asn Ile Ile Lys Arg Ser Ala Trp Glu Ala Arg Glu Thr
 180 185 190
 His Cys Pro Lys Met Asn Leu Pro Ala Lys Tyr Val Ile Ile Ile His
 195 200 205
 Thr Ala Gly Thr Ser Cys Thr Val Ser Thr Asp Cys Gln Thr Val Val
 210 215 220
 Arg Asn Ile Gln Ser Phe His Met Asp Thr Arg Asn Phe Cys Asp Ile
 225 230 235 240
 Gly Tyr Gln

<210> 3

<211> 1876

<212> DNA

<213> Homo sapiens

<400> 3

gggcaagctg actgcaccct gacctgctgg gctgggacag cacaggaccc acagatatct 60
 gctgccatcc acactctcca gattggtgtc ctgggaccac gtggggatgc tgctgtggct 120
 tcttgtcttc tctgctctgg gtatccaggc ctgggggtgat tcctcctgga acaaaacaca 180
 agctaaacag gtatcagagg ggctccagta cctatttgag aacatctccc agctcactga 240
 aaaagatgtc tccaccacgg tctctcgcaa ggcattgggg gcagaagctg ttggctgcag 300
 tattcagctg accacgccag tgaatgtcct tgttatacac catgtccctg gactggagtg 360
 tcacgaccag acagtctgca gccagagact gcgggaactg caggcccatc atgtccacaa 420
 caacagtggg tgtgatgtgg cctacaactt cctggttggg gatgatggca ggggtgatga 480
 aggtgttggc tggaaatatcc aaggagtgca cacccaaggc tacaacaaca tctccctggg 540
 ctttgccttc ttcggcacta agaaaggcca cagtcccagc cctgctgccc tgtcggccat 600

```

ggaaaaaccta atcacctatg ctgtccagaa gggccacctg tcatccagtt atgttcagcc 660
acttcttggg aaagcgagga actgcctggc ccctcggcag aagacaagcc tgaagaagct 720
tgccccggca ttgtcccacg gtctgtgtgg ggagccaggg agaccactgt ccaggatgac 780
tctcccagcg aagtatggca tcattatcca cactgccggg aggacctgca acatttctga 840
tgagtggcgc ctgctgggtcc gggacatcca gtctttctac atagacaggc tcaagtcattg 900
cgacattggt tataacttcc tgggtgggcca ggatggcgcc atttatgaag ggggtgggctg 960
gaatgtccaa ggctcctcca cccctggcta cgatgacatt gccctgggca ttaccttcat 1020
gggcaccttc acaggtatac cacccaatgc tgcagcacta gaggcagccc aagacctgat 1080
ccagtgtgcc atgggtcaaa ggtacctgac tcccaactac ctgctggtgg gccacagtga 1140
tgtggcccga accttgtctc ctgggcaggc tttgtacaac atcatcagca cctggcctca 1200
tttcaaacac tgagagaagc cccaggctct tctgagactg ctttccctcc cctgtcaggt 1260
ctctcctgtc ctaaccatcc agcttggctc aacacctttt gccctcctcc cctgccacac 1320
agtcctgtgc ctctttttc aggttgggat gatcatgcct ctcttgccaa catcctccaa 1380
gggcctccaa actcatagct ggacattcac agccctctga gtctgagtcc agatttcttc 1440
tctccttact tcctctccct tggaaaccca actcctcagc caggtgagac aatgggctgg 1500
ttcttgtttc atttctctct ctctctccat tccctctgct ctgggtgagc ttcccttggt 1560
gtctgcctgg cagccccccac caccaccta tcaccctca cccataactc aggtcaacgt 1620
gaccaacctt ccttgccttac acataaactt gtatatattt ggatgtagcc cttatttaat 1680
ggctgtcatt atttatagat atgtctatcc ttgctacttg gttgtgagtt tctccagggg 1740
aggaactgtg ttttattcat ctctatgtcc tctgtttctc agcagtgtct gaaatttaat 1800
gggttctact gatgtttatt agagaaatgg atgaataaat gaatgaagag atccaaaaaa 1860
aaaaaaaaa aaaaaa

```

<210> 4

<211> 368

<212> PRT

<213> Homo sapiens

<400> 4

```

Met Leu Leu Trp Leu Leu Val Phe Ser Ala Leu Gly Ile Gln Ala Trp
  1              5              10              15

Gly Asp Ser Ser Trp Asn Lys Thr Gln Ala Lys Gln Val Ser Glu Gly
      20              25              30

Leu Gln Tyr Leu Phe Glu Asn Ile Ser Gln Leu Thr Glu Lys Asp Val
      35              40              45

Ser Thr Thr Val Ser Arg Lys Ala Trp Gly Ala Glu Ala Val Gly Cys
      50              55              60

Ser Ile Gln Leu Thr Thr Pro Val Asn Val Leu Val Ile His His Val
      65              70              75              80

Pro Gly Leu Glu Cys His Asp Gln Thr Val Cys Ser Gln Arg Leu Arg
      85              90              95

Glu Leu Gln Ala His His Val His Asn Asn Ser Gly Cys Asp Val Ala
      100              105              110

Tyr Asn Phe Leu Val Gly Asp Asp Gly Arg Val Tyr Glu Gly Val Gly
      115              120              125

Trp Asn Ile Gln Gly Val His Thr Gln Gly Tyr Asn Asn Ile Ser Leu
      130              135              140

Gly Phe Ala Phe Phe Gly Thr Lys Lys Gly His Ser Pro Ser Pro Ala
      145              150              155              160

```

Ala Leu Ser Ala Met Glu Asn Leu Ile Thr Tyr Ala Val Gln Lys Gly
 165 170 175

His Leu Ser Ser Ser Tyr Val Gln Pro Leu Leu Gly Lys Gly Glu Asn
 180 185 190

Cys Leu Ala Pro Arg Gln Lys Thr Ser Leu Lys Lys Leu Ala Pro Ala
 195 200 205

Leu Ser His Gly Leu Cys Gly Glu Pro Gly Arg Pro Leu Ser Arg Met
 210 215 220

Thr Leu Pro Ala Lys Tyr Gly Ile Ile Ile His Thr Ala Gly Arg Thr
 225 230 235 240

Cys Asn Ile Ser Asp Glu Cys Arg Leu Leu Val Arg Asp Ile Gln Ser
 245 250 255

Phe Tyr Ile Asp Arg Leu Lys Ser Cys Asp Ile Gly Tyr Asn Phe Leu
 260 265 270

Val Gly Gln Asp Gly Ala Ile Tyr Glu Gly Val Gly Trp Asn Val Gln
 275 280 285

Gly Ser Ser Thr Pro Gly Tyr Asp Asp Ile Ala Leu Gly Ile Thr Phe
 290 295 300

Met Gly Thr Phe Thr Gly Ile Pro Pro Asn Ala Ala Ala Leu Glu Ala
 305 310 315 320

Ala Gln Asp Leu Ile Gln Cys Ala Met Val Lys Gly Tyr Leu Thr Pro
 325 330 335

Asn Tyr Leu Leu Val Gly His Ser Asp Val Ala Arg Thr Leu Ser Pro
 340 345 350

Gly Gln Ala Leu Tyr Asn Ile Ile Ser Thr Trp Pro His Phe Lys His
 355 360 365

<210> 5

<211> 749

<212> DNA

<213> Homo sapiens

<400> 5

gatcccccg gctgcaggaa ttcggcacga gccggaccct gccgccctgc cactatgtcc 60
 cgccgctcta tgctgcttgc ctgggctctc ccagcctcc ttcgactcgg agcggctcag 120
 gagacagaag acccggcctg ctgcagcccc atagtgcgcc ggaacgagtg gaaggccctg 180
 gcatcagagt gcgcccagca cctgagcctg cccttacgct atgtggtggt atcgcacacg 240
 gcgggcagca gctgcaacac cccgcctcg tgccagcagc aggcccgaa tgtgcagcac 300
 taccacatga agacactggg ctgggtgcgac gtgggctaca acttcctgat tggagaagac 360
 gggctcgtat acgagggccg tggtggaac ttcacgggtg ccactcagg tcacttatgg 420
 aaccccatgt ccattggcat cagcttcctg ggcaactaca tggatcgggt gccacaccc 480

5

caggccatcc gggcagccca gggctctactg gcctgcggtg tggctcaggg agccctgagg 540
 tccaactatg tgctcaaagg acaccgggat gtgcagcgta cactctctcc aggcaaccag 600
 ctctaccacc tcatccagaa ttggccacac taccgctccc cctgaggccc tgctgatccg 660
 caccgccattc ctcccctccc atggccaaaa accccactgt ctcttctcc aataaagatg 720
 tagctcaaaa aaaaaaaaaa aaaaaaaaaa 749

<210> 6

<211> 196

<212> PRT

<213> Homo sapiens

<400> 6

Met	Ser	Arg	Arg	Ser	Met	Leu	Leu	Ala	Trp	Ala	Leu	Pro	Ser	Leu	Leu
1				5					10					15	
Arg	Leu	Gly	Ala	Ala	Gln	Glu	Thr	Glu	Asp	Pro	Ala	Cys	Cys	Ser	Pro
			20					25					30		
Ile	Val	Pro	Arg	Asn	Glu	Trp	Lys	Ala	Leu	Ala	Ser	Glu	Cys	Ala	Gln
		35					40					45			
His	Leu	Ser	Leu	Pro	Leu	Arg	Tyr	Val	Val	Val	Ser	His	Thr	Ala	Gly
	50					55					60				
Ser	Ser	Cys	Asn	Thr	Pro	Ala	Ser	Cys	Gln	Gln	Gln	Ala	Arg	Asn	Val
	65				70				75					80	
Gln	His	Tyr	His	Met	Lys	Thr	Leu	Gly	Trp	Cys	Asp	Val	Gly	Tyr	Asn
				85					90					95	
Phe	Leu	Ile	Gly	Glu	Asp	Gly	Leu	Val	Tyr	Glu	Gly	Arg	Gly	Trp	Asn
			100				105						110		
Phe	Thr	Gly	Ala	His	Ser	Gly	His	Leu	Trp	Asn	Pro	Met	Ser	Ile	Gly
		115					120					125			
Ile	Ser	Phe	Met	Gly	Asn	Tyr	Met	Asp	Arg	Val	Pro	Thr	Pro	Gln	Ala
	130				135						140				
Ile	Arg	Ala	Ala	Gln	Gly	Leu	Leu	Ala	Cys	Gly	Val	Ala	Gln	Gly	Ala
145				150					155					160	
Leu	Arg	Ser	Asn	Tyr	Val	Leu	Lys	Gly	His	Arg	Asp	Val	Gln	Arg	Thr
			165					170					175		
Leu	Ser	Pro	Gly	Asn	Gln	Leu	Tyr	His	Leu	Ile	Gln	Asn	Trp	Pro	His
		180					185					190			
Tyr	Arg	Ser	Pro												
		195													

<210> 7

<211> 151

<212> PRT

<213> Homo sapiens

<400> 7

6

Met Leu Phe Ala Cys Ala Leu Leu Ala Leu Leu Gly Leu Ala Thr Ser
 1 5 10 15
 Cys Ser Phe Ile Val Pro Arg Ser Glu Trp Arg Ala Leu Pro Ser Glu
 20 25 30
 Cys Ser Ser Arg Leu Gly His Pro Val Arg Tyr Val Val Ile Ser His
 35 40 45
 Thr Ala Gly Ser Phe Cys Asn Ser Pro Asp Ser Cys Glu Gln Gln Ala
 50 55 60
 Arg Asn Val Gln His Tyr His Lys Asn Glu Leu Gly Trp Cys Asp Val
 65 70 75 80
 Ala Tyr Asn Phe Leu Ile Gly Glu Asp Gly His Val Tyr Glu Gly Arg
 85 90 95
 Gly Trp Asn Ile Lys Gly Asp His Thr Gly Pro Ile Trp Asn Pro Met
 100 105 110
 Ser Ile Gly Ile Thr Phe Met Gly Asn Phe Met Asp Arg Val Arg Lys
 115 120 125
 Ala Gly Pro Pro Cys Cys Pro Lys Ser Ser Gly Ile Trp Gly Val Ser
 130 135 140
 Gly Leu Pro Glu Ile Gln Leu
 145 150

<210> 8
 <211> 43
 <212> DNA
 <213> Homo sapiens

<400> 8
 gcagcacata tggattcctc ctggaacaaa acacaagcta aac 43

<210> 9
 <211> 46
 <212> DNA
 <213> Homo sapiens

<400> 9
 gcagcaggta ccttagtggt tgaaatgagg ccagggtgctg atgatg 46

<210> 10
 <211> 48
 <212> DNA
 <213> Homo sapiens

<400> 10
 gcagcaggat ccgccatcat ggggacgctg ccatggcttc ttgccttc 48

<210> 11
 <211> 46
 <212> DNA

<213> Homo sapiens

<400> 11
gcagcaggta ccttattgat atccaatgtc acaaaaagttc cgtgtg 46

<210> 12
<211> 48
<212> DNA
<213> Homo sapiens

<400> 12
gcagcaggat ccgccatcat ggggacgctg ccatggcttc ttgccttc 48

<210> 13
<211> 46
<212> DNA
<213> Homo sapiens

<400> 13
gcagcaggta ccttattgat atccaatgtc acaaaaagttc cgtgtg 46

<210> 14
<211> 733
<212> DNA
<213> Homo sapiens

<400> 14
gggatccgga gcccaaatct tctgacaaaa ctcacacatg cccaccgtgc ccagcacctg 60
aattcgaggg tgcaccgtca gtcttctctt tcccccaaa acccaaggac accctcatga 120
tctcccggac tcttgaggtc acatgcgtgg tgggtggacgt aagccacgaa gacctgagg 180
tcaagttcaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccgcggg 240
aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg caccaggact 300
ggctgaatgg caaggagtac aagtgaagg tctccaacaa agccctccca acccccatcg 360
agaaaaccat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac accctgcccc 420
catcccggga tgagctgacc aagaaccagg tcagcctgac ctgcctggtc aaaggcttct 480
atccaagcga catcgccgtg gagtgggaga gcaatgggca gccggagaac aactacaaga 540
ccacgcctcc cgtgctggac tccgacggct ccttcttctt ctacagcaag ctcaccgtgg 600
acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcat gaggctctgc 660
acaaccacta cagcagaag agcctctccc tgtctccggg taaatgagtg cgacggccgc 720
gactctagag gat 733

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/30736

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12Q 1/100; G01N 33/53, 33/48; C12P 21/06; C12N 15/00; C07H 21/02; C07K 16/00, 1/00; A61K 39/00
US CL : 435/4, 69.1, 7.1, 320.1; 536/23.1; 530,387.1, 350+; 424/185.1; 436/63.64

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/4, 69.1, 7.1, 320.1; 536/23.1; 530,387.1, 350+; 424/185.1; 436/63.64

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
STN, WEST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/02686 A1 (BOEHRINGER INGELHEIM INTERNATIONAL) 11 July 1997 (11.07.97) entire document.	1-23
X	KANG et al. A peptidoglycan recognition protein in innate immunity conserved from insects to humans. Proc. Natl. Acad. Sci. USA. August 1998, Vol. 95, pages 10078-10082, entire document.	1-16, 20-23
Y,E	US 6,034,217 A (ASHIDA et al.) 07 March 2000 (12.10.1997) column 1, lines 40-43.	17-19

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 16 March 2000 (16.03.2000)	Date of mailing of the international search report 11 APR 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>Gary Nickol</i> Gary Nickol Telephone No. 703-308-0196

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)